



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:37 PM UTC

PDB ID : 2FJ1 / pdb_00002fj1
Title : Crystal Structure Analysis of Tet Repressor (class D) in Complex with 7-Chl
ortetracycline-Nickel(II)
Authors : Orth, P.; Saenger, W.; Palm, G.J.; Hinrichs, W.
Deposited on : 2005-12-30
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 1819 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tetracycline repressor protein class D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	0	4	0
			1680	1055	303	319	3			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	SER	ALA	engineered mutation	UNP P0ACT4


- Molecule 2 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ni	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Cl	0	0
			3	3		

- Molecule 4 is 7-CHLOROTETRACYCLINE (CCD ID: CTC) (formula: C₂₂H₂₃ClN₂O₈).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 33	C 22	Cl 1	N 2	O 8	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	102	Total O 102 102	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tetracycline repressor protein class D



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	68.94Å 68.94Å 180.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.55 – 2.20 64.42 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.6 (64.55-2.20) 97.6 (64.42-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.146 , 0.215 0.156 , 0.221	Depositor DCC
R_{free} test set	544 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	30.0	Xtriage
Anisotropy	0.322	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1819	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NI, CTC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.89	22/1710 (1.3%)	1.47	8/2314 (0.3%)

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	177	MET	SD-CE	20.31	2.30	1.79
1	A	136	ALA	CA-CB	-7.13	1.41	1.53
1	A	118	ARG	NE-CZ	6.91	1.40	1.33
1	A	84	MET	SD-CE	-6.53	1.63	1.79
1	A	34	LEU	CA-C	6.49	1.61	1.52
1	A	88	ARG	NE-CZ	6.24	1.40	1.33
1	A	64	HIS	C-O	-6.21	1.16	1.23
1	A	13	ALA	CA-CB	6.04	1.62	1.53
1	A	184	GLN	CD-OE1	5.97	1.34	1.23
1	A	92	ARG	C-O	-5.49	1.17	1.24
1	A	123	ASN	C-O	-5.36	1.16	1.24
1	A	6	ARG	CA-C	5.31	1.59	1.52
1	A	98	LYS	C-O	-5.30	1.17	1.24
1	A	133	ALA	CA-CB	-5.29	1.45	1.53
1	A	200	GLN	C-O	-5.29	1.18	1.24
1	A	8	SER	CA-C	-5.28	1.46	1.52
1	A	6	ARG	C-O	5.23	1.30	1.24
1	A	192	SER	CA-C	5.17	1.59	1.52
1	A	136	ALA	C-O	-5.17	1.17	1.24
1	A	131	LEU	N-CA	-5.16	1.40	1.46
1	A	168	PRO	C-O	-5.08	1.17	1.24
1	A	118	ARG	CD-NE	5.06	1.53	1.46

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	139	HIS	N-CA-C	6.30	117.81	111.07
1	A	167	PRO	CB-CA-C	6.26	117.06	111.17
1	A	134	ILE	N-CA-C	6.14	116.81	110.36
1	A	94	ARG	CB-CG-CD	5.85	124.76	111.30
1	A	195	ARG	NE-CZ-NH2	5.25	123.93	119.20
1	A	11	ASP	CB-CA-C	-5.13	102.60	110.81
1	A	204	LEU	CB-CA-C	-5.11	110.28	117.23
1	A	207	ILE	N-CA-CB	5.04	116.03	110.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1680	0	1663	20	0
2	A	1	0	0	0	0
3	A	3	0	0	0	0
4	A	33	0	20	1	0
5	A	102	0	0	9	0
All	All	1819	0	1683	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:MET:SD	1:A:177:MET:CE	2.30	1.20
1:A:65[B]:ASP:OD2	5:A:374:HOH:O	2.04	0.74
4:A:222:CTC:H6'1	4:A:222:CTC:CL7	2.38	0.60
1:A:63:HIS:CD2	1:A:92:ARG:HD3	2.37	0.59
1:A:63:HIS:CD2	1:A:92:ARG:CD	2.87	0.58
1:A:49:ARG:NH2	5:A:371:HOH:O	2.17	0.57
1:A:148[A]:GLN:NE2	5:A:363:HOH:O	2.37	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:ARG:HD3	5:A:339:HOH:O	2.04	0.56
1:A:58:GLU:HG2	5:A:398:HOH:O	2.05	0.55
1:A:155:LEU:O	1:A:155:LEU:HD23	2.07	0.54
1:A:21:GLY:CA	1:A:94:ARG:HD2	2.39	0.52
1:A:151:HIS:HE1	5:A:385:HOH:O	1.94	0.51
1:A:197[B]:PHE:CE2	5:A:393:HOH:O	2.65	0.49
1:A:128[B]:ARG:NE	5:A:342:HOH:O	2.47	0.47
1:A:63:HIS:CD2	1:A:92:ARG:HD2	2.51	0.46
1:A:63:HIS:NE2	1:A:92:ARG:HD3	2.31	0.46
1:A:21:GLY:HA2	1:A:94:ARG:HD2	2.00	0.43
1:A:155:LEU:O	1:A:155:LEU:CG	2.67	0.42
1:A:128[B]:ARG:HD3	5:A:366:HOH:O	2.20	0.41
1:A:158:ARG:HA	1:A:159:PRO:HD3	1.80	0.41
1:A:163:ASP:HA	1:A:166:LEU:HD22	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	209/207 (101%)	199 (95%)	8 (4%)	2 (1%)	12 11

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	160	ALA
1	A	162	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/173 (102%)	162 (92%)	15 (8%)	10	11

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	A	14	LEU
1	A	27	THR
1	A	29	LYS
1	A	55	LEU
1	A	79	LEU
1	A	113	VAL
1	A	127	LEU
1	A	142	LEU
1	A	155	LEU
1	A	164	GLU
1	A	166	LEU
1	A	177	MET
1	A	205	LEU
1	A	206	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	HIS
1	A	151	HIS
1	A	200	GLN
1	A	206	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	CTC	A	222	2	35,36,36	2.14	11 (31%)	42,60,60	2.12	13 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CTC	A	222	2	-	0/8/74/74	0/4/4/4

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	222	CTC	O11-C11	5.87	1.35	1.23
4	A	222	CTC	C6-C6A	-4.51	1.48	1.53
4	A	222	CTC	C5B-C12	3.66	1.40	1.36
4	A	222	CTC	C5B-C11	-3.59	1.38	1.46
4	A	222	CTC	O4B-C4B	3.38	1.47	1.42
4	A	222	CTC	C4A-C4	3.36	1.57	1.54
4	A	222	CTC	O1-C1	3.34	1.28	1.22
4	A	222	CTC	O10-C10	-2.65	1.31	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	222	CTC	C4-C3	-2.59	1.46	1.51
4	A	222	CTC	C4B-C4A	-2.59	1.51	1.53
4	A	222	CTC	C5-C4A	-2.07	1.50	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	222	CTC	O11-C11-C6B	-5.45	112.20	122.06
4	A	222	CTC	C11-C5B-C12	5.32	123.01	118.80
4	A	222	CTC	C4B-C12-C5B	-3.54	119.47	123.06
4	A	222	CTC	C4'-N4-C4	-3.35	106.47	114.10
4	A	222	CTC	C8-C7-CL7	-3.26	112.00	118.42
4	A	222	CTC	C6'-C6-C6A	-3.14	104.78	111.64
4	A	222	CTC	O6-C6-C6'	-3.09	100.34	108.15
4	A	222	CTC	O6-C6-C6A	3.07	115.31	108.84
4	A	222	CTC	C4A-C4B-C1	-2.99	107.62	111.05
4	A	222	CTC	C5-C4A-C4	-2.79	109.69	113.73
4	A	222	CTC	O2'-C2'-N2'	-2.39	117.54	122.89
4	A	222	CTC	O4B-C4B-C12	-2.20	106.62	110.14
4	A	222	CTC	C10-C6B-C11	-2.01	118.47	121.45

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	222	CTC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/207 (100%)	-0.50	2 (0%) 79 77	9, 32, 71, 112	4 (1%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	162	PRO	2.2
1	A	161	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CL	A	303	1/1	0.87	0.14	63,63,63,63	1
3	CL	A	302	1/1	0.97	0.05	42,42,42,42	0
4	CTC	A	222	33/33	0.97	0.05	18,23,31,32	0
3	CL	A	301	1/1	0.99	0.04	33,33,33,33	0
2	NI	A	223	1/1	1.00	0.01	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.