



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:37 AM UTC

PDB ID : 6FKV / pdb_00006fkv
Title : Structure and function of aldehyde dehydrogenase from *Thermus thermophilus*: An enzyme with an evolutionarily-distinct C-terminal arm (Recombinant protein with shortened C-terminal, ADH508)
Authors : Hayes, K.A.; Noor, M.R.; Djeghader, A.; Soulimane, T.
Deposited on : 2018-01-24
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	517	
1	B	517	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7994 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aldehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	504	Total	C	N	O	S	0	0	0
			3946	2530	692	717	7			
1	B	505	Total	C	N	O	S	0	0	0
			3954	2536	693	718	7			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	expression tag	UNP Q72KD3
A	-4	HIS	-	expression tag	UNP Q72KD3
A	-3	HIS	-	expression tag	UNP Q72KD3
A	-2	HIS	-	expression tag	UNP Q72KD3
A	-1	HIS	-	expression tag	UNP Q72KD3
A	0	HIS	-	expression tag	UNP Q72KD3
A	1	HIS	-	expression tag	UNP Q72KD3
B	-5	MET	-	expression tag	UNP Q72KD3
B	-4	HIS	-	expression tag	UNP Q72KD3
B	-3	HIS	-	expression tag	UNP Q72KD3
B	-2	HIS	-	expression tag	UNP Q72KD3
B	-1	HIS	-	expression tag	UNP Q72KD3
B	0	HIS	-	expression tag	UNP Q72KD3
B	1	HIS	-	expression tag	UNP Q72KD3

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		
2	A	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	4	Total Cl 4 4	0	0

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C N O 8 4 1 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	7	Total 7	O 7	0	0
6	B	10	Total 10	O 10	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aldehyde dehydrogenase



- Molecule 1: Aldehyde dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	105.19Å 105.19Å 314.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.04 – 2.90 48.04 – 2.90	Depositor EDS
% Data completeness (in resolution range)	97.7 (48.04-2.90) 97.7 (48.04-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.190 , 0.229 0.201 , 0.234	Depositor DCC
R_{free} test set	2000 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	57.7	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7994	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, SO4, TRS, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/4049	1.09	9/5500 (0.2%)
1	B	0.82	0/4057	1.07	4/5511 (0.1%)
All	All	0.82	0/8106	1.08	13/11011 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	10
1	B	0	13
All	All	0	23

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	380	ASP	CA-C-N	7.25	127.42	119.87
1	B	380	ASP	C-N-CA	7.25	127.42	119.87
1	B	204	PRO	N-CA-CB	7.24	107.25	103.19
1	A	380	ASP	CA-C-N	7.13	127.29	119.87
1	A	380	ASP	C-N-CA	7.13	127.29	119.87
1	A	204	PRO	N-CA-CB	6.54	106.85	103.19
1	A	470	PRO	N-CA-CB	6.09	108.17	103.30
1	A	267	PRO	N-CA-CB	6.05	108.68	103.36
1	B	342	ARG	CB-CA-C	-5.39	102.18	110.81
1	A	178	THR	CB-CA-C	5.22	119.79	109.66
1	A	266	ASN	CB-CA-C	5.10	115.94	110.08
1	A	182	LYS	CB-CA-C	5.09	115.52	110.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	267	PRO	CB-CA-C	-5.03	105.38	111.46

There are no chirality outliers.

All (23) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	126	ARG	Sidechain
1	A	146	ARG	Sidechain
1	A	245	ARG	Sidechain
1	A	272	ARG	Sidechain
1	A	352	ARG	Sidechain
1	A	376	ARG	Sidechain
1	A	399	ARG	Sidechain
1	A	417	GLY	Peptide
1	A	68	ARG	Sidechain
1	A	74	ARG	Sidechain
1	B	127	ARG	Sidechain
1	B	146	ARG	Sidechain
1	B	245	ARG	Sidechain
1	B	257	ARG	Sidechain
1	B	32	ARG	Sidechain
1	B	360	ARG	Sidechain
1	B	365	ARG	Sidechain
1	B	370	ARG	Sidechain
1	B	399	ARG	Sidechain
1	B	417	GLY	Peptide
1	B	452	ARG	Sidechain
1	B	68	ARG	Sidechain
1	B	94	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3946	0	3896	46	0
1	B	3954	0	3907	34	0
2	A	14	0	20	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	21	0	30	3	0
3	A	20	0	0	2	0
3	B	10	0	0	0	0
4	A	4	0	0	0	0
5	B	8	0	12	2	0
6	A	7	0	0	0	0
6	B	10	0	0	0	0
All	All	7994	0	7865	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:277:ASP:OD1	1:A:314:ARG:NH1	1.97	0.97
1:B:277:ASP:OD1	1:B:314:ARG:NH1	2.00	0.94
1:A:59:ALA:HB1	1:A:178:THR:HG23	1.51	0.91
1:A:59:ALA:O	1:A:178:THR:HG21	1.87	0.74
1:A:261:GLU:OE2	1:A:483:GLU:HG3	1.94	0.68
1:A:501:GLU:OE1	1:A:507:GLN:NE2	2.27	0.67
1:B:261:GLU:OE2	1:B:483:GLU:HG3	1.95	0.66
1:A:95:LEU:HD23	1:A:190:LEU:HD23	1.80	0.63
1:B:294:ARG:HH12	5:B:704:TRS:H11	1.69	0.57
1:A:59:ALA:HB1	1:A:178:THR:CG2	2.29	0.57
1:B:294:ARG:HH22	5:B:704:TRS:H12	1.68	0.56
1:A:414:LYS:HD3	2:A:601:PEG:C1	2.35	0.56
1:B:95:LEU:HD23	1:B:190:LEU:HD23	1.86	0.56
1:A:79:PHE:CE2	1:A:83:LYS:HE2	2.42	0.55
1:B:198:PHE:CD1	1:B:203:LEU:HD22	2.42	0.55
1:B:474:VAL:HG23	2:B:701:PEG:H11	1.89	0.55
1:B:79:PHE:CE2	1:B:83:LYS:HE2	2.41	0.54
1:A:257:ARG:NH2	1:B:491:GLU:OE1	2.39	0.54
1:B:351:TYR:CE2	1:B:366:GLY:HA2	2.45	0.52
1:A:149:LEU:O	2:B:701:PEG:H41	2.10	0.52
1:A:198:PHE:CD1	1:A:203:LEU:HD22	2.45	0.52
1:B:7:LYS:HE2	1:B:36:ASP:OD2	2.11	0.51
1:A:294:ARG:NH1	3:A:604:SO4:O3	2.43	0.51
1:B:7:LYS:CE	1:B:36:ASP:OD2	2.59	0.51
1:B:7:LYS:NZ	1:B:36:ASP:OD2	2.43	0.51
1:A:16:HIS:CD2	1:A:51:LEU:CD2	2.95	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:414:LYS:HD3	2:A:601:PEG:H12	1.93	0.50
1:B:237:PHE:CZ	1:B:239:GLY:HA3	2.47	0.50
1:A:237:PHE:CZ	1:A:239:GLY:HA3	2.47	0.49
1:B:305:ALA:HB3	1:B:306:PRO:HD3	1.95	0.49
1:B:266:ASN:ND2	1:B:296:THR:HA	2.28	0.49
1:B:29:LEU:O	1:B:43:ARG:HG2	2.12	0.48
1:A:253:ARG:HB3	1:B:249:GLU:HB2	1.95	0.48
1:A:414:LYS:HD3	2:A:601:PEG:H11	1.95	0.48
1:A:41:VAL:HG13	1:A:98:ARG:NH1	2.29	0.47
1:A:95:LEU:HD21	1:A:189:THR:HG22	1.96	0.47
1:A:13:GLU:HA	1:A:43:ARG:O	2.14	0.47
1:B:16:HIS:CD2	1:B:51:LEU:CD2	2.96	0.47
1:B:41:VAL:HG13	1:B:98:ARG:NH1	2.28	0.47
1:B:16:HIS:CD2	1:B:51:LEU:HD21	2.49	0.47
1:A:16:HIS:CD2	1:A:51:LEU:HD21	2.50	0.47
1:B:13:GLU:HA	1:B:43:ARG:O	2.15	0.47
1:B:95:LEU:HD21	1:B:189:THR:HG22	1.97	0.46
1:B:63:PHE:CE2	1:B:150:GLY:HA2	2.51	0.46
1:A:457:SER:HB3	1:A:460:ASN:HB3	1.98	0.46
1:A:27:PRO:HG2	1:A:46:GLU:HB3	1.98	0.46
1:A:351:TYR:CE2	1:A:366:GLY:HA2	2.51	0.46
1:A:266:ASN:ND2	1:A:296:THR:HA	2.31	0.44
1:A:53:ARG:NH2	3:A:606:SO4:O4	2.39	0.44
1:A:325:GLY:N	1:A:333:VAL:HG21	2.33	0.44
1:B:219:THR:HA	1:B:222:TRP:CD1	2.52	0.43
1:A:448:LYS:HD2	1:B:498:VAL:HG11	2.00	0.43
1:B:325:GLY:N	1:B:333:VAL:HG21	2.33	0.43
1:A:71:ALA:N	1:A:72:PRO:CD	2.81	0.43
1:B:467:ALA:O	1:B:484:SER:HB3	2.18	0.43
1:A:178:THR:OG1	1:A:207:VAL:HA	2.19	0.42
1:A:215:GLY:HA3	2:A:602:PEG:H22	2.01	0.42
1:B:44:PHE:CE1	1:B:188:PRO:HG3	2.53	0.42
1:A:146:ARG:HH11	1:A:146:ARG:HG2	1.84	0.42
1:B:139:ASP:CB	1:B:503:SER:HB3	2.50	0.42
1:A:264:GLY:HA2	1:A:430:TYR:CG	2.55	0.42
1:A:149:LEU:O	2:B:701:PEG:H22	2.19	0.42
1:B:264:GLY:HA2	1:B:430:TYR:CG	2.55	0.42
1:B:457:SER:HB3	1:B:460:ASN:HB3	2.02	0.42
1:A:313:ARG:NH1	1:A:317:GLU:OE1	2.52	0.42
1:A:219:THR:HA	1:A:222:TRP:CD1	2.54	0.41
1:A:474:VAL:HG22	1:A:475:LYS:N	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:LYS:HE3	1:A:183:PRO:O	2.20	0.41
1:A:305:ALA:HB3	1:A:306:PRO:HD3	2.03	0.41
1:A:467:ALA:O	1:A:484:SER:HB3	2.20	0.41
1:A:59:ALA:C	1:A:178:THR:HG21	2.43	0.41
1:A:63:PHE:CE2	1:A:150:GLY:HA2	2.55	0.41
1:B:16:HIS:CE1	1:B:45:PRO:HG2	2.55	0.41
1:A:311:PHE:CD1	1:A:311:PHE:C	2.98	0.41
1:B:71:ALA:N	1:B:72:PRO:CD	2.83	0.41
1:A:16:HIS:CE1	1:A:45:PRO:HG2	2.56	0.41
1:A:506:LEU:O	1:A:507:GLN:C	2.64	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	502/517 (97%)	481 (96%)	21 (4%)	0	100	100
1	B	503/517 (97%)	484 (96%)	19 (4%)	0	100	100
All	All	1005/1034 (97%)	965 (96%)	40 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	402/414 (97%)	378 (94%)	24 (6%)	17	47
1	B	403/414 (97%)	381 (94%)	22 (6%)	19	50
All	All	805/828 (97%)	759 (94%)	46 (6%)	18	49

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	GLU
1	A	28	LEU
1	A	35	SER
1	A	38	GLU
1	A	61	GLU
1	A	105	LYS
1	A	178	THR
1	A	182	LYS
1	A	199	GLU
1	A	209	ASN
1	A	295	CYS
1	A	306	PRO
1	A	310	GLU
1	A	312	LYS
1	A	314	ARG
1	A	317	GLU
1	A	333	VAL
1	A	349	GLU
1	A	350	HIS
1	A	400	LEU
1	A	404	GLU
1	A	420	GLU
1	A	427	SER
1	A	452	ARG
1	B	13	GLU
1	B	25	GLU
1	B	35	SER
1	B	61	GLU
1	B	91	THR
1	B	105	LYS
1	B	209	ASN
1	B	216	LYS
1	B	295	CYS
1	B	306	PRO
1	B	310	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	312	LYS
1	B	314	ARG
1	B	333	VAL
1	B	349	GLU
1	B	350	HIS
1	B	371	GLU
1	B	395	ARG
1	B	400	LEU
1	B	404	GLU
1	B	420	GLU
1	B	427	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	GLN
1	A	266	ASN
1	A	411	ASN
1	B	76	GLN
1	B	266	ASN
1	B	372	ASN
1	B	411	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PEG	A	601	-	6,6,6	0.81	0	5,5,5	0.87	0
3	SO4	A	605	-	4,4,4	0.59	0	6,6,6	0.42	0
3	SO4	A	606	-	4,4,4	0.56	0	6,6,6	0.62	0
3	SO4	B	706	-	4,4,4	0.40	0	6,6,6	1.06	0
2	PEG	B	701	-	6,6,6	1.10	0	5,5,5	1.84	1 (20%)
2	PEG	A	602	-	6,6,6	0.70	0	5,5,5	0.94	0
3	SO4	A	603	-	4,4,4	0.77	0	6,6,6	0.74	0
3	SO4	B	705	-	4,4,4	0.50	0	6,6,6	0.55	0
3	SO4	A	604	-	4,4,4	0.57	0	6,6,6	0.83	0
5	TRS	B	704	-	7,7,7	0.50	0	9,9,9	1.11	0
2	PEG	B	702	-	6,6,6	0.62	0	5,5,5	0.53	0
2	PEG	B	703	-	6,6,6	0.73	0	5,5,5	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	A	601	-	-	1/4/4/4	-
2	PEG	B	701	-	-	2/4/4/4	-
2	PEG	A	602	-	-	1/4/4/4	-
5	TRS	B	704	-	-	3/9/9/9	-
2	PEG	B	702	-	-	1/4/4/4	-
2	PEG	B	703	-	-	3/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	PEG	O2-C2-C1	3.34	124.82	110.11

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	701	PEG	C4-C3-O2-C2
2	A	602	PEG	C1-C2-O2-C3
2	A	601	PEG	O2-C3-C4-O4
5	B	704	TRS	C1-C-C3-O3
2	B	703	PEG	O2-C3-C4-O4
2	B	702	PEG	O1-C1-C2-O2
2	B	703	PEG	O1-C1-C2-O2
5	B	704	TRS	C2-C-C3-O3
5	B	704	TRS	N-C-C3-O3
2	B	701	PEG	O1-C1-C2-O2
2	B	703	PEG	C1-C2-O2-C3

There are no ring outliers.

6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	PEG	3	0
3	A	606	SO4	1	0
2	B	701	PEG	3	0
2	A	602	PEG	1	0
3	A	604	SO4	1	0
5	B	704	TRS	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	504/517 (97%)	-0.18	5 (0%)	79 73	37, 54, 76, 113	0
1	B	505/517 (97%)	-0.14	3 (0%)	85 81	38, 55, 79, 97	0
All	All	1009/1034 (97%)	-0.16	8 (0%)	82 77	37, 55, 78, 113	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	ALA	3.4
1	A	4	ALA	3.2
1	A	24	LEU	2.7
1	B	508	LEU	2.6
1	B	440	HIS	2.5
1	A	506	LEU	2.5
1	A	483	GLU	2.2
1	B	352	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PEG	B	703	7/7	0.81	0.21	54,66,72,74	0
5	TRS	B	704	8/8	0.86	0.16	62,89,93,106	0
2	PEG	B	701	7/7	0.87	0.18	61,72,77,81	0
2	PEG	A	601	7/7	0.88	0.16	57,62,70,70	0
2	PEG	B	702	7/7	0.89	0.17	56,67,69,69	0
3	SO4	A	606	5/5	0.92	0.10	77,96,101,107	0
4	CL	A	610	1/1	0.93	0.08	66,66,66,66	0
3	SO4	B	706	5/5	0.93	0.15	78,80,85,85	0
2	PEG	A	602	7/7	0.94	0.13	53,68,76,76	0
3	SO4	A	605	5/5	0.94	0.12	84,101,109,118	0
4	CL	A	609	1/1	0.95	0.08	67,67,67,67	0
4	CL	A	607	1/1	0.95	0.10	63,63,63,63	0
4	CL	A	608	1/1	0.95	0.07	60,60,60,60	0
3	SO4	A	604	5/5	0.96	0.13	66,68,86,93	0
3	SO4	A	603	5/5	0.97	0.06	55,64,71,72	0
3	SO4	B	705	5/5	0.97	0.15	60,68,73,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.