



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:35 AM UTC

PDB ID : 4FL6 / pdb_00004fl6
Title : Crystal structure of the complex of the 3-MBT repeat domain of L3MBTL3 and UNC1215
Authors : Zhong, N.; Tempel, W.; Ravichandran, M.; Dong, A.; Ingerman, L.A.; Graslund, S.; Frye, S.V.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.M.; Brown, P.J.; Structural Genomics Consortium (SGC)
Deposited on : 2012-06-14
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	5% 85% 8% 7%
1	B	332	8% 83% 9% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	UNX	A	1005	-	-	-	X
3	UNX	A	1026	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4804 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lethal(3)malignant brain tumor-like protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	309	Total	C	N	O	S	0	0	0
			2394	1555	400	424	15			
1	B	306	Total	C	N	O	S	0	0	1
			2303	1494	386	408	15			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	228	MET	-	expression tag	UNP Q96JM7
A	554	HIS	-	expression tag	UNP Q96JM7
A	555	HIS	-	expression tag	UNP Q96JM7
A	556	HIS	-	expression tag	UNP Q96JM7
A	557	HIS	-	expression tag	UNP Q96JM7
A	558	HIS	-	expression tag	UNP Q96JM7
A	559	HIS	-	expression tag	UNP Q96JM7
B	228	MET	-	expression tag	UNP Q96JM7
B	554	HIS	-	expression tag	UNP Q96JM7
B	555	HIS	-	expression tag	UNP Q96JM7
B	556	HIS	-	expression tag	UNP Q96JM7
B	557	HIS	-	expression tag	UNP Q96JM7
B	558	HIS	-	expression tag	UNP Q96JM7
B	559	HIS	-	expression tag	UNP Q96JM7

- Molecule 2 is [2-(phenylamino)benzene-1,4-diyl]bis{[4-(pyrrolidin-1-yl)piperidin-1-yl]methanone} (CCD ID: UWN) (formula: C₃₂H₄₃N₅O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 39	C 32	N 5	O 2	0	0
2	A	1	Total 39	C 32	N 5	O 2	0	0


- Molecule 3 is UNKNOWN ATOM OR ION (CCD ID: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	24	Total X 24 24	0	0
3	B	5	Total X 5 5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lethal(3)malignant brain tumor-like protein 3



- Molecule 1: Lethal(3)malignant brain tumor-like protein 3



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	228.03Å 54.37Å 65.55Å 90.00° 97.11° 90.00°	Depositor
Resolution (Å)	44.11 – 2.55 44.11 – 2.55	Depositor EDS
% Data completeness (in resolution range)	98.9 (44.11-2.55) 98.9 (44.11-2.55)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 2.54Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.10.0, BUSTER 2.10.0	Depositor
R, R_{free}	0.195 , 0.259 0.211 , 0.267	Depositor DCC
R_{free} test set	1330 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	66.7	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 61.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4804	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UWN, UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	1/2485 (0.0%)	1.26	6/3409 (0.2%)
1	B	0.84	0/2391	1.25	6/3286 (0.2%)
All	All	0.87	1/4876 (0.0%)	1.26	12/6695 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	390	VAL	CA-C	5.58	1.59	1.52

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	445	VAL	N-CA-C	-6.90	104.99	111.48
1	B	409	ASP	CA-CB-CG	6.76	119.36	112.60
1	A	486	LYS	CA-C-N	5.89	128.43	120.65
1	A	486	LYS	C-N-CA	5.89	128.43	120.65
1	B	485	ASP	CA-CB-CG	5.84	118.44	112.60
1	A	417	GLU	CB-CG-CD	5.62	122.16	112.60
1	B	452	LYS	CA-C-N	5.46	127.59	120.28
1	B	452	LYS	C-N-CA	5.46	127.59	120.28
1	B	475	PHE	CA-CB-CG	5.43	119.23	113.80
1	B	517	ASP	CA-CB-CG	5.17	117.77	112.60
1	A	468	LYS	CA-C-N	5.02	127.30	120.63
1	A	468	LYS	C-N-CA	5.02	127.30	120.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2394	0	2120	9	0
1	B	2303	0	1973	9	0
2	A	78	0	86	2	0
3	A	24	0	0	0	0
3	B	5	0	0	0	0
All	All	4804	0	4179	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:407:ASN:HB2	1:A:440:PRO:HD2	1.84	0.60
1:B:380:VAL:HG22	1:B:388:ILE:HG12	1.92	0.51
2:A:1002:UWN:CAG	2:A:1002:UWN:H12	2.40	0.51
1:B:387:PHE:HD2	1:B:437:ILE:HG13	1.78	0.48
1:B:493:ARG:HG2	1:B:541:GLN:HB2	1.95	0.48
1:B:402:LEU:HD11	1:B:413:ASP:HB3	1.96	0.48
1:A:393:VAL:HG13	1:A:401:PHE:HB2	1.95	0.47
1:A:294:ILE:HD12	1:A:296:LEU:HD21	1.97	0.46
1:B:484:VAL:HG22	1:B:492:ILE:HG12	1.97	0.45
1:A:274:ASP:OD2	2:A:1002:UWN:H17	2.17	0.44
1:B:294:ILE:HD12	1:B:296:LEU:HD21	2.00	0.44
1:B:319:VAL:HG13	1:B:342:TRP:CG	2.53	0.43
1:A:380:VAL:HG22	1:A:388:ILE:HG12	2.00	0.43
1:A:484:VAL:HG22	1:A:492:ILE:HG12	2.01	0.42
1:A:234:TRP:CG	1:A:530:VAL:HG13	2.54	0.42
1:B:263:GLY:HA3	1:B:358:LEU:HD22	2.00	0.42
1:A:331:PRO:HB2	1:A:335:TYR:HB3	2.02	0.41
1:A:254:HIS:HD2	1:A:525:PRO:HB3	1.86	0.41
1:B:393:VAL:HG13	1:B:401:PHE:HB2	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	305/332 (92%)	294 (96%)	11 (4%)	0	100	100
1	B	300/332 (90%)	286 (95%)	14 (5%)	0	100	100
All	All	605/664 (91%)	580 (96%)	25 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/295 (79%)	227 (98%)	5 (2%)	45	64
1	B	212/295 (72%)	204 (96%)	8 (4%)	29	44
All	All	444/590 (75%)	431 (97%)	13 (3%)	37	55

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	256	SER
1	A	343	GLN
1	A	377	LEU
1	A	475	PHE
1	A	500	THR
1	B	304	CYS
1	B	343	GLN
1	B	386	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	389	CYS
1	B	496	THR
1	B	500	THR
1	B	501	ASP
1	B	527	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	277	HIS
1	A	424	HIS
1	A	444	ASN
1	B	424	HIS
1	B	503	HIS
1	B	514	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 31 ligands modelled in this entry, 29 are unknown - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UWN	A	1002	-	44,44,44	1.47	5 (11%)	61,61,61	1.32	8 (13%)
2	UWN	A	1001	-	44,44,44	1.68	6 (13%)	61,61,61	1.36	9 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UWN	A	1002	-	-	1/28/62/62	0/6/6/6
2	UWN	A	1001	-	-	0/28/62/62	0/6/6/6

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	UWN	CBG-CBC	-6.79	1.40	1.50
2	A	1002	UWN	CBG-CBC	-5.43	1.42	1.50
2	A	1001	UWN	CBE-CBB	-5.40	1.40	1.50
2	A	1002	UWN	CBE-CBB	-4.95	1.41	1.50
2	A	1001	UWN	CAR-CBI	2.71	1.59	1.52
2	A	1001	UWN	CAX-N4	2.50	1.51	1.47
2	A	1002	UWN	CAR-CBI	2.35	1.58	1.52
2	A	1001	UWN	C3-N4	2.17	1.50	1.47
2	A	1001	UWN	CAF-CBD	2.08	1.42	1.39
2	A	1002	UWN	CBD-N3	-2.08	1.36	1.40
2	A	1002	UWN	C3-N4	2.00	1.50	1.47

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1002	UWN	C8-N5-C6	-4.25	110.01	114.12
2	A	1001	UWN	C7-N5-C6	-3.77	110.48	114.12
2	A	1002	UWN	CBE-CBB-N4	3.17	122.62	118.66
2	A	1001	UWN	CAU-N1-CBI	2.92	116.94	114.12
2	A	1002	UWN	CAN-CAV-N1	-2.78	99.50	104.17
2	A	1001	UWN	CAM-CAU-N1	-2.63	99.76	104.17
2	A	1001	UWN	CBE-CBB-N4	2.58	121.89	118.66
2	A	1002	UWN	C10-C7-N5	-2.55	99.89	104.17
2	A	1002	UWN	CAQ-CAY-N2	-2.54	105.68	110.66
2	A	1001	UWN	CAR-CBI-CAQ	2.53	116.88	111.17
2	A	1001	UWN	C3-N4-CAX	2.52	117.83	112.68
2	A	1001	UWN	CAQ-CAY-N2	-2.52	105.72	110.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1002	UWN	C3-N4-CAX	2.12	117.01	112.68
2	A	1001	UWN	C2-CAF-CBD	2.09	122.14	119.73
2	A	1002	UWN	CBG-CBF-N3	-2.08	116.87	120.50
2	A	1002	UWN	CAV-N1-CBI	2.07	116.12	114.12
2	A	1001	UWN	CBD-N3-CBF	-2.05	121.50	126.73

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1002	UWN	CAR-CBI-N1-CAV

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1002	UWN	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	309/332 (93%)	0.48	15 (4%) 35 34	37, 69, 114, 134	0
1	B	306/332 (92%)	0.75	26 (8%) 16 16	47, 84, 119, 143	0
All	All	615/664 (92%)	0.61	41 (6%) 24 23	37, 76, 117, 143	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	445	VAL	9.9
1	A	517	ASP	4.0
1	A	443	PRO	4.0
1	A	233	CYS	3.5
1	B	369	SER	3.1
1	B	361	ASN	3.1
1	A	367	ILE	3.0
1	B	544	LEU	2.9
1	B	444	ASN	2.9
1	B	233	CYS	2.8
1	B	336	LYS	2.8
1	B	264	PHE	2.7
1	A	366	VAL	2.6
1	A	237	TYR	2.6
1	A	481	LEU	2.5
1	A	512	TRP	2.5
1	B	234	TRP	2.5
1	B	512	TRP	2.4
1	A	522	ALA	2.4
1	B	442	TYR	2.4
1	B	239	GLU	2.4
1	A	244	VAL	2.4
1	B	298	PHE	2.3
1	B	498	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	500	THR	2.3
1	B	494	VAL	2.3
1	B	339	GLU	2.2
1	A	239	GLU	2.2
1	B	428	TRP	2.2
1	B	475	PHE	2.2
1	A	469	VAL	2.1
1	B	232	TRP	2.1
1	B	254	HIS	2.1
1	B	243	ALA	2.1
1	B	447	HIS	2.1
1	B	231	ALA	2.1
1	B	488	ASN	2.1
1	A	476	GLN	2.0
1	B	476	GLN	2.0
1	A	232	TRP	2.0
1	B	509	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	UNX	A	1026	1/1	0.54	0.46	82,82,82,82	0
3	UNX	A	1017	1/1	0.73	0.31	53,53,53,53	0
3	UNX	A	1022	1/1	0.79	0.15	57,57,57,57	0
3	UNX	A	1005	1/1	0.79	0.43	68,68,68,68	0
3	UNX	A	1025	1/1	0.82	0.21	68,68,68,68	0
3	UNX	A	1013	1/1	0.82	0.28	56,56,56,56	0

Continued on next page...



Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	UNX	A	1019	1/1	0.83	0.23	51,51,51,51	0
3	UNX	A	1010	1/1	0.84	0.39	66,66,66,66	0
3	UNX	A	1004	1/1	0.85	0.31	50,50,50,50	0
3	UNX	A	1014	1/1	0.86	0.15	54,54,54,54	0
3	UNX	A	1012	1/1	0.86	0.14	58,58,58,58	0
3	UNX	A	1008	1/1	0.86	0.16	56,56,56,56	0
3	UNX	B	602	1/1	0.86	0.09	70,70,70,70	0
3	UNX	B	603	1/1	0.86	0.18	73,73,73,73	0
3	UNX	A	1020	1/1	0.89	0.14	54,54,54,54	0
3	UNX	A	1011	1/1	0.89	0.18	61,61,61,61	0
3	UNX	A	1023	1/1	0.90	0.13	63,63,63,63	0
3	UNX	B	601	1/1	0.90	0.52	66,66,66,66	0
3	UNX	B	605	1/1	0.90	0.31	66,66,66,66	0
2	UWN	A	1001	39/39	0.92	0.12	43,69,83,83	0
3	UNX	A	1021	1/1	0.92	0.17	65,65,65,65	0
2	UWN	A	1002	39/39	0.93	0.12	41,65,76,78	0
3	UNX	A	1003	1/1	0.93	0.17	58,58,58,58	0
3	UNX	A	1006	1/1	0.93	0.18	48,48,48,48	0
3	UNX	A	1007	1/1	0.94	0.10	52,52,52,52	0
3	UNX	A	1015	1/1	0.94	0.15	44,44,44,44	0
3	UNX	A	1024	1/1	0.94	0.14	46,46,46,46	0
3	UNX	B	604	1/1	0.95	0.27	70,70,70,70	0
3	UNX	A	1016	1/1	0.95	0.11	43,43,43,43	0
3	UNX	A	1018	1/1	0.97	0.13	61,61,61,61	0
3	UNX	A	1009	1/1	0.97	0.27	68,68,68,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UWN A 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.