



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 06:07 PM UTC

PDB ID : 4FLF / pdb_00004ff
Title : Structure of three phase partition treated lipase from *Thermomyces lanuginosa* at 2.15Å resolution.
Authors : Kumar, M.; Mukherjee, J.; Sinha, M.; Kaur, P.; Gupta, M.N.; Sharma, S.; Singh, T.P.
Deposited on : 2012-06-14
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	12% 76% 22% .
1	B	269	15% 71% 25% .

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 4509 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lipase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	269	Total	C	N	O	S	0	0	0
			2071	1303	359	403	6			
1	B	269	Total	C	N	O	S	0	0	0
			2071	1303	359	403	6			

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 4-nitrobenzaldehyde (CCD ID: XXH) (formula: $C_7H_5NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			11	7	1	3		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	148	Total 148	O 148	0	0
6	B	140	Total 140	O 140	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lipase



- Molecule 1: Lipase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	140.00Å 140.00Å 80.50Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.15 50.00 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.0 (50.00-2.15) 99.0 (50.00-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.14Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.224 , 0.268 0.234 , 0.275	Depositor DCC
R_{free} test set	2459 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	33.4	Xtriage
Anisotropy	0.683	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.043 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4509	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, XXH, NAG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.48	13/2121 (0.6%)	1.22	10/2887 (0.3%)
1	B	1.41	9/2121 (0.4%)	1.32	9/2887 (0.3%)
All	All	1.45	22/4242 (0.5%)	1.27	19/5774 (0.3%)

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	255	ILE	CA-C	6.85	1.59	1.52
1	A	174	PRO	CA-C	6.52	1.60	1.52
1	A	229	PRO	CA-C	6.41	1.60	1.52
1	A	222	ILE	N-CA	6.40	1.53	1.46
1	A	125	ARG	C-N	6.24	1.42	1.33
1	B	63	VAL	CA-CB	-6.11	1.47	1.54
1	B	35	THR	C-N	6.01	1.42	1.33
1	B	85	SER	CA-C	-5.98	1.45	1.53
1	B	262	PHE	C-O	-5.96	1.16	1.24
1	A	262	PHE	C-O	-5.94	1.16	1.24
1	A	264	LEU	C-O	-5.56	1.17	1.23
1	A	237	LYS	C-O	-5.52	1.17	1.24
1	B	173	ALA	N-CA	5.47	1.53	1.45
1	A	198	HIS	C-O	-5.45	1.17	1.24
1	A	141	VAL	CA-C	-5.28	1.46	1.52
1	A	77	VAL	C-O	-5.26	1.18	1.24
1	B	209	ARG	C-O	5.25	1.30	1.24
1	A	68	ALA	CA-C	-5.22	1.46	1.52
1	A	14	ALA	N-CA	5.17	1.52	1.46
1	B	260	TRP	N-CA	-5.12	1.39	1.46
1	A	73	ASN	C-O	-5.09	1.17	1.24
1	B	124	LEU	C-O	-5.01	1.18	1.24

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	27	ASP	N-CA-C	-7.64	97.71	113.31
1	A	95	PHE	N-CA-C	6.76	120.42	112.72
1	A	221	TRP	CA-C-N	-6.38	114.40	122.37
1	A	221	TRP	C-N-CA	-6.38	114.40	122.37
1	A	22	CYS	N-CA-C	6.22	120.88	113.16
1	B	63	VAL	N-CA-C	6.06	117.40	109.58
1	B	52	LEU	N-CA-C	-5.94	104.78	112.68
1	B	203	VAL	CB-CA-C	-5.68	108.33	114.35
1	A	176	VAL	CB-CA-C	5.54	116.61	110.62
1	B	2	VAL	CB-CA-C	-5.54	103.98	111.63
1	B	100	ILE	CB-CA-C	-5.37	102.48	111.29
1	B	94	ASN	N-CA-CB	-5.34	101.71	110.52
1	A	77	VAL	N-CA-C	5.21	115.35	107.75
1	B	166	ILE	CB-CA-C	-5.19	104.55	110.73
1	A	49	ALA	N-CA-C	5.14	117.27	108.90
1	B	95	PHE	N-CA-C	5.08	119.17	112.92
1	A	169	PHE	CA-C-N	-5.03	114.77	122.87
1	A	169	PHE	C-N-CA	-5.03	114.77	122.87
1	A	171	TYR	N-CA-C	5.00	116.56	108.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2071	0	1964	52	0
1	B	2071	0	1964	52	0
2	A	14	0	13	0	0
2	B	14	0	13	0	0
3	A	30	0	39	3	0
3	B	6	0	8	0	0
4	A	11	0	5	0	0
5	B	4	0	6	1	0
6	A	148	0	0	1	0
6	B	140	0	0	2	0
All	All	4509	0	4012	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:24:LYS:O	1:A:25:ASN:HB2	1.41	1.15
1:A:25:ASN:HB3	1:A:28:ALA:HB2	1.33	1.08
1:A:252:ILE:HG13	1:A:253:PRO:HD2	1.30	1.05
1:A:86:ILE:O	1:A:90:ILE:HG13	1.71	0.89
1:B:224:SER:HB3	1:B:234:ASP:OD1	1.74	0.87
6:A:546:HOH:O	1:B:227:LEU:HD13	1.76	0.86
1:A:225:GLY:H	3:A:305:GOL:H32	1.37	0.85
1:B:203:VAL:HG21	1:B:258:HIS:CE1	2.12	0.85
1:A:252:ILE:CG1	1:A:253:PRO:HD2	2.11	0.81
1:A:107:CYS:SG	1:A:180:ALA:C	2.65	0.80
1:A:25:ASN:CB	1:A:28:ALA:HB2	2.13	0.79
1:B:97:LEU:HD12	1:B:97:LEU:N	1.99	0.77
1:A:104:CYS:SG	1:A:184:PHE:HB2	2.25	0.76
1:B:26:ASN:HD21	1:B:56:GLU:HB2	1.48	0.76
1:B:9:GLN:NE2	1:B:139:ARG:HH12	1.84	0.76
1:B:26:ASN:O	1:B:27:ASP:HB3	1.84	0.76
1:A:252:ILE:HG13	1:A:253:PRO:CD	2.14	0.74
1:B:26:ASN:C	1:B:26:ASN:OD1	2.30	0.74
1:B:26:ASN:O	1:B:27:ASP:CB	2.34	0.74
1:B:101:ASN:HD22	1:B:105:SER:HA	1.56	0.68
1:B:26:ASN:ND2	1:B:56:GLU:HB2	2.08	0.68
1:B:203:VAL:HB	1:B:204:PRO:HD3	1.76	0.68
1:A:203:VAL:HB	1:A:258:HIS:CE1	2.30	0.67
1:B:97:LEU:N	1:B:97:LEU:CD1	2.58	0.67
1:A:107:CYS:SG	1:A:181:PHE:N	2.69	0.65
1:B:203:VAL:HG21	1:B:258:HIS:NE2	2.11	0.65
1:B:101:ASN:ND2	1:B:106:GLY:H	1.95	0.65
1:A:185:LEU:HB2	1:A:216:SER:HB3	1.78	0.65
1:A:24:LYS:O	1:A:25:ASN:CB	2.28	0.64
1:A:125:ARG:HD3	1:A:159:LEU:HD22	1.80	0.62
1:B:104:CYS:HB2	6:B:473:HOH:O	2.00	0.62
1:A:192:THR:HA	3:A:302:GOL:H2	1.82	0.60
1:A:256:PRO:C	1:A:258:HIS:H	2.10	0.60
1:A:181:PHE:O	1:A:185:LEU:HG	2.01	0.60
1:A:261:TYR:C	1:A:263:GLY:H	2.11	0.58
1:A:255:ILE:N	1:A:256:PRO:CD	2.66	0.58
1:B:98:LYS:HD3	1:B:111:ASP:OD1	2.04	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:ASN:HB3	1:A:28:ALA:CB	2.22	0.56
1:A:252:ILE:CG1	1:A:253:PRO:CD	2.79	0.56
1:B:203:VAL:CG2	1:B:258:HIS:CE1	2.88	0.56
1:B:204:PRO:HB2	1:B:247:ASN:OD1	2.06	0.55
1:B:1:GLU:HG3	1:B:233:ASN:O	2.07	0.54
1:A:107:CYS:SG	1:A:180:ALA:CB	2.96	0.54
1:A:100:ILE:HG13	1:A:107:CYS:O	2.09	0.53
1:B:26:ASN:OD1	1:B:27:ASP:N	2.42	0.52
1:A:101:ASN:HA	1:A:104:CYS:O	2.11	0.51
1:B:162:ASN:O	1:B:162:ASN:OD1	2.27	0.51
1:A:203:VAL:CB	1:A:258:HIS:CE1	2.95	0.50
1:B:60:VAL:HG13	1:B:61:GLY:N	2.27	0.50
1:A:4:GLN:NE2	1:A:232:ARG:HD3	2.27	0.50
1:B:186:THR:HG23	1:B:217:SER:O	2.12	0.50
1:B:101:ASN:HD21	1:B:106:GLY:H	1.60	0.49
1:B:93:LEU:HD12	6:B:403:HOH:O	2.12	0.49
1:A:104:CYS:SG	1:A:184:PHE:CB	2.96	0.49
1:A:225:GLY:H	3:A:305:GOL:C3	2.19	0.49
1:B:98:LYS:HG2	1:B:99:GLU:O	2.13	0.49
1:A:255:ILE:HB	1:A:256:PRO:HD3	1.94	0.48
1:B:97:LEU:HD11	1:B:213:TYR:CE1	2.48	0.48
1:A:201:ASP:OD1	1:A:258:HIS:HB2	2.12	0.48
1:A:198:HIS:O	1:A:199:THR:C	2.56	0.48
1:A:107:CYS:SG	1:A:180:ALA:HB1	2.54	0.48
1:A:206:LEU:HA	1:A:207:PRO:C	2.38	0.47
1:B:9:GLN:HE22	1:B:139:ARG:HH22	1.63	0.47
1:B:60:VAL:HG12	1:B:119:SER:CB	2.44	0.47
1:B:206:LEU:HA	1:B:207:PRO:C	2.40	0.47
1:B:234:ASP:O	1:B:236:VAL:HG23	2.15	0.47
1:A:4:GLN:NE2	1:A:4:GLN:HA	2.31	0.46
1:A:125:ARG:HD3	1:A:159:LEU:CD2	2.44	0.46
1:B:171:TYR:CE2	1:B:196:ILE:HD13	2.51	0.46
1:B:20:ALA:HB1	1:B:81:ARG:HB2	1.98	0.46
1:A:4:GLN:HA	1:A:4:GLN:HE21	1.80	0.46
1:B:81:ARG:CD	1:B:82:GLY:O	2.64	0.46
1:B:220:TYR:CE1	1:B:237:LYS:HG3	2.50	0.46
1:A:255:ILE:N	1:A:256:PRO:HD2	2.30	0.46
1:A:223:LYS:HB2	1:A:223:LYS:HE2	1.63	0.46
1:B:15:GLN:NE2	5:B:303:EDO:H12	2.30	0.45
1:A:203:VAL:HG21	1:A:258:HIS:CE1	2.50	0.45
1:B:156:GLY:O	1:B:160:ARG:HG3	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:261:TYR:C	1:B:263:GLY:H	2.24	0.45
1:B:26:ASN:C	1:B:28:ALA:N	2.73	0.45
1:A:28:ALA:O	1:A:29:PRO:C	2.59	0.44
1:A:108:ARG:HG3	1:A:178:ASN:ND2	2.32	0.44
1:B:57:ASP:HA	1:B:62:ASP:C	2.41	0.44
1:A:256:PRO:C	1:A:258:HIS:N	2.73	0.44
1:B:81:ARG:HD2	1:B:82:GLY:O	2.17	0.43
1:A:252:ILE:CG1	1:A:253:PRO:N	2.82	0.43
1:B:13:PHE:CE1	1:B:141:VAL:HG11	2.54	0.43
1:B:160:ARG:HA	1:B:166:ILE:HD12	2.01	0.43
1:A:187:VAL:O	1:A:188:GLN:C	2.60	0.43
1:A:94:ASN:OD1	1:A:95:PHE:N	2.52	0.43
1:A:203:VAL:CG2	1:A:258:HIS:CE1	3.02	0.43
1:B:24:LYS:HG2	1:B:37:THR:HG23	1.99	0.43
1:B:233:ASN:N	1:B:233:ASN:HD22	2.15	0.43
1:B:227:LEU:H	1:B:260:TRP:CG	2.36	0.42
1:A:4:GLN:HE21	1:A:232:ARG:HD3	1.84	0.42
1:A:203:VAL:N	1:A:204:PRO:CD	2.82	0.42
1:B:5:ASP:O	1:B:9:GLN:HG3	2.20	0.42
1:A:104:CYS:HB3	1:A:107:CYS:HB2	1.23	0.42
1:B:35:THR:HA	1:B:45:GLU:HG2	2.01	0.42
1:A:67:LEU:HA	1:A:77:VAL:O	2.19	0.41
1:B:200:ASN:O	1:B:201:ASP:C	2.63	0.41
1:B:55:PHE:CE1	1:B:63:VAL:HG12	2.56	0.41
1:A:179:ARG:O	1:A:179:ARG:HG3	2.20	0.40
1:B:99:GLU:CD	1:B:99:GLU:C	2.88	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/269 (99%)	253 (95%)	10 (4%)	4 (2%)	8	4
1	B	267/269 (99%)	255 (96%)	9 (3%)	3 (1%)	11	7
All	All	534/538 (99%)	508 (95%)	19 (4%)	7 (1%)	9	5

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	25	ASN
1	B	26	ASN
1	A	257	ALA
1	B	242	ASP
1	A	146	SER
1	B	100	ILE
1	A	253	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/220 (100%)	217 (99%)	3 (1%)	59	66
1	B	220/220 (100%)	214 (97%)	6 (3%)	39	41
All	All	440/440 (100%)	431 (98%)	9 (2%)	48	54

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	SER
1	A	215	HIS
1	A	223	LYS
1	B	1	GLU
1	B	81	ARG
1	B	86	ILE
1	B	183	GLU
1	B	235	ILE
1	B	252	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	249	GLN
1	B	9	GLN
1	B	88	ASN
1	B	92	ASN
1	B	101	ASN
1	B	162	ASN
1	B	233	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	XXH	A	307	-	11,11,11	1.12	1 (9%)	12,14,14	1.92	3 (25%)
3	GOL	A	306	-	5,5,5	0.54	0	5,5,5	0.90	0
3	GOL	B	302	-	5,5,5	0.57	0	5,5,5	0.75	0
3	GOL	A	303	-	5,5,5	0.39	0	5,5,5	0.96	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	B	303	-	3,3,3	0.65	0	2,2,2	0.25	0
3	GOL	A	302	-	5,5,5	0.81	0	5,5,5	0.43	0
3	GOL	A	304	-	5,5,5	1.13	0	5,5,5	0.94	0
2	NAG	B	301	1	14,14,15	0.76	0	17,19,21	1.53	4 (23%)
2	NAG	A	301	1	14,14,15	0.76	0	17,19,21	1.53	4 (23%)
3	GOL	A	305	-	5,5,5	0.98	0	5,5,5	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	XXH	A	307	-	-	0/4/6/6	0/1/1/1
3	GOL	A	306	-	-	2/4/4/4	-
3	GOL	B	302	-	-	2/4/4/4	-
3	GOL	A	303	-	-	4/4/4/4	-
5	EDO	B	303	-	-	1/1/1/1	-
3	GOL	A	302	-	-	2/4/4/4	-
3	GOL	A	304	-	-	1/4/4/4	-
2	NAG	B	301	1	-	2/6/23/26	0/1/1/1
2	NAG	A	301	1	-	2/6/23/26	0/1/1/1
3	GOL	A	305	-	-	4/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	307	XXH	C2-C3	2.06	1.42	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	307	XXH	O1-N1-C1	-4.59	112.50	118.82
2	A	301	NAG	C2-N2-C7	-3.36	118.39	122.90
2	B	301	NAG	C2-N2-C7	-3.34	118.42	122.90
2	A	301	NAG	O5-C5-C4	-3.01	103.51	110.83
2	B	301	NAG	O5-C5-C4	-3.00	103.53	110.83
4	A	307	XXH	O3-C5-C4	-2.72	115.35	124.56
4	A	307	XXH	C2-C3-C4	-2.61	117.85	121.22
2	B	301	NAG	O5-C1-C2	-2.38	107.61	111.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	NAG	O5-C1-C2	-2.37	107.62	111.29
2	B	301	NAG	O5-C5-C6	2.11	111.78	107.66
2	A	301	NAG	O5-C5-C6	2.08	111.72	107.66

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	303	GOL	O1-C1-C2-C3
3	A	305	GOL	O1-C1-C2-O2
3	A	305	GOL	O1-C1-C2-C3
3	A	305	GOL	C1-C2-C3-O3
3	B	302	GOL	O1-C1-C2-C3
2	A	301	NAG	O5-C5-C6-O6
2	B	301	NAG	O5-C5-C6-O6
2	A	301	NAG	C4-C5-C6-O6
2	B	301	NAG	C4-C5-C6-O6
3	A	303	GOL	O1-C1-C2-O2
3	A	303	GOL	O2-C2-C3-O3
3	A	302	GOL	C1-C2-C3-O3
3	A	303	GOL	C1-C2-C3-O3
3	A	306	GOL	O1-C1-C2-C3
3	A	305	GOL	O2-C2-C3-O3
3	B	302	GOL	O1-C1-C2-O2
3	A	302	GOL	O2-C2-C3-O3
3	A	306	GOL	O1-C1-C2-O2
3	A	304	GOL	O1-C1-C2-O2
5	B	303	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	303	EDO	1	0
3	A	302	GOL	1	0
3	A	305	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	269/269 (100%)	0.70	31 (11%)	9 10	17, 34, 56, 79	1 (0%)
1	B	269/269 (100%)	0.91	40 (14%)	5 6	18, 37, 60, 78	1 (0%)
All	All	538/538 (100%)	0.80	71 (13%)	7 8	17, 36, 59, 79	2 (0%)

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	104	CYS	7.0
1	B	234	ASP	6.3
1	A	257	ALA	4.4
1	A	185	LEU	4.4
1	A	60	VAL	4.4
1	A	107	CYS	4.2
1	A	103	ILE	4.0
1	B	1	GLU	4.0
1	B	176	VAL	3.9
1	B	206	LEU	3.8
1	A	106	GLY	3.7
1	A	94	ASN	3.6
1	B	155	ALA	3.4
1	A	255	ILE	3.4
1	B	106	GLY	3.3
1	B	107	CYS	3.3
1	B	246	GLY	3.2
1	A	26	ASN	3.2
1	A	23	GLY	3.2
1	A	102	ASP	3.1
1	A	101	ASN	3.1
1	B	184	PHE	3.1
1	B	185	LEU	3.1
1	A	176	VAL	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	97	LEU	3.0
1	B	163	GLY	3.0
1	A	90	ILE	3.0
1	B	28	ALA	2.9
1	A	30	ALA	2.8
1	B	60	VAL	2.8
1	B	26	ASN	2.7
1	A	125	ARG	2.7
1	A	24	LYS	2.7
1	B	245	GLY	2.7
1	A	163	GLY	2.7
1	B	187	VAL	2.6
1	A	25	ASN	2.6
1	B	25	ASN	2.6
1	A	95	PHE	2.6
1	B	125	ARG	2.6
1	B	154	VAL	2.5
1	B	181	PHE	2.5
1	B	102	ASP	2.5
1	B	151	LEU	2.5
1	B	233	ASN	2.5
1	B	34	ILE	2.5
1	B	100	ILE	2.5
1	B	27	ASP	2.4
1	B	153	THR	2.4
1	A	184	PHE	2.4
1	B	103	ILE	2.4
1	A	161	GLY	2.4
1	A	100	ILE	2.4
1	B	58	SER	2.4
1	B	180	ALA	2.3
1	A	92	ASN	2.3
1	A	154	VAL	2.3
1	A	258	HIS	2.2
1	A	28	ALA	2.2
1	A	27	ASP	2.2
1	B	112	GLY	2.2
1	B	121	ALA	2.2
1	B	152	ALA	2.2
1	A	252	ILE	2.1
1	B	30	ALA	2.1
1	B	99	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	241	ILE	2.1
1	B	162	ASN	2.1
1	A	98	LYS	2.1
1	B	182	ALA	2.1
1	B	212	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	A	301	14/15	0.69	0.15	61,69,72,72	0
2	NAG	B	301	14/15	0.70	0.16	61,69,72,72	0
3	GOL	A	306	6/6	0.76	0.27	70,71,72,72	0
3	GOL	A	305	6/6	0.79	0.19	43,48,52,52	0
3	GOL	B	302	6/6	0.82	0.20	61,64,65,65	0
4	XXH	A	307	11/11	0.82	0.16	35,38,47,53	0
5	EDO	B	303	4/4	0.85	0.15	43,43,44,49	0
3	GOL	A	304	6/6	0.86	0.18	40,53,55,57	0
3	GOL	A	302	6/6	0.87	0.17	49,50,51,52	0
3	GOL	A	303	6/6	0.89	0.14	53,58,59,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.