



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:17 PM UTC

PDB ID : 6FLA / pdb_00006fla
Title : 3H5 Fab bound to EDIII of DenV 2 Xtal form 1
Authors : Flanagan, A.; Renner, M.; Grimes, J.M.
Deposited on : 2018-01-25
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	85% 8% 7%
1	H	227	85% 9% •
2	B	218	90% 8% •
2	L	218	87% 11% •
3	G	98	3% 80% 14% •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	I	101	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	H	301	-	X	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	0	0	0
			1601	1013	266	315	7			
1	H	217	Total	C	N	O	S	0	0	0
			1632	1030	272	323	7			

- Molecule 2 is a protein called Light Chain of 3H5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	216	Total	C	N	O	S	0	0	0
			1671	1039	287	338	7			
2	L	216	Total	C	N	O	S	0	0	0
			1671	1039	287	338	7			

- Molecule 3 is a protein called Domain III of Dengue virus 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	98	Total	C	N	O	S	0	0	0
			776	500	127	144	5			


- Molecule 4 is a protein called Domain III of Dengue virus 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	I	99	Total	C	N	O	S	0	0	0
			777	500	128	145	4			

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		
5	L	1	Total	Cl	0	0
			1	1		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	H	1	Total	C	O	0	0
			6	3	3		
6	L	1	Total	C	O	0	0
			6	3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heavy chain




- Molecule 1: Heavy chain




- Molecule 2: Light Chain of 3H5





- Molecule 2: Light Chain of 3H5



- Molecule 3: Domain III of Dengue virus 2



- Molecule 4: Domain III of Dengue virus 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	40.84Å 263.87Å 272.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.59 – 2.90 39.59 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.0 (39.59-2.90) 99.2 (39.59-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 2.90Å)	Xtriage
Refinement program	BUSTER 2.11.1	Depositor
R, R_{free}	0.203 , 0.221 (Not available) , 0.191	Depositor DCC
R_{free} test set	1689 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	66.7	Xtriage
Anisotropy	0.146	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 66.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8142	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	1/1644 (0.1%)	1.17	11/2245 (0.5%)
1	H	0.78	0/1676	1.18	12/2290 (0.5%)
2	B	0.82	0/1708	1.17	4/2319 (0.2%)
2	L	0.87	1/1708 (0.1%)	1.23	11/2319 (0.5%)
3	G	0.89	1/793 (0.1%)	1.25	8/1070 (0.7%)
4	I	0.91	0/794	1.25	3/1073 (0.3%)
All	All	0.83	3/8323 (0.0%)	1.20	49/11316 (0.4%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	144	CYS	CA-C	-7.19	1.43	1.52
2	L	215	ARG	CA-C	7.05	1.62	1.52
3	G	378	ILE	CG1-CD1	-5.97	1.28	1.51

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	215	ARG	CA-C-N	11.60	142.57	121.70
2	L	215	ARG	C-N-CA	11.60	142.57	121.70
3	G	297	MET	CA-C-N	7.62	135.42	121.70
3	G	297	MET	C-N-CA	7.62	135.42	121.70
1	A	66	GLY	CA-C-N	6.97	129.92	120.44
1	A	66	GLY	C-N-CA	6.97	129.92	120.44
4	I	352	ILE	N-CA-C	-6.96	105.82	111.81
2	B	62	VAL	N-CA-CB	6.65	119.66	111.21
2	L	62	VAL	N-CA-CB	6.58	119.56	111.21
3	G	352	ILE	N-CA-C	-6.49	106.23	111.81
1	H	66	GLY	CA-C-N	6.39	129.13	120.44
1	H	66	GLY	C-N-CA	6.39	129.13	120.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	67	LYS	N-CA-C	6.37	118.02	111.14
2	L	62	VAL	CB-CA-C	6.29	117.19	110.53
1	H	61	ALA	N-CA-C	-6.24	99.84	109.76
1	A	61	ALA	N-CA-C	-6.16	99.96	109.76
1	A	187	VAL	N-CA-CB	-6.10	105.16	111.87
3	G	311	GLU	CB-CG-CD	6.09	122.96	112.60
1	H	132	SER	N-CA-C	6.06	123.71	110.80
3	G	331	SER	CA-C-O	-5.91	112.06	120.16
1	A	67	LYS	N-CA-C	5.84	117.45	111.14
1	H	74	LYS	CA-C-N	5.75	128.31	120.54
1	H	74	LYS	C-N-CA	5.75	128.31	120.54
1	A	124	SER	N-CA-C	-5.74	99.88	109.24
2	B	70	GLY	N-CA-C	5.58	117.72	112.08
2	L	17	GLN	CB-CG-CD	5.52	121.98	112.60
2	B	62	VAL	CB-CA-C	5.49	116.35	110.53
1	A	187	VAL	CB-CA-C	5.41	115.73	109.89
1	A	96	CYS	N-CA-C	-5.40	101.36	109.95
3	G	329	ASP	CA-CB-CG	5.39	117.99	112.60
1	A	65	GLN	N-CA-C	5.38	117.58	111.02
1	H	124	SER	N-CA-C	-5.35	100.52	109.24
2	L	155	ASP	CA-CB-CG	5.30	117.91	112.60
3	G	390	ASN	CA-CB-CG	5.29	117.89	112.60
1	H	96	CYS	N-CA-C	-5.24	101.62	109.95
1	A	74	LYS	CA-C-N	5.22	127.59	120.54
1	A	74	LYS	C-N-CA	5.22	127.59	120.54
2	L	70	GLY	N-CA-C	5.22	117.35	112.08
1	H	187	VAL	N-CA-CB	-5.20	103.93	111.21
2	L	215	ARG	O-C-N	-5.19	115.52	122.43
2	B	141	ASN	N-CA-C	5.16	118.48	110.17
4	I	341	ASP	CA-CB-CG	5.15	117.75	112.60
2	L	141	ASN	N-CA-C	5.14	118.45	110.17
4	I	331	SER	CA-C-O	-5.09	113.18	120.16
3	G	311	GLU	CA-CB-CG	5.07	124.24	114.10
1	H	65	GLN	N-CA-C	5.05	117.18	111.11
2	L	61	GLY	CA-C-N	-5.04	118.40	123.04
2	L	61	GLY	C-N-CA	-5.04	118.40	123.04
1	H	121	THR	CB-CA-C	5.03	117.20	109.15

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1601	0	1554	3	0
1	H	1632	0	1583	6	0
2	B	1671	0	1608	7	0
2	L	1671	0	1608	11	0
3	G	776	0	789	12	0
4	I	777	0	788	11	0
5	B	1	0	0	0	0
5	L	1	0	0	0	0
6	H	6	0	8	1	0
6	L	6	0	8	1	0
All	All	8142	0	7946	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:297:MET:HA	3:G:298:SER:HB2	1.49	0.92
1:H:135:GLN:O	1:H:138:SER:HB2	1.70	0.91
4:I:331:SER:HB3	4:I:332:PRO:HD3	1.72	0.71
2:L:65:ARG:HD2	2:L:81:PRO:O	1.90	0.70
2:B:65:ARG:HD2	2:B:81:PRO:O	1.92	0.69
3:G:331:SER:HB3	3:G:332:PRO:HD3	1.76	0.65
1:H:168:HIS:HA	6:H:301:GOL:H12	1.81	0.61
3:G:297:MET:HA	3:G:298:SER:CB	2.28	0.59
4:I:348:LEU:HB3	4:I:372:PRO:HG3	1.88	0.56
2:L:199:GLU:HG2	2:L:210:VAL:HG22	1.87	0.56
2:L:34:LYS:HG2	2:L:54:LEU:HD13	1.87	0.55
3:G:312:ILE:CD1	3:G:378:ILE:HD13	2.37	0.55
1:A:123:PRO:HB3	1:A:149:TYR:HB3	1.90	0.54
3:G:340:MET:HE3	3:G:344:LYS:HG2	1.90	0.54
2:B:199:GLU:HG2	2:B:210:VAL:HG22	1.91	0.52
1:H:123:PRO:HB3	1:H:149:TYR:HB3	1.93	0.50
4:I:339:ILE:HG12	4:I:378:ILE:HD12	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:I:395:GLY:O	4:I:396:SER:HB2	2.12	0.49
2:L:45:GLY:H	6:L:302:GOL:H2	1.77	0.48
3:G:311:GLU:OE1	1:H:147:LYS:NZ	2.45	0.47
3:G:378:ILE:HD11	3:G:391:TRP:HB2	1.99	0.45
2:L:37:MET:HE2	2:L:37:MET:HB2	1.83	0.45
4:I:344:LYS:HE3	4:I:377:TYR:HE2	1.82	0.45
4:I:332:PRO:HA	4:I:358:VAL:O	2.17	0.45
3:G:312:ILE:HD13	3:G:378:ILE:HD13	1.99	0.44
2:L:18:ARG:O	2:L:18:ARG:HG3	2.16	0.44
4:I:377:TYR:OH	4:I:388:LYS:HD3	2.18	0.44
3:G:312:ILE:HD11	3:G:378:ILE:HD13	1.99	0.44
4:I:394:LYS:HD3	4:I:395:GLY:H	1.82	0.44
4:I:331:SER:HB3	4:I:332:PRO:CD	2.45	0.44
3:G:332:PRO:HA	3:G:358:VAL:O	2.18	0.43
1:H:136:THR:HG22	1:H:137:ASN:OD1	2.19	0.43
2:B:6:GLN:HE21	2:B:21:ILE:CD1	2.32	0.43
2:L:194:ASN:HD21	2:L:216:ASN:C	2.26	0.43
3:G:360:GLU:HB2	3:G:363:SER:HB3	2.01	0.42
1:A:57:ASP:HA	2:L:192:ARG:NH2	2.33	0.42
2:B:18:ARG:HG3	2:B:80:ASP:HB2	2.01	0.42
1:H:138:SER:HB3	1:H:189:SER:OG	2.19	0.42
4:I:383:GLU:HG3	2:L:98:VAL:HG23	2.01	0.42
2:B:59:GLU:HB2	2:B:62:VAL:HG13	2.01	0.42
2:B:34:LYS:HG2	2:B:54:LEU:HD13	2.00	0.42
3:G:378:ILE:HD11	3:G:391:TRP:CB	2.50	0.41
4:I:348:LEU:CB	4:I:372:PRO:HG3	2.50	0.41
2:L:59:GLU:HB2	2:L:62:VAL:HG13	2.03	0.41
2:B:37:MET:HE2	2:B:37:MET:HB2	1.83	0.41
2:L:214:ASN:C	2:L:216:ASN:HA	2.46	0.41
1:A:158:TRP:HZ3	1:A:214:ILE:HD12	1.86	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/227 (92%)	203 (98%)	4 (2%)	1 (0%)	24	54
1	H	215/227 (95%)	210 (98%)	4 (2%)	1 (0%)	24	54
2	B	214/218 (98%)	210 (98%)	3 (1%)	1 (0%)	24	54
2	L	214/218 (98%)	210 (98%)	3 (1%)	1 (0%)	24	54
3	G	96/98 (98%)	90 (94%)	4 (4%)	2 (2%)	5	21
4	I	97/101 (96%)	91 (94%)	4 (4%)	2 (2%)	5	21
All	All	1044/1089 (96%)	1014 (97%)	22 (2%)	8 (1%)	16	44

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	G	331	SER
4	I	331	SER
3	G	298	SER
4	I	396	SER
1	H	132	SER
2	B	72	ARG
2	L	72	ARG
1	A	138	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/189 (94%)	170 (96%)	7 (4%)	28	62
1	H	180/189 (95%)	173 (96%)	7 (4%)	28	63
2	B	191/193 (99%)	186 (97%)	5 (3%)	40	73
2	L	191/193 (99%)	185 (97%)	6 (3%)	35	69
3	G	88/88 (100%)	81 (92%)	7 (8%)	11	34
4	I	88/90 (98%)	83 (94%)	5 (6%)	18	49

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	915/942 (97%)	878 (96%)	37 (4%)	28	62

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	VAL
1	A	57	ASP
1	A	63	LYS
1	A	75	SER
1	A	140	VAL
1	A	187	VAL
1	A	195	GLU
2	B	7	SER
2	B	11	LEU
2	B	21	ILE
2	B	27	GLU
2	B	62	VAL
3	G	297	MET
3	G	311	GLU
3	G	321	VAL
3	G	343	GLU
3	G	370	GLU
3	G	379	ILE
3	G	390	ASN
1	H	11	VAL
1	H	57	ASP
1	H	63	LYS
1	H	75	SER
1	H	138	SER
1	H	187	VAL
1	H	195	GLU
4	I	321	VAL
4	I	343	GLU
4	I	370	GLU
4	I	376	SER
4	I	394	LYS
2	L	11	LEU
2	L	18	ARG
2	L	21	ILE
2	L	27	GLU
2	L	62	VAL
2	L	71	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1	GLN
1	A	3	GLN
1	A	35	GLN
2	B	17	GLN
2	B	57	ASN
2	B	96	ASN
3	G	325	GLN
3	G	355	ASN
3	G	390	ASN
1	H	39	GLN
1	H	43	GLN
4	I	355	ASN
2	L	42	GLN
2	L	57	ASN
2	L	96	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	L	302	-	5,5,5	1.01	0	5,5,5	0.86	0
6	GOL	H	301	-	5,5,5	1.74	2 (40%)	5,5,5	2.05	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	L	302	-	-	2/4/4/4	-
6	GOL	H	301	-	-	4/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	H	301	GOL	O1-C1	2.71	1.53	1.42
6	H	301	GOL	C1-C2	2.55	1.61	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	H	301	GOL	O1-C1-C2	3.29	125.20	110.38

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	H	301	GOL	O1-C1-C2-C3
6	H	301	GOL	C1-C2-C3-O3
6	H	301	GOL	O2-C2-C3-O3
6	L	302	GOL	C1-C2-C3-O3
6	H	301	GOL	O1-C1-C2-O2
6	L	302	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	L	302	GOL	1	0
6	H	301	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	212/227 (93%)	0.03	2 (0%) 81 75	46, 68, 93, 116	0
1	H	217/227 (95%)	-0.01	2 (0%) 81 75	34, 61, 89, 101	0
2	B	216/218 (99%)	-0.01	1 (0%) 87 83	45, 64, 87, 100	0
2	L	216/218 (99%)	-0.06	2 (0%) 81 75	34, 57, 83, 95	0
3	G	98/98 (100%)	0.56	3 (3%) 51 42	46, 80, 116, 129	0
4	I	99/101 (98%)	0.38	4 (4%) 42 34	50, 75, 101, 117	0
All	All	1058/1089 (97%)	0.08	14 (1%) 75 67	34, 65, 97, 129	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	131	GLY	3.5
2	L	23	CYS	3.2
4	I	397	SER	2.7
3	G	344	LYS	2.6
1	H	1	GLN	2.5
2	B	216	ASN	2.3
1	H	215	GLU	2.3
3	G	373	PHE	2.3
3	G	342	LEU	2.2
1	A	215	GLU	2.2
4	I	354	VAL	2.1
4	I	299	TYR	2.1
2	L	213	PHE	2.1
4	I	390	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	GOL	L	302	6/6	0.82	0.15	69,70,71,73	0
6	GOL	H	301	6/6	0.84	0.14	57,59,59,60	0
5	CL	B	301	1/1	0.98	0.09	50,50,50,50	0
5	CL	L	301	1/1	0.98	0.10	52,52,52,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.