



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:21 PM UTC

PDB ID : 7FLX / pdb_00007flx
Title : PanDDA analysis group deposition – Aar2/RNaseH in complex with fragment P05G08 from the F2X-Universal Library
Authors : Barthel, T.; Wollenhaupt, J.; Lima, G.M.A.; Wahl, M.C.; Weiss, M.S.
Deposited on : 2022-08-26
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	36% 78% 13% 8%
2	B	308	39% 86% 11% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9202 atoms, of which 4524 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	237	4068	1287	2060	336	373	12	0	21	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	300	5044	1654	2464	421	485	20	0	17	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	LEU	conflict	UNP P32357
B	167	SER	LYS	conflict	UNP P32357
B	?	-	LEU	deletion	UNP P32357
B	?	-	GLN	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	GLY	deletion	UNP P32357
B	?	-	SER	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ASN	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	170	SER	ASP	conflict	UNP P32357

- Molecule 3 is 2-chlorobenzohydrazide (CCD ID: 5G1) (formula: $C_7H_7ClN_2O$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0	0
			11	7	1	2	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	40	Total	O	0	0
			40	40		
4	B	39	Total	O	0	0
			39	39		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pre-mRNA-splicing factor 8



• Molecule 2: A1 cistron-splicing factor AAR2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.28Å 81.96Å 93.49Å 90.00° 108.25° 90.00°	Depositor
Resolution (Å)	44.39 – 1.75 44.39 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.3 (44.39-1.75) 99.3 (44.39-1.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.92 (at 1.75Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.256 , 0.287 0.266 , 0.297	Depositor DCC
R_{free} test set	2098 reflections (3.30%)	wwPDB-VP
Wilson B-factor (Å ²)	44.3	Xtriage
Anisotropy	0.317	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9202	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 5G1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	1/2149 (0.0%)	0.86	4/2911 (0.1%)
2	B	0.65	0/2739	0.79	2/3699 (0.1%)
All	All	0.71	1/4888 (0.0%)	0.82	6/6610 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2040	TRP	C-N	16.22	1.53	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	VAL	N-CA-C	-6.25	101.97	108.15
1	A	2040	TRP	CA-C-N	6.24	126.21	120.03
1	A	2040	TRP	C-N-CA	6.24	126.21	120.03
2	B	252	SER	CA-CB-OG	-5.84	99.43	111.10
1	A	1992	TYR	CA-C-O	-5.40	112.90	119.43
1	A	1993	ASP	CA-CB-CG	5.27	117.87	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2008	2060	1974	21	0
2	B	2580	2464	2398	26	0
3	A	11	0	0	0	0
4	A	40	0	0	0	0
4	B	39	0	0	2	0
All	All	4678	4524	4372	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1890:PHE:CD2	1:A:1988:LEU:HD21	2.13	0.82
1:A:1890:PHE:CE2	1:A:1988:LEU:HD21	2.16	0.81
2:B:70:GLN:HB3	2:B:81:MET:HE2	1.64	0.80
1:A:2064:GLY:O	1:A:2068:ASN:N	2.20	0.74
2:B:96:PHE:HB2	2:B:102:MET:HE3	1.72	0.70
1:A:1962:ARG:O	1:A:2013:ARG:NH1	2.27	0.67
1:A:1878:CYS:SG	1:A:1969:MET:HE3	2.34	0.67
1:A:1890:PHE:CD2	1:A:1988:LEU:CD2	2.77	0.66
1:A:2061:THR:O	1:A:2064:GLY:N	2.30	0.64
2:B:50:ASP:OD1	2:B:51:ASN:N	2.31	0.63
2:B:68:TYR:CE2	2:B:81:MET:HE3	2.34	0.63
2:B:1:MET:N	4:B:401:HOH:O	2.33	0.60
2:B:277:GLU:CD	2:B:277:GLU:H	2.13	0.57
2:B:96:PHE:CB	2:B:102:MET:HE3	2.34	0.57
2:B:43:VAL:HG13	2:B:43:VAL:O	2.06	0.56
1:A:1979[B]:MET:HA	1:A:1979[B]:MET:HE2	1.87	0.55
1:A:1895:HIS:O	1:A:1898[A]:VAL:HG22	2.07	0.55
1:A:1848:ILE:H	1:A:1931[A]:LYS:HZ2	1.57	0.53
2:B:70:GLN:CB	2:B:81:MET:HE2	2.38	0.53
2:B:258:LYS:HD2	2:B:258:LYS:H	1.73	0.52
2:B:53:SER:O	2:B:54[A]:MET:HB3	2.09	0.52
1:A:2049:ILE:O	1:A:2053[A]:SER:OG	2.28	0.52
1:A:1962:ARG:HD3	1:A:1962:ARG:H	1.76	0.51
2:B:187:GLU:H	2:B:187:GLU:CD	2.21	0.49
1:A:1862:VAL:HG22	1:A:1872:THR:HG22	1.96	0.48
1:A:1946:VAL:O	1:A:1949:LEU:HG	2.14	0.48
1:A:1843:LEU:HA	1:A:1849:LYS:HD2	1.96	0.47
2:B:77:LEU:HD21	2:B:79:LYS:HE3	1.97	0.47
2:B:144:MET:HE2	2:B:176:LEU:HG	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:28:GLN:HG3	2:B:29:PRO:HD2	1.97	0.46
2:B:40:HIS:HD2	4:B:419:HOH:O	1.98	0.45
1:A:1920:LEU:HD13	1:A:1986:MET:HE1	1.98	0.45
1:A:2058:LEU:C	1:A:2058:LEU:HD23	2.42	0.45
2:B:1:MET:HB3	2:B:35:ASP:HA	2.00	0.44
2:B:237:TYR:CE2	2:B:241:LEU:HD11	2.53	0.43
2:B:28:GLN:CG	2:B:29:PRO:HD2	2.48	0.43
2:B:110:GLU:HG2	2:B:110:GLU:O	2.19	0.42
1:A:1974:LEU:O	1:A:1977:VAL:HG12	2.19	0.42
1:A:2062:GLU:OE1	1:A:2065:ARG:NH1	2.52	0.42
2:B:147:VAL:HG23	2:B:177[A]:ASN:HA	2.02	0.41
2:B:77:LEU:N	2:B:77:LEU:HD23	2.35	0.41
1:A:1857:VAL:HG21	1:A:1913:THR:OG1	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/258 (100%)	248 (96%)	9 (4%)	1 (0%)	30	15
2	B	315/308 (102%)	304 (96%)	10 (3%)	1 (0%)	36	21
All	All	573/566 (101%)	552 (96%)	19 (3%)	2 (0%)	30	21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2061	THR
2	B	106	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/233 (102%)	231 (98%)	6 (2%)	42	22
2	B	294/284 (104%)	293 (100%)	1 (0%)	86	83
All	All	531/517 (103%)	524 (99%)	7 (1%)	63	46

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1986	MET
1	A	1988	LEU
1	A	1991	ILE
1	A	2053[A]	SER
1	A	2053[B]	SER
1	A	2066	LYS
2	B	317	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1869	ASN
1	A	1947	HIS
2	B	2	ASN
2	B	23	ASN
2	B	47	GLN
2	B	135	ASN
2	B	150	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	5G1	A	2101	-	11,11,11	2.77	3 (27%)	14,14,14	2.23	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5G1	A	2101	-	-	4/6/6/6	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2101	5G1	C6-N	7.60	1.41	1.33
3	A	2101	5G1	C3-CL	-3.91	1.64	1.73
3	A	2101	5G1	C4-C3	-2.37	1.36	1.39

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2101	5G1	C6-N-N1	6.85	132.24	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2101	5G1	O-C6-N	-2.76	119.15	122.46
3	A	2101	5G1	C4-C3-CL	-2.39	117.49	121.01
3	A	2101	5G1	C4-C6-N	2.02	120.47	116.67

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2101	5G1	C4-C6-N-N1
3	A	2101	5G1	O-C6-N-N1
3	A	2101	5G1	C3-C4-C6-N
3	A	2101	5G1	C5-C4-C6-N

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/258 (91%)	2.05	94 (39%) 1 1	25, 70, 132, 202	12 (5%)
2	B	300/308 (97%)	1.88	119 (39%) 1 1	27, 72, 143, 207	9 (3%)
All	All	537/566 (94%)	1.95	213 (39%) 1 1	25, 71, 139, 207	21 (3%)

All (213) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1979[A]	MET	13.2
1	A	1878	CYS	7.7
1	A	2069	VAL	7.6
1	A	1855[A]	THR	7.2
1	A	2039	LEU	6.8
2	B	246	MET	6.5
1	A	2040	TRP	6.4
2	B	152	LEU	6.2
1	A	1834	ALA	6.1
2	B	101	MET	6.0
2	B	122[A]	GLN	5.9
1	A	2060	LEU	5.9
2	B	93	VAL	5.9
2	B	54[A]	MET	5.8
2	B	213	ILE	5.7
2	B	203[A]	TYR	5.7
1	A	2027	LEU	5.6
2	B	49	ALA	5.5
1	A	1988	LEU	5.3
2	B	316	LEU	5.3
2	B	297	PHE	5.1
1	A	2059	ILE	5.0
1	A	2063	TYR	4.9
2	B	108	ILE	4.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2017[A]	THR	4.8
2	B	104	SER	4.7
1	A	2038	HIS	4.7
2	B	170	SER	4.7
2	B	256	VAL	4.5
2	B	1	MET	4.3
1	A	1998	ARG	4.2
1	A	1873	LYS	4.2
1	A	1967	ALA	4.2
1	A	1995	TRP	4.2
2	B	3	THR	4.1
2	B	107	LYS	3.9
1	A	1866	PHE	3.9
2	B	22	PHE	3.9
1	A	1872	THR	3.8
2	B	275	LEU	3.8
1	A	1860	VAL	3.8
2	B	193	HIS	3.8
1	A	2067	TYR	3.8
2	B	180	VAL	3.8
1	A	1981	ALA	3.8
2	B	102	MET	3.7
2	B	284	LEU	3.7
2	B	129	ILE	3.7
1	A	1863	HIS	3.6
2	B	96	PHE	3.6
2	B	257	PRO	3.6
1	A	1987	VAL	3.6
1	A	2037	TYR	3.6
2	B	313	TYR	3.6
1	A	1989	PHE	3.6
2	B	111	ASP	3.6
2	B	294	TYR	3.6
1	A	1961[A]	LEU	3.5
1	A	2064	GLY	3.5
1	A	2051	ILE	3.5
2	B	173	ALA	3.5
2	B	97	LYS	3.5
2	B	109	ASP	3.4
1	A	2028	SER	3.4
1	A	2034	ILE	3.4
2	B	174	HIS	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	255	THR	3.4
2	B	276	PRO	3.4
2	B	280	SER	3.4
2	B	172	PRO	3.3
1	A	1840	TYR	3.3
2	B	279	TYR	3.3
1	A	1833	GLY	3.3
1	A	1843	LEU	3.3
2	B	41[A]	VAL	3.3
1	A	2048	TRP	3.2
1	A	2058	LEU	3.2
1	A	1844	PHE	3.2
1	A	1965	PHE	3.2
2	B	217	SER	3.2
1	A	1958	PRO	3.2
1	A	1999	ILE	3.1
1	A	1838	SER	3.1
1	A	1841	ALA	3.1
1	A	1955	ALA	3.1
2	B	100	GLN	3.1
2	B	103	VAL	3.1
1	A	1980[A]	LYS	3.0
2	B	126	ILE	3.0
1	A	2061	THR	3.0
2	B	130	VAL	3.0
2	B	260	MET	3.0
1	A	1951	PHE	3.0
1	A	2001[A]	SER	3.0
1	A	2053[A]	SER	3.0
2	B	259	HIS	3.0
2	B	261	LEU	3.0
1	A	1927	GLU	3.0
2	B	254	ALA	2.9
2	B	89	PHE	2.9
2	B	288	VAL	2.9
2	B	317	LEU	2.9
2	B	92	ILE	2.9
1	A	2056[A]	ARG	2.9
2	B	19	GLN	2.9
2	B	176	LEU	2.9
1	A	1837	SER	2.9
2	B	252	SER	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2041	PRO	2.8
1	A	1974	LEU	2.8
2	B	241	LEU	2.8
2	B	44[A]	ILE	2.8
2	B	31	HIS	2.8
1	A	1906	SER	2.8
1	A	1836	ASN	2.8
2	B	208	VAL	2.8
2	B	289	TRP	2.7
2	B	132	LYS	2.7
2	B	291	ILE	2.7
2	B	53	SER	2.7
1	A	1929	GLN	2.7
1	A	1950	ASP	2.7
2	B	293	LEU	2.7
2	B	302	LEU	2.7
1	A	1835	MET	2.7
1	A	1874	ALA	2.7
1	A	2065	ARG	2.7
2	B	204[A]	TYR	2.7
1	A	2068	ASN	2.7
2	B	258	LYS	2.7
1	A	2023	LYS	2.6
1	A	2050	THR	2.6
2	B	179	THR	2.6
1	A	2015	LEU	2.6
1	A	2049	ILE	2.6
1	A	2033	THR	2.6
2	B	249	LEU	2.6
1	A	1954	ILE	2.6
1	A	2025	ILE	2.6
2	B	171	ASP	2.5
2	B	214	PHE	2.5
1	A	2026	LEU	2.5
1	A	2024[A]	MET	2.5
2	B	134	GLU	2.5
2	B	38	ILE	2.5
1	A	2030	PRO	2.5
1	A	1931[A]	LYS	2.5
1	A	1972	ASP	2.5
1	A	1905	LEU	2.5
2	B	6	PHE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	199	LEU	2.5
2	B	121	VAL	2.5
1	A	1868	GLY	2.4
2	B	267	ILE	2.4
1	A	1953	ASN	2.4
2	B	23	ASN	2.4
2	B	95	ASN	2.4
1	A	1994	ASP	2.4
2	B	29	PRO	2.4
1	A	2043	PHE	2.4
2	B	233	PHE	2.4
1	A	1968	ALA	2.4
2	B	80	MET	2.4
2	B	28	GLN	2.4
2	B	216	ASN	2.4
2	B	282	ILE	2.4
2	B	147	VAL	2.4
2	B	64[A]	MET	2.4
1	A	1920	LEU	2.3
1	A	1963	LEU	2.3
1	A	1890	PHE	2.3
1	A	1986	MET	2.3
2	B	234	PHE	2.3
2	B	142	SER	2.3
2	B	5	PRO	2.3
2	B	106	PRO	2.3
2	B	136	GLN	2.3
1	A	2066	LYS	2.3
2	B	181	ILE	2.3
2	B	272	ILE	2.3
2	B	314	PRO	2.2
2	B	84	ARG	2.2
2	B	131	ARG	2.2
2	B	206	ASN	2.2
2	B	52	SER	2.2
2	B	143	SER	2.2
2	B	30	PHE	2.2
2	B	60	PHE	2.2
2	B	237	TYR	2.2
1	A	1893	ILE	2.2
2	B	218	SER	2.2
2	B	307	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2011	LEU	2.2
2	B	287	ARG	2.2
2	B	74	LYS	2.2
2	B	87	ALA	2.2
1	A	1949	LEU	2.2
2	B	269	TYR	2.2
2	B	33	ILE	2.2
1	A	1861	THR	2.2
1	A	2022	ALA	2.1
1	A	1944	LEU	2.1
2	B	285	ASN	2.1
2	B	290	ASN	2.1
2	B	73	PRO	2.1
2	B	146	THR	2.1
1	A	1956	ILE	2.1
2	B	62	CYS	2.1
2	B	264	LEU	2.1
2	B	81	MET	2.1
2	B	94	HIS	2.1
1	A	1901	GLY	2.0
1	A	1969	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	5G1	A	2101	11/11	0.67	0.23	20,20,20,20	11

6.5 Other polymers [i](#)

There are no such residues in this entry.