



Full wwPDB NMR Structure Validation Report ⓘ

Mar 24, 2026 – 02:58 AM UTC

PDB ID : 1FMH / pdb_00001fmh
Title : NMR SOLUTION STRUCTURE OF A DESIGNED HETERODIMERIC
LEUCINE ZIPPER
Authors : Marti, D.N.; Jelesarov, I.; Bosshard, H.R.
Deposited on : 2000-08-17

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	33	
2	B	33	

2 Ensemble composition and analysis

This entry contains 25 models. Model 14 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:27, B:1-B:29 (56)	0.42	14

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 7, 8, 9, 10, 12, 13, 14, 15, 17, 18, 20, 23, 24, 25
2	16, 21, 22
3	11, 19
Single-model clusters	6

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1005 atoms, of which 505 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called GENERAL CONTROL PROTEIN GCN4.

Mol	Chain	Residues	Atoms						Trace
1	A	33	Total	C	H	N	O	S	1
			475	150	226	41	57	1	

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLU	ARG	engineered mutation	UNP P03069
A	2	VAL	MET	engineered mutation	UNP P03069
A	3	ALA	LYS	engineered mutation	UNP P03069
A	7	LYS	ASP	engineered mutation	UNP P03069
A	8	GLU	LYS	engineered mutation	UNP P03069
A	10	ALA	GLU	engineered mutation	UNP P03069
A	11	GLN	GLU	engineered mutation	UNP P03069
A	12	ALA	LEU	engineered mutation	UNP P03069
A	13	GLU	LEU	engineered mutation	UNP P03069
A	14	ALA	SER	engineered mutation	UNP P03069
A	15	GLU	LYS	engineered mutation	UNP P03069
A	18	GLN	HIS	engineered mutation	UNP P03069
A	21	GLN	ASN	engineered mutation	UNP P03069
A	25	GLN	ARG	engineered mutation	UNP P03069
A	27	GLU	LYS	engineered mutation	UNP P03069
A	28	HIS	LYS	engineered mutation	UNP P03069
A	29	GLU	LEU	engineered mutation	UNP P03069
A	30	CYS	VAL	engineered mutation	UNP P03069

- Molecule 2 is a protein called GENERAL CONTROL PROTEIN GCN4.

Mol	Chain	Residues	Atoms						Trace
2	B	33	Total	C	H	N	O	S	1
			530	156	279	54	40	1	

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	GLU	ARG	engineered mutation	UNP P03069
B	2	VAL	MET	engineered mutation	UNP P03069

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	3	GLN	LYS	engineered mutation	UNP P03069
B	4	ALA	GLN	engineered mutation	UNP P03069
B	6	LYS	GLU	engineered mutation	UNP P03069
B	7	LYS	ASP	engineered mutation	UNP P03069
B	8	ARG	LYS	engineered mutation	UNP P03069
B	10	GLN	GLU	engineered mutation	UNP P03069
B	11	ALA	GLU	engineered mutation	UNP P03069
B	13	LYS	LEU	engineered mutation	UNP P03069
B	14	ALA	SER	engineered mutation	UNP P03069
B	15	ARG	LYS	engineered mutation	UNP P03069
B	18	ALA	HIS	engineered mutation	UNP P03069
B	19	ALA	LEU	engineered mutation	UNP P03069
B	20	LYS	GLU	engineered mutation	UNP P03069
B	21	GLN	ASN	engineered mutation	UNP P03069
B	22	LYS	GLU	engineered mutation	UNP P03069
B	24	GLN	ALA	engineered mutation	UNP P03069
B	25	ALA	ARG	engineered mutation	UNP P03069
B	27	ARG	LYS	engineered mutation	UNP P03069
B	28	HIS	LYS	engineered mutation	UNP P03069
B	29	LYS	LEU	engineered mutation	UNP P03069
B	30	CYS	VAL	engineered mutation	UNP P03069

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: GENERAL CONTROL PROTEIN GCN4



- Molecule 2: GENERAL CONTROL PROTEIN GCN4



4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: GENERAL CONTROL PROTEIN GCN4



- Molecule 2: GENERAL CONTROL PROTEIN GCN4



4.2.2 Score per residue for model 2

- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  70% 12% 18%



- Molecule 2: GENERAL CONTROL PROTEIN GCN4


Chain B:  67% 21% 12%



4.2.3 Score per residue for model 3


- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  73% 9% 18%



- Molecule 2: GENERAL CONTROL PROTEIN GCN4

Chain B:  45% 42% 12%



4.2.4 Score per residue for model 4

- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  64% 18% 18%



- Molecule 2: GENERAL CONTROL PROTEIN GCN4

Chain B:  58% 30% 12%




4.2.5 Score per residue for model 5

- Molecule 1: GENERAL CONTROL PROTEIN GCN4



- Molecule 2: GENERAL CONTROL PROTEIN GCN4




4.2.6 Score per residue for model 6

- Molecule 1: GENERAL CONTROL PROTEIN GCN4




- Molecule 2: GENERAL CONTROL PROTEIN GCN4




4.2.7 Score per residue for model 7

- Molecule 1: GENERAL CONTROL PROTEIN GCN4




- Molecule 2: GENERAL CONTROL PROTEIN GCN4



4.2.8 Score per residue for model 8

- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  73% 9% 18%



- Molecule 2: GENERAL CONTROL PROTEIN GCN4

Chain B:  58% 30% 12%



4.2.9 Score per residue for model 9


- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  61% 18% 18%



- Molecule 2: GENERAL CONTROL PROTEIN GCN4

Chain B:  42% 45% 12%



4.2.10 Score per residue for model 10

- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  64% 18% 18%




- Molecule 2: GENERAL CONTROL PROTEIN GCN4

Chain B:  64% 24% 12%




4.2.11 Score per residue for model 11

- Molecule 1: GENERAL CONTROL PROTEIN GCN4



- Molecule 2: GENERAL CONTROL PROTEIN GCN4




4.2.12 Score per residue for model 12

- Molecule 1: GENERAL CONTROL PROTEIN GCN4




- Molecule 2: GENERAL CONTROL PROTEIN GCN4




4.2.13 Score per residue for model 13

- Molecule 1: GENERAL CONTROL PROTEIN GCN4



- Molecule 2: GENERAL CONTROL PROTEIN GCN4



4.2.14 Score per residue for model 14 (medoid)

- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  70% 12% 18%



- Molecule 2: GENERAL CONTROL PROTEIN GCN4

Chain B:  58% 30% 12%



4.2.15 Score per residue for model 15


- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  64% 18% 18%



- Molecule 2: GENERAL CONTROL PROTEIN GCN4

Chain B:  76% 12% 12%



4.2.16 Score per residue for model 16

- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  64% 18% 18%




- Molecule 2: GENERAL CONTROL PROTEIN GCN4

Chain B:  61% 27% 12%




4.2.17 Score per residue for model 17

- Molecule 1: GENERAL CONTROL PROTEIN GCN4




- Molecule 2: GENERAL CONTROL PROTEIN GCN4




4.2.18 Score per residue for model 18

- Molecule 1: GENERAL CONTROL PROTEIN GCN4




- Molecule 2: GENERAL CONTROL PROTEIN GCN4



4.2.19 Score per residue for model 19

- Molecule 1: GENERAL CONTROL PROTEIN GCN4




- Molecule 2: GENERAL CONTROL PROTEIN GCN4



4.2.20 Score per residue for model 20

- Molecule 1: GENERAL CONTROL PROTEIN GCN4




- Molecule 2: GENERAL CONTROL PROTEIN GCN4




4.2.21 Score per residue for model 21

- Molecule 1: GENERAL CONTROL PROTEIN GCN4




- Molecule 2: GENERAL CONTROL PROTEIN GCN4




4.2.22 Score per residue for model 22

- Molecule 1: GENERAL CONTROL PROTEIN GCN4




- Molecule 2: GENERAL CONTROL PROTEIN GCN4



4.2.23 Score per residue for model 23


- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  73% 9% 18%



- Molecule 2: GENERAL CONTROL PROTEIN GCN4

Chain B:  67% 18% 12%



4.2.24 Score per residue for model 24

- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  67% 15% 18%



- Molecule 2: GENERAL CONTROL PROTEIN GCN4

Chain B:  61% 24% 12%



4.2.25 Score per residue for model 25

- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A:  64% 18% 18%



- Molecule 2: GENERAL CONTROL PROTEIN GCN4

Chain B:  73% 15% 12%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *1. DISTANCE GEOMETRY/REGULARIZATION, 2. SIMULATED ANNEALING, 3. MOLECULAR DYNAMICS SIMULATION INCLUDING 8 ANGSTROMS WATER SHELL.*

Of the 50 calculated structures, 25 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.851
X-PLOR	refinement	3.851

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.38±0.02	0±0/217 (0.0± 0.0%)	0.77±0.02	0±0/294 (0.0± 0.0%)
2	B	0.44±0.04	0±0/239 (0.0± 0.0%)	0.82±0.06	0±1/317 (0.1± 0.2%)
All	All	0.42	0/11400 (0.0%)	0.80	5/15275 (0.0%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	28	HIS	CA-CB-CG	5.89	119.69	113.80	17	3
2	B	28	HIS	CB-CA-C	-5.16	102.74	110.90	5	2

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	216	201	201	4±2
2	B	237	267	267	5±2
All	All	11325	11700	11700	188

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 8.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:1:GLU:HB3	1:A:4:GLN:HG3	0.87	1.46	1	1
1:A:1:GLU:HB3	1:A:4:GLN:HG2	0.78	1.56	12	9
2:B:1:GLU:HB3	2:B:4:ALA:HB3	0.75	1.56	17	2
2:B:25:ALA:HA	2:B:28:HIS:CD2	0.69	2.22	17	3
1:A:15:GLU:O	1:A:19:LEU:HG	0.69	1.87	21	25
1:A:23:VAL:HG13	2:B:26:LEU:HD11	0.67	1.66	24	5
1:A:17:TYR:O	1:A:21:GLN:HG2	0.65	1.92	1	1
2:B:22:LYS:O	2:B:26:LEU:HG	0.58	1.98	8	14
1:A:15:GLU:O	1:A:18:GLN:HG3	0.57	2.00	9	1
2:B:21:GLN:HA	2:B:24:GLN:HG2	0.56	1.78	9	2
2:B:1:GLU:HG3	2:B:4:ALA:H	0.56	1.59	18	7
1:A:26:LEU:HB3	2:B:26:LEU:HB3	0.55	1.77	22	7
2:B:17:TYR:O	2:B:21:GLN:HG2	0.54	2.02	8	6
1:A:26:LEU:CD1	2:B:27:ARG:HG3	0.52	2.35	9	6
1:A:15:GLU:HB3	2:B:16:ASN:ND2	0.52	2.19	17	1
1:A:15:GLU:HB2	2:B:16:ASN:ND2	0.51	2.19	6	1
1:A:2:VAL:O	1:A:6:GLU:HG3	0.51	2.06	10	1
1:A:4:GLN:O	1:A:7:LYS:HG2	0.50	2.07	4	1
1:A:22:GLU:O	1:A:26:LEU:HG	0.50	2.05	25	3
2:B:9:VAL:O	2:B:13:LYS:HG2	0.49	2.08	3	2
2:B:1:GLU:HG2	2:B:4:ALA:HB2	0.49	1.83	13	2
1:A:26:LEU:HD11	2:B:27:ARG:HD2	0.49	1.85	23	1
2:B:19:ALA:O	2:B:23:VAL:HG23	0.48	2.07	3	16
1:A:26:LEU:HD11	2:B:27:ARG:HG3	0.48	1.86	22	3
2:B:3:GLN:O	2:B:7:LYS:HG2	0.47	2.09	23	7
1:A:1:GLU:HB3	1:A:4:GLN:CG	0.47	2.39	17	2
2:B:9:VAL:O	2:B:13:LYS:HD3	0.47	2.09	22	5
2:B:1:GLU:HB3	2:B:4:ALA:CB	0.47	2.36	17	2
1:A:12:ALA:HB1	2:B:12:LEU:HB3	0.46	1.87	14	6
1:A:12:ALA:HB3	2:B:12:LEU:HD13	0.46	1.87	5	5
1:A:13:GLU:OE2	2:B:12:LEU:HD21	0.46	2.10	20	1
2:B:9:VAL:HG13	2:B:13:LYS:HE3	0.46	1.85	19	1
1:A:26:LEU:HD13	2:B:26:LEU:HB2	0.46	1.88	8	4
2:B:3:GLN:HE21	2:B:7:LYS:HD3	0.46	1.71	17	1
2:B:5:LEU:O	2:B:9:VAL:HG23	0.45	2.12	19	6
2:B:2:VAL:O	2:B:6:LYS:HG3	0.44	2.12	2	8
1:A:22:GLU:HG2	2:B:23:VAL:HG13	0.44	1.90	16	1
2:B:3:GLN:NE2	2:B:7:LYS:HD3	0.44	2.27	17	1
2:B:18:ALA:O	2:B:21:GLN:HG2	0.43	2.13	9	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:17:TYR:O	1:A:21:GLN:HG3	0.43	2.13	5	1
1:A:1:GLU:HA	1:A:1:GLU:OE1	0.43	2.13	10	1
2:B:21:GLN:HA	2:B:24:GLN:NE2	0.43	2.29	9	1
2:B:1:GLU:HG2	2:B:4:ALA:CB	0.43	2.44	13	1
1:A:19:LEU:O	1:A:23:VAL:HG23	0.42	2.15	24	1
1:A:14:ALA:O	1:A:18:GLN:HG2	0.42	2.14	12	1
2:B:7:LYS:N	2:B:7:LYS:HD2	0.42	2.30	1	1
1:A:16:ASN:O	1:A:20:GLU:HG3	0.41	2.14	17	1
1:A:5:LEU:O	1:A:9:VAL:HG23	0.41	2.16	5	1
2:B:16:ASN:O	2:B:20:LYS:HB2	0.41	2.15	4	1
2:B:6:LYS:O	2:B:10:GLN:HG2	0.41	2.14	5	1
1:A:3:ALA:O	1:A:7:LYS:HG3	0.41	2.16	6	1
2:B:21:GLN:O	2:B:24:GLN:HG2	0.41	2.16	6	1
1:A:26:LEU:HD13	2:B:27:ARG:HG3	0.41	1.91	9	1
1:A:15:GLU:HB3	2:B:16:ASN:HD21	0.41	1.75	17	1
2:B:1:GLU:O	2:B:5:LEU:HG	0.41	2.16	16	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	27/33 (82%)	27±0 (100±1%)	0±0 (0±1%)	0±0 (0±0%)	100	100
2	B	29/33 (88%)	29±0 (100±1%)	0±0 (0±1%)	0±0 (0±0%)	100	100
All	All	1400/1650 (85%)	1397 (100%)	3 (0%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	22/25 (88%)	22±0 (100±1%)	0±0 (0±1%)	81	97
2	B	23/24 (96%)	22±1 (96±4%)	1±1 (4±4%)	29	79
All	All	1125/1225 (92%)	1100 (98%)	25 (2%)	45	90

All 9 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	22	LYS	5
2	B	21	GLN	5
2	B	1	GLU	4
2	B	27	ARG	3
2	B	7	LYS	3
2	B	3	GLN	2
1	A	18	GLN	1
2	B	15	ARG	1
1	A	25	GLN	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry ⓘ

There are no ligands in this entry.

6.7 Other polymers ⓘ

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided