



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:11 PM UTC

PDB ID : 6FMD / pdb_00006fmd
Title : Targeting myeloid differentiation using potent human dihydroorotate dehydrogenase (hDHODH) inhibitors based on 2-hydroxypyrazolo[1,5-a]pyridine scaffold
Authors : Goyal, P.; Jarva, M.; Andersson, M.; Lolli, M.L.; Friemann, R.
Deposited on : 2018-01-30
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	A	407	-	-	X	-

2 Entry composition [i](#)


There are 9 unique types of molecules in this entry. The entry contains 3128 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydroorotate dehydrogenase (quinone), mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	357	Total	C	N	O	S	0	9	0
			2792	1750	516	522	4			

- Molecule 2 is 2-oxidanyl- {N}-[2,3,5,6-tetrakis(fluoranyl)-4-phenyl-phenyl]pyrazolo[1,5-a]pyridine-3-carboxamide (CCD ID: DUH) (formula: C₂₀H₁₁F₄N₃O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			29	20	4	3	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		

- Molecule 4 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula: $C_{17}H_{21}N_4O_9P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is OROTIC ACID (CCD ID: ORO) (formula: $C_5H_4N_2O_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			11	5	2	4		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			10	6	4		
8	A	1	Total	C	O	0	0
			10	6	4		
8	A	1	Total	C	O	0	0
			10	6	4		
8	A	1	Total	C	O	0	0
			10	6	4		
8	A	1	Total	C	O	0	0
			10	6	4		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	155	Total	O	0	0
			155	155		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydroorotate dehydrogenase (quinone), mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	90.44Å 90.44Å 122.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	78.32 – 1.58 78.32 – 1.58	Depositor EDS
% Data completeness (in resolution range)	99.5 (78.32-1.58) 99.5 (78.32-1.58)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 1.58Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.137 , 0.160 (Not available) , 0.165	Depositor DCC
R_{free} test set	4016 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtriage
Anisotropy	0.167	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3128	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, GOL, PGE, FMN, CL, DUH, ORO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/2846 (0.0%)	0.86	0/3843

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	212	ASN	N-CA	6.06	1.53	1.46

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2792	0	2851	13	0
2	A	29	0	0	0	0
3	A	2	0	0	0	0
4	A	31	0	19	0	0
5	A	24	0	18	3	0
6	A	11	0	3	0	0
7	A	24	0	32	0	0
8	A	60	0	84	2	0
9	A	155	0	0	0	0
All	All	3128	0	3007	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:ARG:NH2	1:A:249:ARG:HD2	1.67	1.10
1:A:246:ARG:CZ	1:A:249:ARG:HD2	1.88	1.04
1:A:246:ARG:HD2	5:A:407:ACT:OXT	1.74	0.86
1:A:246:ARG:HD2	5:A:407:ACT:C	2.15	0.77
1:A:246:ARG:NH2	1:A:249:ARG:CD	2.53	0.65
1:A:246:ARG:NH1	1:A:249:ARG:HD2	2.22	0.52
1:A:246:ARG:NE	1:A:246:ARG:HA	2.26	0.49
1:A:247:VAL:HG23	1:A:248:HIS:CD2	2.51	0.46
1:A:235:THR:O	1:A:239:GLN:HG3	2.16	0.44
1:A:241:ARG:HD2	1:A:241:ARG:C	2.43	0.42
1:A:57:ARG:HD3	8:A:418:PGE:H4	2.04	0.40
1:A:57:ARG:HD3	8:A:418:PGE:C4	2.52	0.40
1:A:207:ASP:OD1	5:A:405:ACT:O	2.40	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	362/395 (92%)	354 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/316 (94%)	296 (100%)	1 (0%)	86	79

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	188	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	141	GLN
1	A	239	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 2 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	PGE	A	417	-	9,9,9	0.52	0	8,8,8	0.35	0
5	ACT	A	406	-	3,3,3	0.76	0	3,3,3	1.19	0
7	GOL	A	415	-	5,5,5	0.39	0	5,5,5	0.43	0
2	DUH	A	401	-	32,32,32	3.44	17 (53%)	41,47,47	1.73	14 (34%)
5	ACT	A	408	-	3,3,3	1.00	0	3,3,3	1.05	0
5	ACT	A	409	-	3,3,3	0.97	0	3,3,3	0.70	0
8	PGE	A	421	-	9,9,9	0.66	0	8,8,8	0.71	0
7	GOL	A	414	-	5,5,5	0.42	0	5,5,5	0.43	0
8	PGE	A	416	-	9,9,9	0.57	0	8,8,8	0.48	0
5	ACT	A	410	-	3,3,3	1.15	0	3,3,3	0.34	0
8	PGE	A	418	-	9,9,9	0.57	0	8,8,8	0.47	0
5	ACT	A	407	-	3,3,3	1.12	0	3,3,3	0.65	0
6	ORO	A	411	-	11,11,11	0.96	0	14,15,15	2.48	7 (50%)
7	GOL	A	413	-	5,5,5	0.59	0	5,5,5	0.82	0
4	FMN	A	404	-	33,33,33	1.16	3 (9%)	48,50,50	1.22	4 (8%)
5	ACT	A	405	-	3,3,3	0.99	0	3,3,3	1.00	0
7	GOL	A	412	-	5,5,5	0.25	0	5,5,5	0.33	0
8	PGE	A	419	-	9,9,9	0.73	0	8,8,8	0.64	0
8	PGE	A	420	-	9,9,9	0.41	0	8,8,8	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	413	-	-	2/4/4/4	-
7	GOL	A	415	-	-	2/4/4/4	-
2	DUH	A	401	-	-	0/12/12/12	0/4/4/4
8	PGE	A	418	-	-	2/7/7/7	-
4	FMN	A	404	-	-	5/18/18/18	0/3/3/3
8	PGE	A	417	-	-	3/7/7/7	-
8	PGE	A	420	-	-	4/7/7/7	-
8	PGE	A	421	-	-	4/7/7/7	-
7	GOL	A	414	-	-	2/4/4/4	-
7	GOL	A	412	-	-	2/4/4/4	-
8	PGE	A	416	-	-	3/7/7/7	-
8	PGE	A	419	-	-	2/7/7/7	-
6	ORO	A	411	-	-	4/4/4/4	0/1/1/1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	DUH	C2-C3	7.78	1.52	1.38
2	A	401	DUH	C12-C11	6.84	1.48	1.38
2	A	401	DUH	C10-C15	6.30	1.48	1.39
2	A	401	DUH	C5-C4	5.89	1.51	1.39
2	A	401	DUH	C7-C6	5.80	1.51	1.38
2	A	401	DUH	C15-C14	-5.37	1.29	1.38
2	A	401	DUH	C13-C14	5.07	1.47	1.39
2	A	401	DUH	C22-N20	3.77	1.43	1.35
2	A	401	DUH	C10-C11	-3.73	1.33	1.39
2	A	401	DUH	C25-C29	-3.33	1.34	1.41
2	A	401	DUH	N28-N27	3.16	1.40	1.37
2	A	401	DUH	C13-C12	-2.92	1.34	1.39
2	A	401	DUH	C33-C32	2.63	1.40	1.35
2	A	401	DUH	C31-C30	2.61	1.43	1.37
2	A	401	DUH	C33-N28	2.56	1.42	1.37
4	A	404	FMN	C4A-N5	2.48	1.36	1.30
2	A	401	DUH	C6-C5	-2.42	1.34	1.38
4	A	404	FMN	C4-N3	-2.40	1.34	1.38
2	A	401	DUH	C7-C2	-2.40	1.33	1.38
4	A	404	FMN	C9A-C5A	2.15	1.44	1.41

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	411	ORO	C4-N3-C2	-4.54	121.15	125.55
6	A	411	ORO	C6-N1-C2	-4.11	118.26	122.65
6	A	411	ORO	C5-C4-N3	3.80	119.96	115.24
4	A	404	FMN	C4-C4A-N5	3.29	122.75	118.21
2	A	401	DUH	C2-C3-C4	-3.26	116.83	120.54
6	A	411	ORO	O4-C4-C5	-3.08	121.20	125.46
2	A	401	DUH	O23-C22-C25	3.02	126.05	121.53
6	A	411	ORO	N3-C2-N1	2.82	120.18	115.74
2	A	401	DUH	C30-C29-N28	2.82	120.94	118.15
4	A	404	FMN	O2-C2-N1	-2.73	117.26	121.80
2	A	401	DUH	O34-C26-N27	-2.73	118.48	122.94
2	A	401	DUH	C3-C4-C10	-2.73	116.12	120.80
2	A	401	DUH	C15-C10-C11	2.71	119.81	116.05
2	A	401	DUH	C29-N28-N27	2.67	115.12	112.94
2	A	401	DUH	C5-C4-C3	2.67	122.45	117.68
2	A	401	DUH	C4-C10-C15	-2.43	118.02	122.12
2	A	401	DUH	C14-C13-C12	2.42	120.21	115.99
6	A	411	ORO	C5-C6-N1	2.37	123.65	120.87

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	404	FMN	O4-C4-C4A	-2.27	120.55	126.53
2	A	401	DUH	C10-C11-C12	-2.25	119.29	121.99
2	A	401	DUH	F16-C12-C13	2.25	122.87	119.54
2	A	401	DUH	C6-C5-C4	-2.23	118.00	120.54
2	A	401	DUH	F17-C11-C10	2.06	122.56	119.78
6	A	411	ORO	C6-C5-C4	-2.00	116.79	120.34
4	A	404	FMN	C5A-C9A-N10	2.00	119.78	117.97

There are no chirality outliers.

All (35) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	411	ORO	N1-C6-C7-O71
6	A	411	ORO	N1-C6-C7-O72
6	A	411	ORO	C5-C6-C7-O72
7	A	412	GOL	O1-C1-C2-O2
7	A	412	GOL	O1-C1-C2-C3
7	A	413	GOL	O1-C1-C2-C3
7	A	414	GOL	C1-C2-C3-O3
8	A	418	PGE	O2-C3-C4-O3
8	A	421	PGE	O2-C3-C4-O3
8	A	418	PGE	O1-C1-C2-O2
8	A	420	PGE	O1-C1-C2-O2
8	A	416	PGE	O3-C5-C6-O4
8	A	419	PGE	O3-C5-C6-O4
8	A	421	PGE	O3-C5-C6-O4
7	A	413	GOL	O1-C1-C2-O2
4	A	404	FMN	C2'-C3'-C4'-O4'
8	A	417	PGE	O1-C1-C2-O2
8	A	417	PGE	O3-C5-C6-O4
8	A	420	PGE	O2-C3-C4-O3
7	A	414	GOL	O2-C2-C3-O3
6	A	411	ORO	C5-C6-C7-O71
7	A	415	GOL	O1-C1-C2-C3
8	A	421	PGE	C4-C3-O2-C2
8	A	419	PGE	C1-C2-O2-C3
8	A	420	PGE	C3-C4-O3-C5
7	A	415	GOL	O1-C1-C2-O2
4	A	404	FMN	O3'-C3'-C4'-O4'
4	A	404	FMN	C4'-C5'-O5'-P
4	A	404	FMN	C5'-O5'-P-O1P
8	A	416	PGE	C1-C2-O2-C3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
8	A	420	PGE	O3-C5-C6-O4
8	A	417	PGE	C4-C3-O2-C2
8	A	421	PGE	C1-C2-O2-C3
8	A	416	PGE	O1-C1-C2-O2
4	A	404	FMN	O3'-C3'-C4'-C5'


There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	418	PGE	2	0
5	A	407	ACT	2	0
5	A	405	ACT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	357/395 (90%)	-0.21	13 (3%) 46 57	7, 20, 48, 89	9 (2%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	216	PRO	5.8
1	A	31	ALA	4.4
1	A	73	PHE	3.9
1	A	189	VAL	3.3
1	A	226	GLY	2.9
1	A	396	ARG	2.7
1	A	246	ARG	2.7
1	A	215	SER	2.6
1	A	213	VAL	2.3
1	A	32	THR	2.3
1	A	71	ALA	2.2
1	A	249	ARG	2.1
1	A	362	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	ACT	A	407	4/4	0.67	0.27	61,69,75,87	0
5	ACT	A	409	4/4	0.71	0.22	66,70,70,74	0
5	ACT	A	410	4/4	0.78	0.17	48,52,55,58	0
8	PGE	A	417	10/10	0.83	0.18	58,65,84,86	0
7	GOL	A	414	6/6	0.84	0.22	54,76,79,81	0
8	PGE	A	416	10/10	0.87	0.18	32,67,79,79	0
8	PGE	A	418	10/10	0.87	0.17	63,68,73,77	0
8	PGE	A	420	10/10	0.87	0.16	57,63,67,71	0
7	GOL	A	413	6/6	0.88	0.14	56,56,65,68	0
8	PGE	A	419	10/10	0.90	0.13	44,54,59,60	0
7	GOL	A	412	6/6	0.90	0.12	58,62,66,79	0
8	PGE	A	421	10/10	0.90	0.15	35,57,66,74	0
5	ACT	A	408	4/4	0.91	0.12	34,35,38,40	0
7	GOL	A	415	6/6	0.91	0.15	61,62,68,69	0
5	ACT	A	406	4/4	0.92	0.14	34,44,48,49	0
5	ACT	A	405	4/4	0.95	0.08	24,25,26,27	0
4	FMN	A	404	31/31	0.99	0.03	11,13,15,15	0
6	ORO	A	411	11/11	0.99	0.02	14,15,17,18	0
2	DUH	A	401	29/29	0.99	0.04	16,20,26,29	0
3	CL	A	402	1/1	1.00	0.02	26,26,26,26	0
3	CL	A	403	1/1	1.00	0.06	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FMN A 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.