



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 04:54 PM UTC

PDB ID : 4FNL / pdb_00004fnl
Title : Crystal structure of broadly neutralizing antibody C05
Authors : Ekiert, D.C.; Wilson, I.A.
Deposited on : 2012-06-19
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	247	8% 86% 7% 6%
1	I	247	8% 87% 7% 5%
2	L	214	6% 86% 12% .
2	M	214	4% 86% 11% .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7360 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Antibody C05, Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	233	Total	C	N	O	S	0	4	0
			1766	1104	299	355	8			
1	I	234	Total	C	N	O	S	0	3	0
			1771	1107	301	355	8			

- Molecule 2 is a protein called Antibody C05, Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	213	Total	C	N	O	S	0	1	0
			1648	1033	281	330	4			
2	M	213	Total	C	N	O	S	0	1	0
			1648	1033	281	330	4			

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	H	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	154	Total	O	0	0
			154	154		
4	L	142	Total	O	0	0
			142	142		
4	I	100	Total	O	0	0
			100	100		
4	M	123	Total	O	0	0
			123	123		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Antibody C05, Heavy Chain




- Molecule 1: Antibody C05, Heavy Chain



- Molecule 2: Antibody C05, Light Chain



- Molecule 2: Antibody C05, Light Chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	160.13Å 160.13Å 135.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.22 – 2.30 49.22 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.22-2.30) 99.3 (49.22-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.19	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.2_869)	Depositor
R, R_{free}	0.212 , 0.240 0.202 , 0.231	Depositor DCC
R_{free} test set	3950 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	40.9	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7360	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.29	0/1796	0.67	0/2443
1	I	0.28	0/1801	0.71	0/2450
2	L	0.29	0/1682	0.68	0/2280
2	M	0.28	0/1682	0.68	0/2280
All	All	0.29	0/6961	0.69	0/9453

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1766	0	1709	12	0
1	I	1771	0	1714	10	0
2	L	1648	0	1614	17	0
2	M	1648	0	1614	19	0
3	H	8	0	12	0	0
4	H	154	0	0	1	0
4	I	100	0	0	0	0
4	L	142	0	0	0	0
4	M	123	0	0	2	0
All	All	7360	0	6663	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:4:LEU:HD13	1:H:92:CYS:SG	2.41	0.61
1:I:4:LEU:HD13	1:I:92:CYS:SG	2.42	0.59
2:M:187:GLU:O	2:M:211:ARG:NH2	2.35	0.59
1:I:178:LEU:C	1:I:178:LEU:HD23	2.27	0.59
1:H:178:LEU:C	1:H:178:LEU:HD23	2.29	0.58
1:I:19[B]:ARG:NH2	1:I:79:TYR:CE1	2.72	0.58
2:L:11:LEU:C	2:L:11:LEU:HD12	2.29	0.57
1:I:128:SER:O	1:I:129:LYS:HB3	2.04	0.57
2:M:11:LEU:HD12	2:M:11:LEU:C	2.29	0.57
1:I:97:SER:OG	1:I:100(N):ASP:OD2	2.21	0.57
2:L:90:GLN:OE1	2:L:92:ASP:N	2.37	0.57
2:M:11:LEU:HD21	2:M:19:VAL:HG13	1.86	0.56
2:L:11:LEU:HD21	2:L:19:VAL:HG13	1.87	0.56
1:I:185:PRO:O	1:I:186:SER:C	2.50	0.55
1:H:128:SER:O	1:H:129:LYS:HB3	2.09	0.53
2:L:150:VAL:HG13	2:L:192:TYR:CE1	2.45	0.52
2:L:55:GLN:HE21	2:L:55:GLN:HA	1.75	0.52
1:I:183:THR:HG21	2:M:137:ASN:ND2	2.26	0.50
2:M:181:LEU:CD1	2:M:186:TYR:HB2	2.42	0.50
2:L:90:GLN:CD	2:L:90:GLN:C	2.80	0.49
2:M:90:GLN:CD	2:M:90:GLN:C	2.81	0.49
2:M:55:GLN:HE21	2:M:55:GLN:HA	1.78	0.48
1:I:145:TYR:CE2	1:I:150:VAL:HG13	2.49	0.48
1:H:97:SER:OG	1:H:100(N):ASP:OD2	2.24	0.47
1:H:183:THR:HG21	2:L:137:ASN:ND2	2.30	0.47
2:L:188:LYS:HB3	2:L:189:HIS:CD2	2.49	0.47
2:L:37:GLN:HB2	2:L:47:LEU:HD11	1.96	0.47
2:L:83:VAL:CG1	2:L:106:ILE:HG12	2.45	0.47
2:M:56:ARG:NE	4:M:390:HOH:O	2.48	0.46
1:H:185:PRO:O	1:H:186:SER:C	2.59	0.46
2:L:181:LEU:CD1	2:L:186:TYR:HB2	2.47	0.45
2:L:33:LEU:HD22	2:L:71:PHE:CG	2.51	0.45
1:I:128:SER:O	1:I:129:LYS:CB	2.64	0.45
2:L:145:LYS:HB3	2:L:197:THR:HB	1.98	0.45
2:M:83:VAL:CG1	2:M:106:ILE:HG12	2.47	0.45
2:M:55:GLN:NE2	4:M:380:HOH:O	2.49	0.44
2:M:140:TYR:CG	2:M:141:PRO:HA	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:184:VAL:O	1:H:185:PRO:C	2.61	0.43
1:H:82(B)[A]:ASN:ND2	4:H:544:HOH:O	2.51	0.43
2:L:140:TYR:CG	2:L:141:PRO:HA	2.53	0.43
2:M:188:LYS:HB3	2:M:189:HIS:CD2	2.54	0.43
1:H:129:LYS:O	1:H:129:LYS:HG2	2.19	0.43
2:M:33:LEU:HD22	2:M:71:PHE:CG	2.54	0.43
2:M:150:VAL:HG13	2:M:192:TYR:CE1	2.54	0.43
2:M:125:LEU:C	2:M:127:SER:N	2.77	0.42
1:H:128:SER:O	1:H:129:LYS:CB	2.67	0.42
2:M:175:LEU:C	2:M:175:LEU:HD23	2.45	0.42
2:M:154:LEU:C	2:M:154:LEU:HD23	2.45	0.42
2:L:154:LEU:C	2:L:154:LEU:HD23	2.45	0.42
2:L:125:LEU:C	2:L:127:SER:N	2.77	0.41
2:M:30:ARG:HD2	2:M:30:ARG:N	2.35	0.41
2:L:30:ARG:N	2:L:30:ARG:HD2	2.36	0.41
2:M:33:LEU:HG	2:M:34:ASN:N	2.35	0.41
1:I:95:HIS:CE1	1:I:100(P):PHE:CE2	3.08	0.41
1:H:145:TYR:CE2	1:H:150:VAL:HG13	2.56	0.40
1:H:95:HIS:CE1	1:H:100(P):PHE:CE2	3.09	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	233/247 (94%)	227 (97%)	5 (2%)	1 (0%)	30	38
1	I	233/247 (94%)	228 (98%)	4 (2%)	1 (0%)	30	38
2	L	212/214 (99%)	208 (98%)	4 (2%)	0	100	100
2	M	212/214 (99%)	207 (98%)	5 (2%)	0	100	100
All	All	890/922 (96%)	870 (98%)	18 (2%)	2 (0%)	43	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	149	PRO
1	I	149	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	197/206 (96%)	193 (98%)	4 (2%)	48	67
1	I	197/206 (96%)	193 (98%)	4 (2%)	48	67
2	L	187/187 (100%)	178 (95%)	9 (5%)	23	35
2	M	187/187 (100%)	179 (96%)	8 (4%)	26	39
All	All	768/786 (98%)	743 (97%)	25 (3%)	33	50

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	78	LEU
1	H	92	CYS
1	H	178	LEU
1	H	209	LYS
2	L	3	GLN
2	L	11	LEU
2	L	24	GLN
2	L	55	GLN
2	L	90	GLN
2	L	123	GLU
2	L	147	GLN
2	L	181	LEU
2	L	188	LYS
1	I	78	LEU
1	I	92	CYS
1	I	178	LEU
1	I	209	LYS
2	M	3	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	M	11	LEU
2	M	24	GLN
2	M	55	GLN
2	M	90	GLN
2	M	147	GLN
2	M	181	LEU
2	M	188	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	3	GLN
1	H	39	GLN
1	H	164	HIS
2	L	38	GLN
2	L	55	GLN
2	L	138	ASN
2	L	189	HIS
2	L	199	GLN
1	I	3	GLN
1	I	164	HIS
1	I	192	GLN
2	M	55	GLN
2	M	137	ASN
2	M	138	ASN
2	M	189	HIS
2	M	199	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	H	1	1	7,8,9	2.31	2 (28%)	9,10,12	2.31	5 (55%)
1	PCA	I	1	1	7,8,9	2.32	2 (28%)	9,10,12	2.24	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	H	1	1	-	0/0/11/13	0/1/1/1
1	PCA	I	1	1	-	0/0/11/13	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	1	PCA	CD-N	4.98	1.46	1.34
1	I	1	PCA	CD-N	4.95	1.46	1.34
1	I	1	PCA	CA-N	3.47	1.50	1.46
1	H	1	PCA	CA-N	3.35	1.50	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	1	PCA	CB-CA-C	-3.68	107.61	112.66
1	I	1	PCA	CB-CA-C	-3.35	108.06	112.66
1	H	1	PCA	OE-CD-CG	-2.99	121.38	126.72
1	I	1	PCA	OE-CD-CG	-2.99	121.38	126.72
1	H	1	PCA	CA-N-CD	-2.97	103.41	113.58
1	I	1	PCA	CA-N-CD	-2.92	103.59	113.58
1	H	1	PCA	CB-CA-N	2.74	110.78	103.24
1	I	1	PCA	CB-CA-N	2.65	110.54	103.24
1	H	1	PCA	CG-CD-N	2.60	114.74	108.39
1	I	1	PCA	CG-CD-N	2.59	114.74	108.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TRS	H	301	-	7,7,7	0.36	0	9,9,9	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	H	301	-	-	3/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	H	301	TRS	C3-C-C1-O1
3	H	301	TRS	N-C-C1-O1
3	H	301	TRS	C2-C-C1-O1

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	H	232/247 (93%)	0.37	19 (8%)	17 19	16, 38, 83, 134	4 (1%)
1	I	233/247 (94%)	0.50	20 (8%)	16 17	25, 43, 85, 134	3 (1%)
2	L	213/214 (99%)	0.08	12 (5%)	30 32	19, 34, 79, 114	1 (0%)
2	M	213/214 (99%)	0.15	8 (3%)	44 46	19, 36, 83, 116	1 (0%)
All	All	891/922 (96%)	0.28	59 (6%)	24 26	16, 40, 85, 134	9 (1%)

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	185	PRO	7.5
1	H	129	LYS	7.4
1	I	186	SER	7.1
1	H	185	PRO	6.7
1	I	214	LYS	6.6
1	H	127	SER	6.4
1	I	129	LYS	6.4
1	I	127	SER	6.3
1	H	214	LYS	5.9
1	H	128	SER	5.9
1	I	130	SER	5.8
1	H	133	GLY	5.7
1	H	186	SER	5.6
1	I	192	GLN	5.2
1	H	134	GLY	4.9
1	H	130	SER	4.9
2	L	129	THR	4.8
2	M	129	THR	4.8
1	H	193	THR	4.7
1	I	204[A]	ASN	4.5
2	M	213	GLU	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	I	133	GLY	4.3
2	L	213	GLU	4.0
1	I	128	SER	3.9
2	L	126	LYS	3.8
1	I	134	GLY	3.6
2	L	212	GLY	3.6
2	M	126	LYS	3.2
1	H	131	THR	3.0
1	I	131	THR	3.0
2	M	188	LYS	2.9
2	L	123	GLU	2.8
2	M	212	GLY	2.7
1	H	213	PRO	2.7
2	M	56	ARG	2.7
1	I	213	PRO	2.7
1	H	201	LYS	2.6
1	I	184	VAL	2.6
1	I	132	SER	2.5
2	L	56	ARG	2.5
2	L	188	LYS	2.4
1	I	13	GLN	2.4
2	L	23	CYS	2.4
1	H	184	VAL	2.4
1	H	126	PRO	2.3
1	I	193	THR	2.3
1	H	208	ASP	2.3
1	I	92	CYS	2.3
2	L	181	LEU	2.3
2	M	127	SER	2.3
1	H	210	ARG	2.2
2	M	194	CYS	2.2
2	L	122	ASP	2.1
1	I	210	ARG	2.1
1	I	201	LYS	2.0
1	H	212	GLU	2.0
2	L	156	SER	2.0
1	H	140	CYS	2.0
2	L	194	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	PCA	H	1	8/9	0.94	0.10	36,44,54,65	0
1	PCA	I	1	8/9	0.94	0.10	38,44,56,67	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	TRS	H	301	8/8	0.84	0.22	70,79,85,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.