



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 12:11 PM UTC

PDB ID : 5FNO / pdb_00005fno
Title : Manganese Lipxygenase
Authors : Wennman, A.; Karkehabadi, S.; Oliw, E.H.; Chen, Y.
Deposited on : 2015-11-16
Resolution : 2.04 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	605	84% 10% 6%
1	B	605	85% 8% 6%
2	C	2	100%
2	D	2	50% 50%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9741 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MANGANESE LIPOXYGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	568	Total	C	N	O	S	0	0	0
			4472	2845	788	824	15			
1	B	568	Total	C	N	O	S	0	0	0
			4472	2845	788	824	15			

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLU	-	expression tag	UNP G4NAP4
A	2	PHE	-	expression tag	UNP G4NAP4
A	575	ALA	-	expression tag	UNP G4NAP4
A	576	ARG	-	expression tag	UNP G4NAP4
A	577	ARG	-	expression tag	UNP G4NAP4
A	578	PHE	-	expression tag	UNP G4NAP4
A	579	ASP	-	expression tag	UNP G4NAP4
A	580	ASP	-	expression tag	UNP G4NAP4
A	581	ARG	-	expression tag	UNP G4NAP4
A	582	GLY	-	expression tag	UNP G4NAP4
A	583	LEU	-	expression tag	UNP G4NAP4
A	584	SER	-	expression tag	UNP G4NAP4
A	585	GLN	-	expression tag	UNP G4NAP4
A	586	GLY	-	expression tag	UNP G4NAP4
A	587	MET	-	expression tag	UNP G4NAP4
A	588	PRO	-	expression tag	UNP G4NAP4
A	589	PHE	-	expression tag	UNP G4NAP4
A	590	ILE	-	expression tag	UNP G4NAP4
A	591	TRP	-	expression tag	UNP G4NAP4
A	592	GLN	-	expression tag	UNP G4NAP4
A	593	ALA	-	expression tag	UNP G4NAP4
A	594	LEU	-	expression tag	UNP G4NAP4
A	595	ASP	-	expression tag	UNP G4NAP4
A	596	PRO	-	expression tag	UNP G4NAP4
A	597	GLY	-	expression tag	UNP G4NAP4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	598	ASN	-	expression tag	UNP G4NAP4
A	599	ILE	-	expression tag	UNP G4NAP4
A	600	PRO	-	expression tag	UNP G4NAP4
A	601	PHE	-	expression tag	UNP G4NAP4
A	602	TYR	-	expression tag	UNP G4NAP4
A	603	LEU	-	expression tag	UNP G4NAP4
A	604	SER	-	expression tag	UNP G4NAP4
A	605	VAL	-	expression tag	UNP G4NAP4
B	1	GLU	-	expression tag	UNP G4NAP4
B	2	PHE	-	expression tag	UNP G4NAP4
B	575	ALA	-	expression tag	UNP G4NAP4
B	576	ARG	-	expression tag	UNP G4NAP4
B	577	ARG	-	expression tag	UNP G4NAP4
B	578	PHE	-	expression tag	UNP G4NAP4
B	579	ASP	-	expression tag	UNP G4NAP4
B	580	ASP	-	expression tag	UNP G4NAP4
B	581	ARG	-	expression tag	UNP G4NAP4
B	582	GLY	-	expression tag	UNP G4NAP4
B	583	LEU	-	expression tag	UNP G4NAP4
B	584	SER	-	expression tag	UNP G4NAP4
B	585	GLN	-	expression tag	UNP G4NAP4
B	586	GLY	-	expression tag	UNP G4NAP4
B	587	MET	-	expression tag	UNP G4NAP4
B	588	PRO	-	expression tag	UNP G4NAP4
B	589	PHE	-	expression tag	UNP G4NAP4
B	590	ILE	-	expression tag	UNP G4NAP4
B	591	TRP	-	expression tag	UNP G4NAP4
B	592	GLN	-	expression tag	UNP G4NAP4
B	593	ALA	-	expression tag	UNP G4NAP4
B	594	LEU	-	expression tag	UNP G4NAP4
B	595	ASP	-	expression tag	UNP G4NAP4
B	596	PRO	-	expression tag	UNP G4NAP4
B	597	GLY	-	expression tag	UNP G4NAP4
B	598	ASN	-	expression tag	UNP G4NAP4
B	599	ILE	-	expression tag	UNP G4NAP4
B	600	PRO	-	expression tag	UNP G4NAP4
B	601	PHE	-	expression tag	UNP G4NAP4
B	602	TYR	-	expression tag	UNP G4NAP4
B	603	LEU	-	expression tag	UNP G4NAP4
B	604	SER	-	expression tag	UNP G4NAP4
B	605	VAL	-	expression tag	UNP G4NAP4

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a

cetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		
3	B	1	Total	Mn	0	0
			1	1		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	346	Total	O	0	0
			346	346		
5	B	337	Total	O	0	0
			337	337		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: MANGANESE LIPOXYGENASE



• Molecule 1: MANGANESE LIPOXYGENASE



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.72Å 111.37Å 171.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.70 – 2.04 29.70 – 2.04	Depositor EDS
% Data completeness (in resolution range)	96.8 (29.70-2.04) 96.7 (29.70-2.04)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.06 (at 2.04Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.168 , 0.213 0.170 , 0.215	Depositor DCC
R_{free} test set	4188 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	35.9	Xtriage
Anisotropy	0.532	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9741	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/4585	0.52	0/6243
1	B	0.37	2/4585 (0.0%)	0.54	5/6243 (0.1%)
All	All	0.34	2/9170 (0.0%)	0.53	5/12486 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	556	ARG	CB-CG	-7.30	1.30	1.52
1	B	317	MET	SD-CE	-5.25	1.66	1.79

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	556	ARG	CG-CD-NE	-7.56	95.38	112.00
1	B	548	LEU	CA-C-N	-5.67	111.25	120.72
1	B	548	LEU	C-N-CA	-5.67	111.25	120.72
1	B	581	ARG	NE-CZ-NH1	-5.40	116.10	121.50
1	B	556	ARG	CB-CA-C	-5.33	102.51	110.88

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4472	0	4385	46	0
1	B	4472	0	4384	46	0
2	C	28	0	25	1	0
2	D	28	0	25	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	28	0	26	0	0
4	B	28	0	25	0	0
5	A	346	0	0	5	0
5	B	337	0	0	2	0
All	All	9741	0	8870	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 90 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:VAL:HA	1:A:132:GLN:HG2	1.60	0.84
1:A:581:ARG:NH1	1:B:581:ARG:HH12	1.75	0.84
1:B:556:ARG:N	1:B:556:ARG:HD2	1.93	0.82
1:B:548:LEU:O	1:B:556:ARG:NH2	2.15	0.79
1:A:128:LYS:HD3	1:A:148:SER:HB3	1.67	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	566/605 (94%)	547 (97%)	19 (3%)	0	100	100
1	B	566/605 (94%)	547 (97%)	19 (3%)	0	100	100
All	All	1132/1210 (94%)	1094 (97%)	38 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	468/496 (94%)	465 (99%)	3 (1%)	78	83
1	B	468/496 (94%)	467 (100%)	1 (0%)	87	91
All	All	936/992 (94%)	932 (100%)	4 (0%)	84	88

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	161	ARG
1	A	256	THR
1	A	346	GLN
1	B	284	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	263	ASN
1	B	585	GLN
1	B	559	ASN
1	A	585	GLN
1	B	248	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	1,2	14,14,15	0.37	0	17,19,21	0.75	0
2	NAG	C	2	2	14,14,15	0.33	0	17,19,21	0.69	1 (5%)
2	NAG	D	1	1,2	14,14,15	0.63	0	17,19,21	0.72	1 (5%)
2	NAG	D	2	2	14,14,15	0.28	0	17,19,21	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	NAG	C1-O5-C5	2.35	115.33	112.19
2	D	1	NAG	C1-O5-C5	2.28	115.24	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1	NAG	O5-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	D	1	NAG	O5-C5-C6-O6
2	D	1	NAG	C4-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	NAG	1	0
2	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	702	1	14,14,15	0.48	0	17,19,21	0.86	1 (5%)
4	NAG	A	703	1	14,14,15	0.47	0	17,19,21	0.74	0
4	NAG	B	703	1	14,14,15	0.41	0	17,19,21	0.70	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	B	702	1	14,14,15	0.42	0	17,19,21	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	702	1	-	3/6/23/26	0/1/1/1
4	NAG	A	703	1	-	0/6/23/26	0/1/1/1
4	NAG	B	703	1	-	2/6/23/26	0/1/1/1
4	NAG	B	702	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	703	NAG	C1-O5-C5	2.42	115.43	112.19
4	A	702	NAG	C1-O5-C5	2.10	115.00	112.19

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	703	NAG	O5-C5-C6-O6
4	B	703	NAG	C4-C5-C6-O6
4	A	702	NAG	C4-C5-C6-O6
4	A	702	NAG	C1-C2-N2-C7
4	A	702	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	568/605 (93%)	-0.27	7 (1%) 76 78	22, 40, 71, 99	0
1	B	568/605 (93%)	-0.36	2 (0%) 88 90	25, 40, 62, 96	0
All	All	1136/1210 (93%)	-0.32	9 (0%) 82 84	22, 40, 67, 99	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	38	VAL	3.0
1	A	335	GLY	2.7
1	A	329	ARG	2.6
1	A	38	VAL	2.4
1	A	121	LEU	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	D	2	14/15	0.79	0.12	52,59,65,68	0
2	NAG	C	1	14/15	0.85	0.17	45,54,83,97	0
2	NAG	C	2	14/15	0.88	0.10	57,66,76,76	0
2	NAG	D	1	14/15	0.89	0.09	38,44,58,76	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	B	702	14/15	0.71	0.13	72,78,81,83	0
4	NAG	A	702	14/15	0.75	0.13	63,67,72,72	0
4	NAG	B	703	14/15	0.82	0.12	54,73,82,84	0
4	NAG	A	703	14/15	0.85	0.12	61,65,69,72	0
3	MN	A	701	1/1	0.99	0.09	33,33,33,33	0
3	MN	B	701	1/1	1.00	0.06	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.