



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:33 PM UTC

PDB ID : 2FOR / pdb\_00002for  
Title : Crystal Structure of the Shigella flexneri Farnesyl Pyrophosphate Synthase Complex with an Isopentenyl Pyrophosphate  
Authors : Minasov, G.; Brunzelle, J.S.; Shuvalova, L.; Collart, F.R.; Joachimiak, A.; Anderson, W.F.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2006-01-13  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	 24% 72% 15% 12%
1	B	323	 20% 75% 11% 13%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4743 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Geranyltranstransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	283	2156	1346	385	414	11	0	3	0
1	B	281	2170	1355	386	416	13	0	6	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	expression tag	GB 24050587
A	-22	HIS	-	expression tag	GB 24050587
A	-21	HIS	-	expression tag	GB 24050587
A	-20	HIS	-	expression tag	GB 24050587
A	-19	HIS	-	expression tag	GB 24050587
A	-18	HIS	-	expression tag	GB 24050587
A	-17	HIS	-	expression tag	GB 24050587
A	-16	SER	-	expression tag	GB 24050587
A	-15	SER	-	expression tag	GB 24050587
A	-14	GLY	-	expression tag	GB 24050587
A	-13	VAL	-	expression tag	GB 24050587
A	-12	ASP	-	expression tag	GB 24050587
A	-11	LEU	-	expression tag	GB 24050587
A	-10	GLY	-	expression tag	GB 24050587
A	-9	THR	-	expression tag	GB 24050587
A	-8	GLU	-	expression tag	GB 24050587
A	-7	ASN	-	expression tag	GB 24050587
A	-6	LEU	-	expression tag	GB 24050587
A	-5	TYR	-	expression tag	GB 24050587
A	-4	PHE	-	expression tag	GB 24050587
A	-3	GLN	-	expression tag	GB 24050587
A	-2	SER	-	expression tag	GB 24050587
A	-1	ASN	-	expression tag	GB 24050587
A	0	ALA	-	expression tag	GB 24050587
B	-23	MET	-	expression tag	GB 24050587

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	HIS	-	expression tag	GB 24050587
B	-21	HIS	-	expression tag	GB 24050587
B	-20	HIS	-	expression tag	GB 24050587
B	-19	HIS	-	expression tag	GB 24050587
B	-18	HIS	-	expression tag	GB 24050587
B	-17	HIS	-	expression tag	GB 24050587
B	-16	SER	-	expression tag	GB 24050587
B	-15	SER	-	expression tag	GB 24050587
B	-14	GLY	-	expression tag	GB 24050587
B	-13	VAL	-	expression tag	GB 24050587
B	-12	ASP	-	expression tag	GB 24050587
B	-11	LEU	-	expression tag	GB 24050587
B	-10	GLY	-	expression tag	GB 24050587
B	-9	THR	-	expression tag	GB 24050587
B	-8	GLU	-	expression tag	GB 24050587
B	-7	ASN	-	expression tag	GB 24050587
B	-6	LEU	-	expression tag	GB 24050587
B	-5	TYR	-	expression tag	GB 24050587
B	-4	PHE	-	expression tag	GB 24050587
B	-3	GLN	-	expression tag	GB 24050587
B	-2	SER	-	expression tag	GB 24050587
B	-1	ASN	-	expression tag	GB 24050587
B	0	ALA	-	expression tag	GB 24050587

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	1
2	A	1	Total O P 5 4 1	0	1
2	A	1	Total O P 5 4 1	0	1
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	1

- Molecule 3 is ISOPENTYL PYROPHOSPHATE (CCD ID: IPR) (formula: C<sub>5</sub>H<sub>14</sub>O<sub>7</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O P 14 5 7 2	0	1
3	B	1	Total C O P 14 5 7 2	0	1

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	161	Total O 163 163	0	7

Continued on next page...


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	B	187	Total 191	O 191	0	12


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Geranyltranstransferase



- Molecule 1: Geranyltranstransferase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.66Å 75.66Å 216.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	65.51 – 2.00 65.51 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.7 (65.51-2.00) 97.7 (65.51-2.00)	Depositor EDS
$R_{merge}$	0.02	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.60 (at 1.99Å)	Xtrriage
Refinement program	REFMAC 5.2	Depositor
R, $R_{free}$	0.181 , 0.233 0.185 , 0.184	Depositor DCC
$R_{free}$ test set	2301 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.5	Xtrriage
Anisotropy	0.164	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 62.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.079 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	4743	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IPR, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.63	0/2187	1.02	5/2958 (0.2%)
1	B	0.79	5/2201 (0.2%)	1.04	2/2978 (0.1%)
All	All	0.71	5/4388 (0.1%)	1.03	7/5936 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	12	LYS	C-O	12.33	1.39	1.24
1	B	12	LYS	C-N	8.83	1.45	1.33
1	B	19	SER	C-O	7.41	1.33	1.24
1	B	16[A]	GLN	C-O	6.26	1.31	1.24
1	B	16[B]	GLN	C-O	6.26	1.31	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	255	GLY	N-CA-C	-6.68	104.81	112.29
1	A	255	GLY	N-CA-C	-6.64	103.02	111.72
1	A	154	MET	N-CA-C	6.20	118.55	111.11
1	B	41	LEU	N-CA-C	5.76	120.58	113.50
1	A	41	LEU	N-CA-C	5.51	120.11	112.45
1	A	130	MET	CA-C-N	5.13	126.25	119.84
1	A	130	MET	C-N-CA	5.13	126.25	119.84

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2156	0	2162	33	0
1	B	2170	0	2176	28	0
2	A	25	0	0	1	0
2	B	10	0	0	0	0
3	A	14	0	10	1	0
3	B	14	0	10	4	0
4	A	163	0	0	4	0
4	B	191	0	0	2	0
All	All	4743	0	4358	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:162:LEU:O	1:A:165:GLU:HG2	1.85	0.77
1:A:27:PHE:O	1:A:30:THR:HG22	1.88	0.73
1:A:132:GLU:HB2	4:A:413:HOH:O	1.91	0.70
1:A:83:HIS:HD2	1:B:115:ASP:OD2	1.75	0.70
1:B:165:GLU:HG2	1:B:247:LYS:HE2	1.77	0.67
1:B:175:GLU:HG2	1:B:250:TYR:OH	1.95	0.66
1:A:264:ARG:HA	1:A:295:ILE:CD1	2.28	0.64
1:A:128:ALA:O	1:A:138:ARG:NH2	2.22	0.64
1:B:45:LYS:HE2	3:B:302[B]:IPR:O13	2.02	0.60
1:A:267:ILE:HG13	1:A:295:ILE:HB	1.85	0.59
1:B:222:GLN:HE21	1:B:222:GLN:HA	1.69	0.57
1:A:7:LEU:O	1:A:11:VAL:HG23	2.03	0.57
1:A:32:VAL:HG23	4:A:336:HOH:O	2.03	0.57
1:A:112:LEU:HD21	1:B:89:MET:HE1	1.87	0.57
1:A:112:LEU:CD2	1:B:89:MET:HE1	2.35	0.57
1:B:95:ARG:HH22	3:B:302[B]:IPR:HC41	1.70	0.57
1:A:258[A]:GLN:HG3	4:A:390[A]:HOH:O	2.05	0.56
1:A:264:ARG:HA	1:A:295:ILE:HD12	1.88	0.55
1:B:12:LYS:O	1:B:16[A]:GLN:HG3	2.06	0.55

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:264:ARG:HG3	1:B:295:ILE:HD11	1.88	0.55
1:A:53:TYR:O	1:A:57:HIS:ND1	2.40	0.55
1:B:48:ARG:NH2	3:B:302[B]:IPR:O14	2.39	0.54
1:A:118:GLN:NE2	1:B:119:THR:OG1	2.39	0.54
1:B:165:GLU:HG2	1:B:247:LYS:CE	2.39	0.52
1:A:3:PHE:N	1:A:4:PRO:HD2	2.25	0.52
1:A:38:TYR:CD1	1:A:42:LEU:HD12	2.45	0.52
1:A:68:ALA:HB3	1:A:69:PRO:CD	2.40	0.51
1:B:95:ARG:NH2	3:B:302[B]:IPR:HC41	2.26	0.50
1:B:7:LEU:O	1:B:11:VAL:HG23	2.12	0.49
1:A:158:GLN:OE1	1:A:181:LYS:HE2	2.13	0.48
1:A:68:ALA:HB3	1:A:69:PRO:HD3	1.96	0.48
1:B:18:LEU:HD11	1:B:74:GLU:HG2	1.94	0.48
1:A:264:ARG:HA	1:A:295:ILE:HD11	1.95	0.48
1:A:22:ILE:HG21	1:A:37:GLN:HG2	1.96	0.47
1:A:83:HIS:HE1	1:A:111:ILE:O	1.97	0.47
1:A:45:LYS:NZ	3:A:305[B]:IPR:HC52	2.29	0.47
1:B:178:HIS:CE1	1:B:220:GLN:HG2	2.49	0.47
1:B:79:TYR:CD1	1:B:79:TYR:C	2.93	0.47
1:A:79:TYR:CD1	1:A:79:TYR:C	2.93	0.46
1:A:119:THR:OG1	1:B:118:GLN:NE2	2.45	0.46
1:A:47:LEU:HD23	1:A:47:LEU:HA	1.74	0.45
1:A:179:ARG:NH1	2:A:303:PO4:O2	2.36	0.45
1:B:288:GLU:HG3	4:B:447:HOH:O	2.16	0.45
1:A:19:SER:HB2	4:A:332:HOH:O	2.16	0.44
1:A:217:LEU:HD11	1:A:250:TYR:OH	2.17	0.44
1:B:136:ARG:C	1:B:136:ARG:HD2	2.43	0.43
1:A:48:ARG:HB2	1:A:49:PRO:HD3	2.00	0.43
1:B:32:VAL:HG22	1:B:109:ASN:HD22	1.82	0.43
1:B:206:PRO:HD2	4:B:477:HOH:O	2.18	0.43
1:B:193:GLY:O	1:B:196:SER:HB3	2.19	0.42
1:B:222:GLN:NE2	1:B:298:ASN:HD22	2.17	0.42
1:A:171:LEU:HD22	1:A:254:LEU:HD21	2.01	0.41
1:B:92:ASP:O	1:B:102:HIS:HE1	2.02	0.41
1:B:125:LEU:HD23	1:B:125:LEU:HA	1.94	0.40
1:A:151:ILE:O	1:A:156:GLY:HA3	2.20	0.40
1:B:79:TYR:HB2	1:B:114:GLY:O	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/323 (87%)	276 (98%)	6 (2%)	0	100	100
1	B	283/323 (88%)	280 (99%)	3 (1%)	0	100	100
All	All	565/646 (88%)	556 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/254 (88%)	220 (98%)	4 (2%)	51	58
1	B	227/254 (89%)	226 (100%)	1 (0%)	84	89
All	All	451/508 (89%)	446 (99%)	5 (1%)	65	73

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	ARG
1	A	79	TYR
1	A	210	LYS
1	A	290	LEU
1	B	79	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	28	GLN
1	A	29	ASN
1	A	37	GLN
1	A	83	HIS
1	A	109	ASN
1	A	118	GLN
1	A	178	HIS
1	A	272	GLN
1	A	276	GLN
1	A	296	GLN
1	A	298	ASN
1	B	6	GLN
1	B	13	GLN
1	B	64	ASN
1	B	102	HIS
1	B	109	ASN
1	B	118	GLN
1	B	222	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	A	304	-	4,4,4	0.87	0	6,6,6	0.53	0
3	IPR	A	305[B]	-	12,13,13	1.36	2 (16%)	16,19,19	1.05	1 (6%)
2	PO4	A	300[A]	-	4,4,4	1.12	0	6,6,6	0.59	0
2	PO4	B	300	-	4,4,4	0.88	0	6,6,6	0.70	0
2	PO4	A	303	-	4,4,4	0.90	0	6,6,6	0.55	0
2	PO4	A	301[A]	-	4,4,4	0.86	0	6,6,6	0.41	0
3	IPR	B	302[B]	-	12,13,13	1.39	2 (16%)	16,19,19	1.07	0
2	PO4	B	301[A]	-	4,4,4	0.89	0	6,6,6	0.49	0
2	PO4	A	302[A]	-	4,4,4	0.93	0	6,6,6	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IPR	A	305[B]	-	-	3/13/13/13	-
3	IPR	B	302[B]	-	-	6/13/13/13	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	305[B]	IPR	C1-C2	-3.69	1.32	1.51
3	B	302[B]	IPR	C1-C2	-3.67	1.32	1.51
3	B	302[B]	IPR	P7-O10	2.39	1.62	1.59
3	A	305[B]	IPR	P7-O10	2.30	1.62	1.59

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	305[B]	IPR	C3-C2-C1	2.02	119.55	110.53

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	305[B]	IPR	C2-C4-C5-O6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	B	302[B]	IPR	P7-O10-P11-O12
3	B	302[B]	IPR	C5-O6-P7-O8
3	B	302[B]	IPR	C2-C4-C5-O6
3	B	302[B]	IPR	C1-C2-C4-C5
3	A	305[B]	IPR	C1-C2-C4-C5
3	B	302[B]	IPR	C5-O6-P7-O10
3	A	305[B]	IPR	C3-C2-C4-C5
3	B	302[B]	IPR	P7-O10-P11-O14

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	305[B]	IPR	1	0
2	A	303	PO4	1	0
3	B	302[B]	IPR	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	283/323 (87%)	1.56	79 (27%) <b>1</b> <b>1</b>	24, 50, 57, 68	3 (1%)
1	B	281/323 (86%)	1.51	63 (22%) <b>2</b> <b>2</b>	24, 51, 60, 73	6 (2%)
All	All	564/646 (87%)	1.54	142 (25%) <b>1</b> <b>1</b>	24, 50, 59, 73	9 (1%)

All (142) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	182	THR	6.1
1	B	182	THR	5.8
1	B	183	GLY	5.6
1	B	184	ALA	5.5
1	A	207	VAL	5.1
1	A	121	ALA	5.0
1	B	139	ILE	5.0
1	B	14	ALA	4.9
1	A	186	ILE	4.5
1	B	19	SER	4.4
1	B	22	ILE	4.4
1	A	41	LEU	4.4
1	A	183	GLY	4.3
1	A	181	LYS	4.3
1	B	94[A]	LEU	4.3
1	A	119	THR	3.9
1	A	125	LEU	3.8
1	B	4	PRO	3.7
1	A	124	ILE	3.7
1	A	185	LEU	3.6
1	A	229	VAL	3.6
1	B	162	LEU	3.6
1	A	128	ALA	3.5
1	B	15	ASN	3.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	142	ILE	3.5
1	A	212	ALA	3.5
1	A	164	ALA	3.4
1	A	123	SER	3.4
1	A	21	PHE	3.3
1	A	129	ASN	3.3
1	B	181	LYS	3.3
1	A	31	PRO	3.2
1	B	131	PRO	3.1
1	B	16[A]	GLN	3.1
1	B	20	ARG	3.1
1	A	221	VAL	3.1
1	A	73	VAL	3.0
1	A	162	LEU	3.0
1	B	134	SER	3.0
1	B	143	SER	3.0
1	A	203	ARG	3.0
1	B	81	LEU	3.0
1	A	268	ASP	3.0
1	B	1	MET	3.0
1	A	82	ILE	2.9
1	B	140	SER	2.9
1	A	246	GLY	2.9
1	A	210	LYS	2.9
1	B	185	LEU	2.8
1	A	220	GLN	2.8
1	A	209	ASP	2.8
1	B	17	ALA	2.8
1	B	128	ALA	2.8
1	B	146	ALA	2.8
1	A	20	ARG	2.8
1	B	141	MET	2.8
1	B	199	ASP	2.8
1	B	186	ILE	2.8
1	A	116	ALA	2.7
1	A	22	ILE	2.7
1	B	86	LEU	2.7
1	A	272	GLN	2.7
1	A	208	LEU	2.7
1	B	82	ILE	2.7
1	B	11	VAL	2.6
1	B	52	VAL	2.6

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	103	VAL	2.6
1	A	230	GLY	2.6
1	A	273	ALA	2.6
1	A	192	LEU	2.6
1	A	225	ILE	2.6
1	B	205	LEU	2.6
1	B	226	LEU	2.6
1	B	180	HIS	2.5
1	B	7	LEU	2.5
1	B	164	ALA	2.5
1	A	36	MET	2.5
1	A	56	GLY	2.5
1	A	188	ALA	2.4
1	A	205	LEU	2.4
1	A	114	GLY	2.4
1	A	228	VAL	2.4
1	A	117	LEU	2.4
1	B	80	SER	2.4
1	A	247	LYS	2.4
1	B	298	ASN	2.3
1	B	166	GLY	2.3
1	B	83	HIS	2.3
1	A	206	PRO	2.3
1	A	38	TYR	2.3
1	A	299	LYS	2.3
1	B	295	ILE	2.3
1	A	249	THR	2.3
1	B	13	GLN	2.3
1	B	263	ALA	2.3
1	A	81	LEU	2.3
1	A	195	LEU	2.3
1	A	279	GLU	2.3
1	A	295	ILE	2.3
1	A	24	PRO	2.3
1	A	197	ALA	2.3
1	B	18	LEU	2.3
1	B	79	TYR	2.3
1	A	168	HIS	2.2
1	A	3	PHE	2.2
1	A	27	PHE	2.2
1	A	120	LEU	2.2
1	B	165	GLU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	266	LEU	2.2
1	B	57	HIS	2.2
1	B	147	SER	2.2
1	A	96	ARG	2.2
1	B	267	ILE	2.2
1	A	54	ALA	2.2
1	A	280	GLN	2.2
1	A	7	LEU	2.2
1	B	145	LEU	2.2
1	A	15	ASN	2.2
1	A	180	HIS	2.1
1	A	184	ALA	2.1
1	A	200	LYS	2.1
1	A	250	TYR	2.1
1	A	224	ASP	2.1
1	B	229	VAL	2.1
1	A	194	ALA	2.1
1	A	204	ALA	2.1
1	B	200	LYS	2.1
1	A	29	ASN	2.1
1	B	64	ASN	2.1
1	B	227	ASP	2.1
1	A	78	ALA	2.1
1	B	110	ALA	2.1
1	B	278	ALA	2.1
1	A	39	GLY	2.1
1	B	56	GLY	2.1
1	B	104	LYS	2.1
1	A	103	VAL	2.1
1	A	47	LEU	2.0
1	B	41	LEU	2.0
1	A	214	SER	2.0
1	B	204	ALA	2.0
1	A	35	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	IPR	B	302[B]	14/14	0.66	0.32	46,51,53,54	14
3	IPR	A	305[B]	14/14	0.75	0.37	71,73,75,75	14
2	PO4	A	303	5/5	0.75	0.17	55,56,58,58	5
2	PO4	A	304	5/5	0.76	0.16	70,71,73,74	5
2	PO4	A	301[A]	5/5	0.78	0.21	51,51,52,52	5
2	PO4	B	301[A]	5/5	0.79	0.23	143,143,144,144	5
2	PO4	A	302[A]	5/5	0.82	0.17	56,58,59,59	5
2	PO4	B	300	5/5	0.95	0.18	45,49,50,51	0
2	PO4	A	300[A]	5/5	0.97	0.13	36,38,40,42	5

### 6.5 Other polymers [i](#)

There are no such residues in this entry.