



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 01:05 PM UTC

PDB ID : 4FOR / pdb_00004for
Title : Crystal Structure of C-lobe of Bovine lactoferrin Complexed with Flurbiprofen at 1.58 Å Resolution
Authors : Shukla, P.K.; Gautam, L.; Sinha, M.; Kaur, P.; Sharma, S.; Singh, T.P.
Deposited on : 2012-06-21
Resolution : 1.58 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	6% 78% 21% .
2	B	6	67% 83% 17%
3	C	2	50% 50%
3	D	2	100%

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 3098 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	0	0
			2560	1593	448	499	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	565	LYS	ASN	SEE REMARK 999	UNP P24627
A	608	GLU	LYS	SEE REMARK 999	UNP P24627

- Molecule 2 is a protein called C-terminal peptide from Lactotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	6	Total	C	N	O	S	0	0	0
			45	29	6	9	1			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Fe	0	0
			1	1		

- Molecule 5 is CARBONATE ION (CCD ID: CO3) (formula: CO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	1	3		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Zn	0	0
			2	2		

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 9 is FLURBIPROFEN (CCD ID: FLP) (formula: C₁₅H₁₃FO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	F	O	0	0
			18	15	1	2		

- Molecule 10 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	385	Total 385	O 385	0	0
11	B	2	Total 2	O 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Lactotransferrin




• Molecule 2: C-terminal peptide from Lactotransferrin



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.45Å 50.01Å 65.47Å 90.00° 107.14° 90.00°	Depositor
Resolution (Å)	62.56 – 1.58 62.56 – 1.58	Depositor EDS
% Data completeness (in resolution range)	99.8 (62.56-1.58) 99.5 (62.56-1.58)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.22 (at 1.58Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.165 , 0.205 (Not available) , 0.206	Depositor DCC
R_{free} test set	2691 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	18.2	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.022 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3098	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FLP, ZN, GOL, SO4, NAG, CO3, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.78	30/2608 (1.2%)	1.55	12/3533 (0.3%)
2	B	1.02	0/45	1.04	0/58
All	All	1.77	30/2653 (1.1%)	1.54	12/3591 (0.3%)

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	458	HIS	ND1-CE1	10.57	1.43	1.32
1	A	344	ARG	C-O	8.28	1.34	1.23
1	A	599	SER	C-O	-7.46	1.14	1.23
1	A	601	SER	C-O	-7.25	1.15	1.24
1	A	521	GLU	N-CA	7.12	1.54	1.46
1	A	590	ALA	CA-C	-6.98	1.44	1.52
1	A	578	ARG	C-O	6.38	1.31	1.24
1	A	534	ALA	N-CA	6.32	1.54	1.46
1	A	668	ALA	N-CA	6.27	1.53	1.46
1	A	520	LYS	N-CA	6.22	1.54	1.46
1	A	650	LYS	C-O	6.18	1.31	1.23
1	A	531	ARG	C-O	-6.08	1.17	1.24
1	A	457	CYS	N-CA	6.06	1.53	1.46
1	A	571	LEU	C-O	5.89	1.30	1.23
1	A	514	LYS	C-O	-5.70	1.17	1.23
1	A	385	LEU	N-CA	5.70	1.53	1.46
1	A	600	ARG	CD-NE	-5.49	1.38	1.46
1	A	521	GLU	CA-C	5.47	1.60	1.52
1	A	347	TRP	N-CA	5.39	1.52	1.45
1	A	579	LYS	N-CA	-5.37	1.39	1.45
1	A	632	PHE	N-CA	5.28	1.52	1.46
1	A	384	VAL	N-CA	5.27	1.52	1.46
1	A	619	GLY	C-O	-5.26	1.18	1.23
1	A	519	SER	N-CA	5.25	1.53	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	598	VAL	C-O	-5.22	1.18	1.24
1	A	613	HIS	CG-CD2	5.17	1.41	1.35
1	A	588	HIS	CD2-NE2	5.16	1.43	1.37
1	A	467	TRP	CA-CB	5.14	1.62	1.53
1	A	452	LYS	C-O	-5.08	1.17	1.23
1	A	603	ARG	CZ-NH1	5.05	1.39	1.32

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	563	ASN	N-CA-C	7.15	122.03	113.16
1	A	675	CYS	N-CA-C	6.58	120.56	112.54
1	A	605	ALA	N-CA-C	6.11	118.02	111.36
1	A	474	ILE	N-CA-C	5.88	116.62	110.62
1	A	519	SER	CA-C-O	-5.67	111.95	119.10
1	A	521	GLU	N-CA-C	-5.50	101.71	109.96
1	A	530	PHE	CA-C-O	-5.30	114.80	120.42
1	A	455	LYS	CA-CB-CG	-5.21	103.68	114.10
1	A	558	ALA	N-CA-C	5.17	116.96	110.24
1	A	375	SER	CA-CB-OG	-5.15	100.80	111.10
1	A	454	LYS	O-C-N	5.12	128.54	122.20
1	A	436	VAL	CA-CB-CG2	5.08	119.04	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2560	0	2480	29	0
2	B	45	0	39	1	0
3	C	28	0	25	1	0
3	D	28	0	25	0	0
4	A	1	0	0	0	0
5	A	4	0	0	0	0
6	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	2	0	0	0	0
8	A	14	0	13	0	0
9	A	18	0	12	2	0
10	A	6	0	8	0	0
11	A	385	0	0	1	0
11	B	2	0	0	0	0
All	All	3098	0	2602	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:565:LYS:HE3	1:A:567:GLU:HB2	1.69	0.74
1:A:565:LYS:HE3	1:A:567:GLU:H	1.55	0.72
1:A:565:LYS:CE	1:A:567:GLU:HB2	2.21	0.71
9:A:711:FLP:O1	9:A:711:FLP:H8	1.94	0.67
1:A:671:ASN:HA	1:A:674:LYS:HD3	1.78	0.65
1:A:565:LYS:HE3	1:A:567:GLU:N	2.12	0.64
1:A:395:ASP:HA	1:A:595:HIS:CD2	2.38	0.57
1:A:660:TYR:O	9:A:711:FLP:H132	2.06	0.56
1:A:342:TYR:HB2	1:A:606:HIS:HE1	1.71	0.54
1:A:669:ILE:CG2	1:A:673:LYS:HE3	2.40	0.51
1:A:565:LYS:HE3	1:A:567:GLU:CB	2.39	0.50
1:A:364:GLN:HG3	1:A:629:PHE:HB2	1.94	0.48
1:A:352:PRO:HG2	1:A:520:LYS:HD3	1.95	0.48
1:A:364:GLN:OE1	1:A:628:LYS:HE2	2.15	0.47
1:A:653:GLY:O	1:A:654:ARG:C	2.60	0.44
1:A:633:LYS:HD3	1:A:633:LYS:HA	1.83	0.44
1:A:366:GLY:C	1:A:367:GLN:HG2	2.42	0.44
1:A:626:PRO:HA	1:A:630:CYS:SG	2.59	0.43
1:A:527:THR:CG2	1:A:560:TRP:HZ2	2.31	0.43
1:A:668:ALA:HB1	3:C:1:NAG:H83	2.01	0.43
1:A:565:LYS:HD2	1:A:566:ARG:N	2.33	0.42
1:A:483:PHE:CD2	1:A:483:PHE:C	2.97	0.42
1:A:509:ASP:HA	11:A:1161:HOH:O	2.20	0.41
1:A:566:ARG:HG2	1:A:581:VAL:HG21	2.01	0.41
1:A:455:LYS:HB3	1:A:504:LEU:HD11	2.02	0.41
1:A:520:LYS:HA	1:A:520:LYS:HD2	1.71	0.41
1:A:565:LYS:CE	1:A:567:GLU:H	2.29	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:522:LYS:HD3	1:A:523:TYR:CE1	2.56	0.41
1:A:404:LYS:NZ	2:B:681:LEU:O	2.54	0.40
1:A:573:CYS:SG	1:A:579:LYS:HG3	2.61	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/335 (99%)	324 (97%)	8 (2%)	1 (0%)	36	20
2	B	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
All	All	337/341 (99%)	327 (97%)	9 (3%)	1 (0%)	36	20

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	432	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/278 (100%)	275 (99%)	3 (1%)	65	43
2	B	4/4 (100%)	3 (75%)	1 (25%)	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	282/282 (100%)	278 (99%)	4 (1%)	59	34

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	515	CYS
1	A	567	GLU
1	A	621	ASN
2	B	681	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	355	GLN
1	A	360	GLN
1	A	609	GLN
1	A	613	HIS
1	A	642	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	1,3	14,14,15	1.14	2 (14%)	17,19,21	1.97	4 (23%)
3	NAG	C	2	3	14,14,15	0.63	0	17,19,21	2.04	5 (29%)
3	NAG	D	1	1,3	14,14,15	0.87	0	17,19,21	2.21	7 (41%)
3	NAG	D	2	3	14,14,15	0.61	0	17,19,21	2.51	10 (58%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1	NAG	C4-C3	2.13	1.57	1.52
3	C	1	NAG	C3-C2	2.07	1.56	1.52

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	2	NAG	O4-C4-C3	5.52	123.40	110.38
3	D	1	NAG	C1-O5-C5	4.67	118.45	112.19
3	C	2	NAG	C2-N2-C7	4.29	128.65	122.90
3	C	1	NAG	C1-C2-N2	-4.25	103.73	110.43
3	D	2	NAG	C2-N2-C7	4.01	128.28	122.90
3	D	1	NAG	C1-C2-N2	-3.94	104.23	110.43
3	C	1	NAG	C1-O5-C5	3.84	117.34	112.19
3	C	2	NAG	O3-C3-C2	-3.81	101.49	109.40
3	D	1	NAG	C2-N2-C7	-3.72	117.92	122.90
3	C	1	NAG	O5-C1-C2	-3.68	105.59	111.29
3	C	2	NAG	O7-C7-C8	-3.15	116.44	122.05
3	D	2	NAG	O7-C7-N2	3.07	127.40	121.98
3	C	2	NAG	C4-C3-C2	3.04	115.47	111.02
3	D	2	NAG	C1-O5-C5	2.98	116.18	112.19
3	D	2	NAG	C4-C3-C2	-2.98	106.65	111.02
3	D	2	NAG	C3-C4-C5	-2.83	105.11	110.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	NAG	O4-C4-C5	-2.62	102.86	109.32
3	D	2	NAG	O7-C7-C8	-2.56	117.50	122.05
3	D	1	NAG	C8-C7-N2	2.32	119.97	116.12
3	D	1	NAG	C4-C3-C2	-2.22	107.76	111.02
3	D	2	NAG	C6-C5-C4	-2.22	107.56	113.02
3	D	2	NAG	O3-C3-C2	-2.17	104.90	109.40
3	C	1	NAG	C3-C4-C5	2.11	114.07	110.23
3	D	2	NAG	O5-C1-C2	2.10	114.54	111.29
3	D	1	NAG	O7-C7-C8	-2.08	118.35	122.05
3	C	2	NAG	C1-O5-C5	2.08	114.97	112.19

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	FLP	A	711	-	19,19,19	2.26	2 (10%)	26,26,26	2.17	9 (34%)
10	GOL	A	712	-	5,5,5	0.60	0	5,5,5	0.84	0
8	NAG	A	706	1	14,14,15	1.33	1 (7%)	17,19,21	3.18	8 (47%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SO4	A	703	-	4,4,4	0.53	0	6,6,6	0.86	0
5	CO3	A	702	4	3,3,3	0.60	0	2,3,3	2.73	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	GOL	A	712	-	-	1/4/4/4	-
8	NAG	A	706	1	-	0/6/23/26	0/1/1/1
9	FLP	A	711	-	-	4/12/12/12	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	711	FLP	C6-C11	8.21	1.53	1.39
8	A	706	NAG	O7-C7	3.82	1.31	1.23
9	A	711	FLP	C12-C14	2.59	1.59	1.52

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	706	NAG	O5-C1-C2	-7.44	99.78	111.29
8	A	706	NAG	O7-C7-N2	5.37	131.48	121.98
8	A	706	NAG	C8-C7-N2	-5.35	107.25	116.12
9	A	711	FLP	F-C11-C6	5.07	127.03	119.00
8	A	706	NAG	C2-N2-C7	4.88	129.44	122.90
9	A	711	FLP	O1-C14-O	-4.55	113.75	124.08
8	A	706	NAG	C1-O5-C5	3.95	117.48	112.19
5	A	702	CO3	O3-C-O1	3.81	129.41	119.68
9	A	711	FLP	C10-C11-C6	-3.66	117.73	123.59
9	A	711	FLP	C2-C6-C11	3.58	128.53	122.86
9	A	711	FLP	C13-C12-C14	2.84	118.81	110.68
9	A	711	FLP	C7-C6-C2	-2.70	113.34	118.70
9	A	711	FLP	C9-C12-C14	2.64	116.30	108.75
9	A	711	FLP	O1-C14-C12	2.49	124.81	114.25
8	A	706	NAG	C1-C2-N2	2.49	114.36	110.43
8	A	706	NAG	O5-C5-C6	2.28	112.09	107.66
8	A	706	NAG	O3-C3-C4	-2.19	105.21	110.38
9	A	711	FLP	C9-C10-C11	2.07	122.04	119.52

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	A	711	FLP	C14-C12-C9-C10
9	A	711	FLP	C14-C12-C9-C8
9	A	711	FLP	C9-C12-C14-O
9	A	711	FLP	C9-C12-C14-O1
10	A	712	GOL	C1-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	711	FLP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	335/335 (100%)	0.13	21 (6%)	26 35	10, 20, 49, 90	1 (0%)
2	B	6/6 (100%)	2.99	4 (66%)	0 0	36, 43, 102, 109	0
All	All	341/341 (100%)	0.18	25 (7%)	21 30	10, 20, 50, 109	1 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	681	LEU	6.4
1	A	342	TYR	6.4
1	A	418	SER	3.8
1	A	609	GLN	3.4
1	A	431	GLU	3.3
2	B	683	ALA	3.3
1	A	422	SER	3.3
1	A	416	LYS	3.2
1	A	420	HIS	3.2
1	A	667	THR	3.0
2	B	685	ALA	2.9
1	A	663	THR	2.9
1	A	627	ASP	2.9
1	A	421	SER	2.9
1	A	602	ASP	2.7
1	A	636	THR	2.6
1	A	423	LEU	2.5
1	A	343	THR	2.5
1	A	419	LYS	2.4
1	A	373	THR	2.3
1	A	621	ASN	2.3
1	A	623	LYS	2.2
2	B	682	GLU	2.2
1	A	613	HIS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	628	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	D	2	14/15	0.82	0.13	38,44,65,69	0
3	NAG	C	2	14/15	0.83	0.12	43,52,62,68	0
3	NAG	C	1	14/15	0.90	0.09	31,35,42,46	0
3	NAG	D	1	14/15	0.95	0.07	24,27,35,36	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	FLP	A	711	18/18	0.58	0.26	32,54,60,61	18
8	NAG	A	706	14/15	0.77	0.13	34,47,56,67	0
10	GOL	A	712	6/6	0.84	0.13	30,39,43,49	0
6	SO4	A	703	5/5	0.96	0.07	37,38,40,51	0
7	ZN	A	705	1/1	0.97	0.04	21,21,21,21	0
5	CO3	A	702	4/4	0.99	0.04	11,12,12,13	0
7	ZN	A	704	1/1	0.99	0.02	17,17,17,17	0
4	FE	A	701	1/1	1.00	0.01	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.