



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:54 PM UTC

PDB ID : 5FO8 / pdb\_00005fo8  
Title : Crystal Structure of Human Complement C3b in Complex with MCP (CCP1-4)  
Authors : Forneris, F.; Wu, J.; Xue, X.; Gros, P.  
Deposited on : 2015-11-18  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	645	 4% 84% 15% ..
2	B	915	 7% 82% 15% ..
3	C	252	 4% 46% 5% 5% 49%
4	D	2	 50% 50%
5	E	3	 33% 67%

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 13810 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called COMPLEMENT C3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	639	4979	3169	844	951	15	0	0	0

- Molecule 2 is a protein called COMPLEMENT C3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	894	7145	4530	1201	1376	38	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1013	GLU	GLN	SEE REMARK 999	UNP P01024

- Molecule 3 is a protein called MEMBRANE COFACTOR PROTEIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	129	999	644	157	189	9	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	2	28	16	2	10	0	0	0

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
			Total	C	N				O
5	E	3	39	22	2	15	0	0	0

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	A	1	4	2	2	0	0
6	A	1	4	2	2	0	0
6	A	1	4	2	2	0	0
6	A	1	4	2	2	0	0
6	B	1	4	2	2	0	0
6	B	1	4	2	2	0	0
6	B	1	4	2	2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	172	Total O 172 172	0	0
7	B	293	Total O 293 293	0	0
7	C	27	Total O 27 27	0	0


### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: COMPLEMENT C3





#### • Molecule 2: COMPLEMENT C3





- Molecule 3: MEMBRANE COFACTOR PROTEIN



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.83Å 130.63Å 233.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.02 – 2.40 57.02 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.8 (57.02-2.40) 95.2 (57.02-2.40)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 2.40Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.188 , 0.219 0.193 , 0.223	Depositor DCC
$R_{free}$ test set	4928 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.1	Xtrriage
Anisotropy	0.221	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 55.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	13810	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/5079	0.76	4/6902 (0.1%)
2	B	0.36	0/7291	0.74	1/9872 (0.0%)
3	C	0.29	0/1029	0.72	0/1399
All	All	0.34	0/13399	0.75	5/18173 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	23	SER	CA-C-N	-5.92	113.84	119.76
1	A	23	SER	C-N-CA	-5.92	113.84	119.76
1	A	90	ILE	CA-C-N	5.52	126.30	120.11
1	A	90	ILE	C-N-CA	5.52	126.30	120.11
2	B	1440	ASP	N-CA-C	-5.20	106.89	113.18

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4979	0	5037	57	0
2	B	7145	0	7070	83	0
3	C	999	0	962	6	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	28	0	25	2	0
5	E	39	0	34	0	0
6	A	16	0	24	1	0
6	B	112	0	168	12	0
7	A	172	0	0	4	0
7	B	293	0	0	19	1
7	C	27	0	0	0	0
All	All	13810	0	13320	141	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 141 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1586:SER:OG	7:B:2246:HOH:O	1.83	0.96
2:B:873:CYS:HG	2:B:1513:CYS:HG	1.07	0.86
1:A:351:SER:O	7:A:2090:HOH:O	2.00	0.79
2:B:1010:CYS:SG	7:B:2078:HOH:O	2.42	0.77
2:B:819:ASP:O	7:B:2035:HOH:O	2.03	0.76

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:2137:HOH:O	7:B:2291:HOH:O[1_545]	2.02	0.18

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	635/645 (98%)	617 (97%)	17 (3%)	1 (0%)	43	58
2	B	889/915 (97%)	860 (97%)	29 (3%)	0	100	100
3	C	127/252 (50%)	123 (97%)	4 (3%)	0	100	100
All	All	1651/1812 (91%)	1600 (97%)	50 (3%)	1 (0%)	48	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	659	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	563/567 (99%)	539 (96%)	24 (4%)	26	44
2	B	792/810 (98%)	751 (95%)	41 (5%)	21	36
3	C	114/222 (51%)	109 (96%)	5 (4%)	25	43
All	All	1469/1599 (92%)	1399 (95%)	70 (5%)	23	40

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	1523	SER
2	B	1526	LYS
3	C	179	GLU
2	B	793	ILE
1	A	660	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	1127	GLN
2	B	1277	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	C	238	GLN
2	B	1355	GLN
2	B	857	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	D	1	1,4	14,14,15	0.53	0	17,19,21	1.04	1 (5%)
4	NAG	D	2	4	14,14,15	0.52	0	17,19,21	0.76	0
5	NAG	E	1	2,5	14,14,15	0.50	0	17,19,21	1.01	1 (5%)
5	NAG	E	2	5	14,14,15	0.56	0	17,19,21	1.13	2 (11%)
5	BMA	E	3	5	11,11,12	0.81	0	15,15,17	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	1,4	-	3/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	E	1	2,5	-	0/6/23/26	0/1/1/1
5	NAG	E	2	5	-	0/6/23/26	0/1/1/1
5	BMA	E	3	5	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	1	NAG	C1-O5-C5	2.77	115.91	112.19
5	E	2	NAG	C3-C4-C5	2.64	115.03	110.23
5	E	2	NAG	C4-C3-C2	2.49	114.67	111.02
4	D	1	NAG	O5-C1-C2	-2.22	107.86	111.29

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	1	NAG	C8-C7-N2-C2
4	D	1	NAG	O7-C7-N2-C2
4	D	2	NAG	C8-C7-N2-C2
4	D	2	NAG	O7-C7-N2-C2
4	D	1	NAG	C4-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

32 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	EDO	B	2672	-	3,3,3	0.43	0	2,2,2	0.36	0
6	EDO	B	2665	-	3,3,3	0.44	0	2,2,2	0.35	0
6	EDO	B	2687	-	3,3,3	0.42	0	2,2,2	0.41	0
6	EDO	B	2675	-	3,3,3	0.43	0	2,2,2	0.36	0
6	EDO	B	2683	-	3,3,3	0.45	0	2,2,2	0.37	0
6	EDO	B	2668	-	3,3,3	0.44	0	2,2,2	0.27	0
6	EDO	B	2690	-	3,3,3	0.45	0	2,2,2	0.44	0
6	EDO	B	2688	-	3,3,3	0.45	0	2,2,2	0.35	0
6	EDO	B	2666	-	3,3,3	0.42	0	2,2,2	0.35	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	EDO	B	2664	-	3,3,3	0.42	0	2,2,2	0.46	0
6	EDO	A	1669	-	3,3,3	0.44	0	2,2,2	0.34	0
6	EDO	B	2671	-	3,3,3	0.45	0	2,2,2	0.35	0
6	EDO	B	2681	-	3,3,3	0.43	0	2,2,2	0.35	0
6	EDO	B	2678	-	3,3,3	0.41	0	2,2,2	0.41	0
6	EDO	B	2673	-	3,3,3	0.44	0	2,2,2	0.43	0
6	EDO	B	2667	-	3,3,3	0.41	0	2,2,2	0.37	0
6	EDO	B	2682	-	3,3,3	0.44	0	2,2,2	0.34	0
6	EDO	A	1667	-	3,3,3	0.42	0	2,2,2	0.37	0
6	EDO	B	2686	-	3,3,3	0.43	0	2,2,2	0.38	0
6	EDO	B	2685	-	3,3,3	0.40	0	2,2,2	0.44	0
6	EDO	B	2677	-	3,3,3	0.44	0	2,2,2	0.37	0
6	EDO	B	2670	-	3,3,3	0.42	0	2,2,2	0.42	0
6	EDO	A	1666	-	3,3,3	0.43	0	2,2,2	0.37	0
6	EDO	B	2689	-	3,3,3	0.41	0	2,2,2	0.37	0
6	EDO	B	2691	-	3,3,3	0.40	0	2,2,2	0.48	0
6	EDO	A	1668	-	3,3,3	0.44	0	2,2,2	0.37	0
6	EDO	B	2676	-	3,3,3	0.44	0	2,2,2	0.31	0
6	EDO	B	2679	-	3,3,3	0.42	0	2,2,2	0.41	0
6	EDO	B	2669	-	3,3,3	0.44	0	2,2,2	0.36	0
6	EDO	B	2674	-	3,3,3	0.44	0	2,2,2	0.39	0
6	EDO	B	2680	-	3,3,3	0.40	0	2,2,2	0.40	0
6	EDO	B	2684	-	3,3,3	0.44	0	2,2,2	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	B	2672	-	-	0/1/1/1	-
6	EDO	B	2665	-	-	0/1/1/1	-
6	EDO	B	2687	-	-	1/1/1/1	-
6	EDO	B	2675	-	-	0/1/1/1	-
6	EDO	B	2683	-	-	0/1/1/1	-
6	EDO	B	2668	-	-	0/1/1/1	-
6	EDO	B	2690	-	-	0/1/1/1	-
6	EDO	B	2688	-	-	0/1/1/1	-
6	EDO	B	2666	-	-	0/1/1/1	-
6	EDO	B	2664	-	-	0/1/1/1	-
6	EDO	A	1669	-	-	1/1/1/1	-
6	EDO	B	2671	-	-	0/1/1/1	-
6	EDO	B	2681	-	-	0/1/1/1	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	B	2678	-	-	0/1/1/1	-
6	EDO	B	2673	-	-	0/1/1/1	-
6	EDO	B	2667	-	-	1/1/1/1	-
6	EDO	B	2682	-	-	0/1/1/1	-
6	EDO	A	1667	-	-	0/1/1/1	-
6	EDO	B	2686	-	-	0/1/1/1	-
6	EDO	B	2685	-	-	0/1/1/1	-
6	EDO	B	2677	-	-	0/1/1/1	-
6	EDO	B	2670	-	-	0/1/1/1	-
6	EDO	A	1666	-	-	0/1/1/1	-
6	EDO	B	2689	-	-	0/1/1/1	-
6	EDO	B	2691	-	-	0/1/1/1	-
6	EDO	A	1668	-	-	0/1/1/1	-
6	EDO	B	2676	-	-	0/1/1/1	-
6	EDO	B	2679	-	-	1/1/1/1	-
6	EDO	B	2669	-	-	0/1/1/1	-
6	EDO	B	2674	-	-	0/1/1/1	-
6	EDO	B	2680	-	-	0/1/1/1	-
6	EDO	B	2684	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1669	EDO	O1-C1-C2-O2
6	B	2687	EDO	O1-C1-C2-O2
6	B	2679	EDO	O1-C1-C2-O2
6	B	2667	EDO	O1-C1-C2-O2

There are no ring outliers.

12 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	2687	EDO	1	0
6	B	2675	EDO	1	0
6	B	2666	EDO	2	0
6	B	2671	EDO	1	0
6	B	2681	EDO	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	2678	EDO	1	0
6	B	2673	EDO	1	0
6	B	2682	EDO	1	0
6	B	2677	EDO	1	0
6	A	1666	EDO	1	0
6	B	2691	EDO	1	0
6	B	2680	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	639/645 (99%)	0.37	23 (3%) 46 42	38, 64, 99, 164	0
2	B	894/915 (97%)	0.35	61 (6%) 23 20	25, 64, 118, 157	1 (0%)
3	C	129/252 (51%)	0.58	10 (7%) 19 15	43, 71, 101, 122	0
All	All	1662/1812 (91%)	0.38	94 (5%) 29 25	25, 65, 110, 164	1 (0%)

The worst 5 of 94 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	540	ALA	5.6
2	B	1506	CYS	4.1
1	A	461	LEU	4.1
1	A	665	PRO	3.9
2	B	1466	TYR	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	BMA	E	3	11/12	0.29	0.17	122,143,153,153	0
5	NAG	E	2	14/15	0.60	0.15	102,132,138,142	0
4	NAG	D	2	14/15	0.67	0.18	94,112,117,117	0


*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	E	1	14/15	0.77	0.14	94,100,108,121	0
4	NAG	D	1	14/15	0.91	0.11	67,79,90,101	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	EDO	A	1669	4/4	0.46	0.25	99,99,100,101	0
6	EDO	B	2686	4/4	0.73	0.17	70,75,86,92	0
6	EDO	B	2688	4/4	0.74	0.24	76,78,82,85	0
6	EDO	B	2665	4/4	0.76	0.26	88,90,92,93	0
6	EDO	A	1666	4/4	0.78	0.26	79,82,83,85	0
6	EDO	B	2682	4/4	0.79	0.23	77,81,86,91	0
6	EDO	B	2666	4/4	0.80	0.20	59,60,62,63	2
6	EDO	B	2668	4/4	0.81	0.22	82,82,84,84	0
6	EDO	A	1668	4/4	0.82	0.20	92,92,95,96	0
6	EDO	B	2674	4/4	0.84	0.22	82,87,93,96	0
6	EDO	B	2673	4/4	0.85	0.24	70,70,75,77	1
6	EDO	B	2671	4/4	0.85	0.18	76,76,76,85	1
6	EDO	B	2677	4/4	0.85	0.17	85,87,88,88	0
6	EDO	B	2676	4/4	0.86	0.24	65,66,77,82	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	EDO	B	2669	4/4	0.86	0.17	87,87,88,88	0
6	EDO	B	2681	4/4	0.86	0.20	91,92,92,98	0
6	EDO	B	2678	4/4	0.88	0.28	62,63,64,65	0
6	EDO	A	1667	4/4	0.89	0.21	90,90,92,93	0
6	EDO	B	2685	4/4	0.90	0.19	48,58,61,64	0
6	EDO	B	2667	4/4	0.90	0.20	64,65,68,76	1
6	EDO	B	2679	4/4	0.90	0.18	68,73,77,82	0
6	EDO	B	2675	4/4	0.91	0.14	48,50,51,55	3
6	EDO	B	2672	4/4	0.92	0.16	74,74,76,78	0
6	EDO	B	2691	4/4	0.92	0.16	47,54,66,77	0
6	EDO	B	2689	4/4	0.93	0.14	62,65,71,73	0
6	EDO	B	2684	4/4	0.94	0.15	36,43,55,66	0
6	EDO	B	2664	4/4	0.94	0.12	42,56,63,72	0
6	EDO	B	2680	4/4	0.94	0.12	78,86,89,91	0
6	EDO	B	2683	4/4	0.95	0.11	55,56,56,57	0
6	EDO	B	2687	4/4	0.96	0.12	51,53,55,59	0
6	EDO	B	2690	4/4	0.96	0.11	33,38,48,54	0
6	EDO	B	2670	4/4	0.96	0.12	45,47,48,49	3

## 6.5 Other polymers [i](#)

There are no such residues in this entry.