



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 11:39 AM UTC

PDB ID : 5FOC / pdb_00005foc
Title : Crystal structure of the P.falciparum cytosolic leucyl-tRNA synthetase editing domain (space group P21)
Authors : Palencia, A.; Sonoiki, E.; Guo, D.; Ahyong, V.; Dong, C.; Li, X.; Hernandez, V.S.; Gut, J.; Legac, J.; Cooper, R.; Alley, M.R.K.; Freund, Y.R.; DeRisi, J.; Cusack, S.; Rosenthal, P.J.
Deposited on : 2015-11-19
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	 75% 21%
1	B	374	 76% 20%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 5369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LEUCYL-TRNA SYNTHETASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	297	2486	1608	387	476	15	0	7	0
1	B	300	2501	1616	390	481	14	0	6	0

There are 100 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	269	GLY	-	expression tag	UNP C6KT64
A	270	ALA	-	expression tag	UNP C6KT64
A	271	MET	-	expression tag	UNP C6KT64
A	273	SER	CYS	engineered mutation	UNP C6KT64
A	?	-	ILE	deletion	UNP C6KT64
A	?	-	LYS	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	VAL	deletion	UNP C6KT64
A	?	-	GLN	deletion	UNP C6KT64
A	?	-	ILE	deletion	UNP C6KT64
A	?	-	PRO	deletion	UNP C6KT64
A	?	-	LYS	deletion	UNP C6KT64
A	?	-	SER	deletion	UNP C6KT64
A	?	-	GLU	deletion	UNP C6KT64
A	?	-	ASP	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	THR	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASP	deletion	UNP C6KT64
A	?	-	ASP	deletion	UNP C6KT64
A	?	-	ASP	deletion	UNP C6KT64
A	?	-	THR	deletion	UNP C6KT64
A	?	-	LEU	deletion	UNP C6KT64
A	?	-	GLU	deletion	UNP C6KT64
A	?	-	LYS	deletion	UNP C6KT64

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	LYS	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASP	deletion	UNP C6KT64
A	?	-	VAL	deletion	UNP C6KT64
A	?	-	ILE	deletion	UNP C6KT64
A	?	-	THR	deletion	UNP C6KT64
A	?	-	LYS	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	THR	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	GLU	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	MET	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	ASN	deletion	UNP C6KT64
A	?	-	VAL	deletion	UNP C6KT64
A	520	GLY	TYR	engineered mutation	UNP C6KT64
B	269	GLY	-	expression tag	UNP C6KT64
B	270	ALA	-	expression tag	UNP C6KT64
B	271	MET	-	expression tag	UNP C6KT64
B	273	SER	CYS	engineered mutation	UNP C6KT64
B	?	-	ILE	deletion	UNP C6KT64
B	?	-	LYS	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	VAL	deletion	UNP C6KT64
B	?	-	GLN	deletion	UNP C6KT64
B	?	-	ILE	deletion	UNP C6KT64
B	?	-	PRO	deletion	UNP C6KT64
B	?	-	LYS	deletion	UNP C6KT64
B	?	-	SER	deletion	UNP C6KT64
B	?	-	GLU	deletion	UNP C6KT64
B	?	-	ASP	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	THR	deletion	UNP C6KT64

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASP	deletion	UNP C6KT64
B	?	-	ASP	deletion	UNP C6KT64
B	?	-	ASP	deletion	UNP C6KT64
B	?	-	THR	deletion	UNP C6KT64
B	?	-	LEU	deletion	UNP C6KT64
B	?	-	GLU	deletion	UNP C6KT64
B	?	-	LYS	deletion	UNP C6KT64
B	?	-	LYS	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASP	deletion	UNP C6KT64
B	?	-	VAL	deletion	UNP C6KT64
B	?	-	ILE	deletion	UNP C6KT64
B	?	-	THR	deletion	UNP C6KT64
B	?	-	LYS	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	THR	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	MET	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	ASN	deletion	UNP C6KT64
B	?	-	VAL	deletion	UNP C6KT64
B	520	GLY	TYR	engineered mutation	UNP C6KT64

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	219	Total O 219 219	0	0
2	B	163	Total O 163 163	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.60Å 53.49Å 122.79Å 90.00° 97.31° 90.00°	Depositor
Resolution (Å)	50.01 – 1.50 50.01 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.3 (50.01-1.50) 99.3 (50.01-1.50)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.14 (at 1.45Å)	Xtrriage
Refinement program	REFMAC 5.8.0123	Depositor
R, R_{free}	0.183 , 0.234 0.191 , 0.236	Depositor DCC
R_{free} test set	5506 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtrriage
Anisotropy	0.257	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5369	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.83 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4350e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.99	1/2551 (0.0%)	1.00	1/3428 (0.0%)
1	B	0.93	1/2566 (0.0%)	0.94	0/3450
All	All	0.96	2/5117 (0.0%)	0.97	1/6878 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	402	TYR	N-CA	6.15	1.54	1.46
1	B	402	TYR	C-O	5.55	1.31	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	566	GLY	N-CA-C	-5.23	107.03	115.08

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2486	0	2496	15	0
1	B	2501	0	2507	11	0
2	A	219	0	0	2	0
2	B	163	0	0	0	0
All	All	5369	0	5003	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 26 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:663[A]:MET:HE1	1:A:671:LEU:HD12	1.66	0.75
1:B:635:ASP:HB3	1:B:638[A]:LEU:HD13	1.80	0.63
1:A:630:VAL:CG1	1:A:638[A]:LEU:HD11	2.33	0.59
1:A:575[A]:SER:OG	1:A:577:ASP:OD1	2.22	0.58
1:A:663[A]:MET:CE	1:A:671:LEU:HD12	2.34	0.58

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/374 (79%)	291 (98%)	5 (2%)	0	100	100
1	B	298/374 (80%)	293 (98%)	5 (2%)	0	100	100
All	All	594/748 (79%)	584 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/349 (82%)	284 (99%)	3 (1%)	68	45
1	B	289/349 (83%)	288 (100%)	1 (0%)	86	75
All	All	576/698 (82%)	572 (99%)	4 (1%)	81	57

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	638[A]	LEU
1	A	638[B]	LEU
1	A	664	LYS
1	B	599	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	301	GLN
1	B	412	ASN
1	B	597	ASN
1	A	460	ASN
1	A	424	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/374 (79%)	-0.05	4 (1%) 75 78	14, 26, 46, 63	7 (2%)
1	B	300/374 (80%)	0.23	10 (3%) 49 53	16, 29, 54, 71	6 (2%)
All	All	597/748 (79%)	0.09	14 (2%) 61 65	14, 28, 50, 71	13 (2%)

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	434	VAL	5.7
1	B	303	VAL	3.6
1	A	361	THR	3.4
1	A	304	LEU	2.9
1	B	304	LEU	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.