



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:04 AM UTC

PDB ID : 6FOX / pdb_00006fox
Title : The crystal structure of P.fluorescens Kynurenine 3-monooxygenase (KMO) in complex with kynurenine
Authors : Levy, C.W.; Leys, D.
Deposited on : 2018-02-08
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	5% 91% 6% .
1	B	460	8% 86% 10% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15070 atoms, of which 7086 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Kynurenine 3-monooxygenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	449	Total	C	H	N	O	S	0	22	0
			7124	2234	3553	665	653	19			
1	B	448	Total	C	H	N	O	S	0	6	0
			6954	2188	3460	646	641	19			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	252	SER	CYS	engineered mutation	UNP Q84HF5
A	461	SER	-	expression tag	UNP Q84HF5
B	252	SER	CYS	engineered mutation	UNP Q84HF5
B	461	SER	-	expression tag	UNP Q84HF5

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	
			84	27	31	9	15	2	
2	B	1	Total	C	H	N	O	P	
			84	27	31	9	15	2	

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca		
			1	1	0	0

- Molecule 4 is (2S)-2-amino-4-(2-aminophenyl)-4-oxobutanoic acid (CCD ID: KYN) (formula: C₁₀H₁₂N₂O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	H	N	O		
			26	10	11	2	3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	507	Total	O		
			507	507	0	0
5	B	290	Total	O		
			290	290	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Kynurenine 3-monooxygenase



• Molecule 1: Kynurenine 3-monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.91Å 52.54Å 136.17Å 90.00° 104.14° 90.00°	Depositor
Resolution (Å)	44.02 – 1.90 44.02 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.7 (44.02-1.90) 95.9 (44.02-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.170 , 0.215 0.169 , 0.214	Depositor DCC
R_{free} test set	1961 reflections (2.57%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15070	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.95 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9430e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, CA, KYN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/3741	0.40	0/5081
1	B	0.20	0/3593	0.37	0/4878
All	All	0.23	0/7334	0.38	0/9959

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3571	3553	3456	17	2
1	B	3494	3460	3433	36	1
2	A	53	31	31	1	0
2	B	53	31	31	1	0
3	A	1	0	0	0	0
4	B	15	11	11	1	0
5	A	507	0	0	1	0
5	B	290	0	0	5	0
All	All	7984	7086	6962	54	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:113:ARG:NH1	5:B:602:HOH:O	2.20	0.73
1:B:368:GLU:OE2	5:B:601:HOH:O	2.11	0.68
1:A:106[A]:ILE:HD11	1:A:408:THR:CG2	2.23	0.68
1:B:9:GLN:NE2	1:B:32:GLN:OE1	2.33	0.62
1:A:449:GLU:OE1	1:A:452:ARG:NH2	2.33	0.61
1:B:180:VAL:HG11	1:B:299:THR:HG21	1.86	0.57
1:A:148:SER:HB3	1:A:154:ARG:HD3	1.87	0.57
1:B:215:ILE:HG23	1:B:222:MET:HE2	1.87	0.55
1:B:16:GLY:O	1:B:20[B]:THR:HG22	2.07	0.55
1:A:90:GLY:O	1:A:91:THR:OG1	2.22	0.53
1:B:184:GLU:HG3	1:B:295:LEU:HD11	1.92	0.52
1:B:20[B]:THR:CG2	1:B:118:LEU:HB3	2.40	0.52
1:B:242:GLN:NE2	5:B:618:HOH:O	2.43	0.51
1:A:106[A]:ILE:HD11	1:A:408:THR:HG22	1.93	0.50
1:B:51:ARG:CG	1:B:52:SER:N	2.74	0.50
1:B:184:GLU:CG	1:B:295:LEU:HD11	2.41	0.50
1:B:189:GLN:HG3	1:B:191:HIS:H	1.77	0.50
1:A:58:ALA:HB2	1:A:106[A]:ILE:HD12	1.94	0.50
1:A:63:HIS:CG	1:A:411:ARG:HD3	2.48	0.48
1:B:51:ARG:HG3	1:B:52:SER:N	2.27	0.48
1:B:162:LEU:O	1:B:306:ALA:HA	2.14	0.47
1:A:429:LYS:NZ	5:A:616:HOH:O	2.43	0.46
1:B:173:ARG:HD2	1:B:297:LEU:HD11	1.97	0.46
1:B:265:PHE:HA	1:B:268:ARG:HE	1.81	0.45
1:A:158:ARG:HH11	1:A:158:ARG:HG2	1.80	0.45
1:B:57:LEU:HA	1:B:324[A]:MET:HE1	1.98	0.45
1:A:17:LEU:HD13	1:A:324[A]:MET:HG3	1.99	0.45
1:B:384:LEU:HD23	1:B:432:VAL:HG13	1.99	0.44
1:B:146:THR:CG2	1:B:154:ARG:HD2	2.48	0.44
1:B:189:GLN:HB3	1:B:292:LEU:HB2	2.00	0.44
1:B:222:MET:HE1	1:B:373:MET:HB3	1.99	0.44
1:B:53:ILE:HD11	1:B:111:ARG:HE	1.83	0.44
1:A:162:LEU:O	1:A:306:ALA:HA	2.18	0.43
4:B:501:KYN:N1	2:B:502:FAD:O4	2.52	0.43
1:B:193:TYR:HA	1:B:237:LEU:O	2.18	0.43
1:A:400:PHE:HZ	1:A:428:LEU:HD11	1.84	0.42
1:B:63:HIS:CG	1:B:411:ARG:HD3	2.54	0.42
1:B:297:LEU:O	5:B:604:HOH:O	2.21	0.42
1:A:390:GLN:O	1:A:394:GLN:HG2	2.19	0.42
1:B:20[B]:THR:CG2	1:B:118:LEU:HD13	2.49	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:329:GLU:OE2	5:B:603:HOH:O	2.21	0.42
1:B:274:SER:N	1:B:275:PRO:HD2	2.35	0.42
1:B:16:GLY:O	1:B:20[B]:THR:CG2	2.68	0.41
1:B:263:ARG:O	1:B:267:GLN:HG2	2.19	0.41
1:B:143:GLN:NE2	1:B:160:HIS:HA	2.35	0.41
1:A:398:THR:HG23	1:A:399:ARG:HG3	2.03	0.41
1:B:20[B]:THR:HG21	1:B:118:LEU:HB3	2.02	0.41
1:B:51:ARG:O	1:B:52:SER:O	2.39	0.41
1:B:145:LEU:O	1:B:156:GLU:HA	2.20	0.41
1:A:323:GLY:HA3	2:A:501:FAD:H1'2	2.02	0.40
1:B:51:ARG:NH1	1:B:195:GLU:OE2	2.54	0.40
1:A:155:LEU:C	1:A:155:LEU:HD23	2.46	0.40
1:B:166:ASP:OD1	1:B:166:ASP:N	2.54	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:158:ARG:NH2	1:A:336:GLU:OE1[2_854]	2.11	0.09
1:B:47:GLY:O	1:B:72:ARG:HH22[2_853]	1.52	0.08
1:A:158:ARG:HH22	1:A:336:GLU:OE1[2_854]	1.59	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	469/460 (102%)	456 (97%)	12 (3%)	1 (0%)	43	36
1	B	450/460 (98%)	441 (98%)	8 (2%)	1 (0%)	43	36
All	All	919/920 (100%)	897 (98%)	20 (2%)	2 (0%)	43	36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	52	SER
1	A	91	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	382/371 (103%)	374 (98%)	8 (2%)	47	44
1	B	369/371 (100%)	365 (99%)	4 (1%)	65	67
All	All	751/742 (101%)	739 (98%)	12 (2%)	57	54

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	SER
1	A	55[A]	LEU
1	A	55[B]	LEU
1	A	91	THR
1	A	147	LEU
1	A	155	LEU
1	A	381	THR
1	A	444	ASP
1	B	51	ARG
1	B	53	ILE
1	B	267	GLN
1	B	381	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	132	ASN
1	A	160	HIS
1	B	218	HIS
1	B	286	HIS
1	B	301	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	KYN	B	501	-	14,15,15	1.58	2 (14%)	15,20,20	1.14	2 (13%)
2	FAD	B	502	-	58,58,58	2.94	26 (44%)	85,89,89	1.60	19 (22%)
2	FAD	A	501	-	58,58,58	0.60	1 (1%)	85,89,89	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	KYN	B	501	-	-	1/12/12/12	0/1/1/1
2	FAD	B	502	-	-	3/34/50/50	0/6/6/6
2	FAD	A	501	-	-	1/34/50/50	0/6/6/6

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	502	FAD	C2B-C3B	-9.87	1.26	1.53
2	B	502	FAD	C10-N1	7.37	1.48	1.33
2	B	502	FAD	C6A-N6A	5.54	1.48	1.34
2	B	502	FAD	C2-N1	5.24	1.48	1.36
4	B	501	KYN	O-C	5.05	1.37	1.22
2	B	502	FAD	O4B-C1B	-4.93	1.30	1.42
2	B	502	FAD	C5B-C4B	-4.33	1.38	1.51
2	B	502	FAD	PA-O3P	4.30	1.64	1.59
2	B	502	FAD	C5X-N5	-4.06	1.32	1.39
2	B	502	FAD	C2-N3	3.96	1.47	1.39
2	B	502	FAD	C9A-N10	-3.94	1.34	1.41
2	B	502	FAD	O4B-C4B	3.79	1.53	1.45
2	B	502	FAD	C2B-C1B	3.64	1.64	1.53
2	B	502	FAD	C8A-N9A	-3.53	1.31	1.37
2	B	502	FAD	C9A-C5X	-3.49	1.35	1.41
2	B	502	FAD	O3B-C3B	3.29	1.51	1.43
2	A	501	FAD	P-O3P	-3.22	1.56	1.59
2	B	502	FAD	P-O3P	3.06	1.62	1.59
2	B	502	FAD	C4-N3	2.89	1.44	1.38
2	B	502	FAD	C4X-N5	2.87	1.37	1.30
2	B	502	FAD	O4-C4	-2.77	1.18	1.23
2	B	502	FAD	O2-C2	-2.53	1.19	1.24
2	B	502	FAD	C3B-C4B	2.52	1.59	1.53
2	B	502	FAD	C4X-C4	2.49	1.53	1.44
2	B	502	FAD	PA-O5B	2.41	1.68	1.59
2	B	502	FAD	C8A-N7A	2.28	1.36	1.31
2	B	502	FAD	O2B-C2B	2.19	1.48	1.43
2	B	502	FAD	C9-C9A	2.07	1.43	1.39
4	B	501	KYN	OXT-C	-2.06	1.24	1.30

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	502	FAD	N3A-C2A-N1A	-4.59	121.64	128.58
2	B	502	FAD	C5A-C4A-N3A	-4.23	120.89	126.72
2	B	502	FAD	C4-N3-C2	-3.71	119.06	125.64
2	B	502	FAD	N9A-C8A-N7A	-3.41	109.09	113.94
2	B	502	FAD	C10-C4X-N5	-3.35	117.96	124.81
2	B	502	FAD	C2A-N3A-C4A	3.16	119.56	111.83
2	B	502	FAD	C4X-C4-N3	2.91	120.66	113.25
2	B	502	FAD	C4A-N9A-C8A	2.79	108.67	105.74
2	B	502	FAD	C4A-C5A-N7A	-2.78	107.40	110.58
2	B	502	FAD	N3A-C4A-N9A	2.75	131.84	127.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	KYN	OXT-C-O	-2.64	118.08	124.08
2	B	502	FAD	C5A-N7A-C8A	2.61	107.56	103.45
2	B	502	FAD	O4-C4-C4X	-2.57	119.75	126.53
2	B	502	FAD	C4'-C3'-C2'	-2.52	109.37	113.57
2	B	502	FAD	C4X-C10-N10	2.49	120.05	116.48
2	B	502	FAD	C4X-C10-N1	-2.48	118.51	124.59
2	B	502	FAD	C4-C4X-N5	2.42	121.55	118.21
2	B	502	FAD	C10-N1-C2	2.34	121.91	116.85
4	B	501	KYN	CB-C1-CD2	-2.25	117.20	119.82
2	B	502	FAD	C5X-C9A-N10	2.08	119.85	117.97
2	B	502	FAD	C4-C4X-C10	2.07	120.49	116.93

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	502	FAD	C3B-C4B-C5B-O5B
2	B	502	FAD	O4B-C4B-C5B-O5B
2	B	502	FAD	P-O3P-PA-O1A
2	A	501	FAD	O4B-C4B-C5B-O5B
4	B	501	KYN	OXT-C-CA-N


There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	501	KYN	1	0
2	B	502	FAD	1	0
2	A	501	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	449/460 (97%)	-0.16	24 (5%) 32 34	6, 22, 77, 130	11 (2%)
1	B	448/460 (97%)	0.31	35 (7%) 19 20	11, 31, 78, 129	3 (0%)
All	All	897/920 (97%)	0.07	59 (6%) 24 26	6, 27, 78, 130	14 (1%)

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	150	VAL	4.9
1	A	91	THR	4.7
1	A	377	VAL	4.5
1	B	155	LEU	4.4
1	A	90	GLY	4.3
1	B	50	GLY	4.3
1	B	284	PHE	4.3
1	B	259	GLY	4.2
1	A	245	ALA	4.1
1	A	246	ALA	4.0
1	B	249	ALA	4.0
1	B	458	SER	3.8
1	B	248	PRO	3.8
1	A	154	ARG	3.8
1	B	287	HIS	3.7
1	A	448	HIS	3.6
1	B	261	ALA	3.6
1	A	438	LEU	3.5
1	B	258	ASP	3.4
1	B	264	ARG	3.3
1	A	92	PRO	3.3
1	B	244	PRO	3.2
1	A	100	ARG	3.1
1	A	394	GLN	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	286	HIS	3.1
1	B	268	ARG	3.0
1	B	257	VAL	2.9
1	B	51	ARG	2.9
1	B	288	PRO	2.9
1	A	378	ALA	2.9
1	B	182	LEU	2.7
1	B	271	PRO	2.7
1	B	394	GLN	2.6
1	B	260	HIS	2.5
1	B	256	LEU	2.5
1	A	443	LEU	2.5
1	A	452	ARG	2.5
1	B	100	ARG	2.4
1	A	249	ALA	2.4
1	A	439	THR	2.3
1	A	247	GLN	2.3
1	B	265	PHE	2.3
1	A	51	ARG	2.2
1	B	52	SER	2.2
1	B	151	SER	2.2
1	A	89	PRO	2.2
1	B	262	ALA	2.2
1	B	427	LEU	2.2
1	B	8	ARG	2.2
1	A	93	PRO	2.1
1	B	386	ARG	2.1
1	A	155	LEU	2.1
1	A	453	CYS	2.1
1	A	436	SER	2.0
1	B	280	LEU	2.0
1	B	254	ALA	2.0
1	A	153	GLU	2.0
1	B	178	SER	2.0
1	B	49	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	KYN	B	501	15/15	0.92	0.12	22,43,61,73	0
2	FAD	B	502	53/53	0.97	0.09	14,27,98,101	0
2	FAD	A	501	53/53	0.98	0.04	9,13,22,33	0
3	CA	A	502	1/1	0.99	0.07	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD B 502:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.