



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:22 AM UTC

PDB ID : 6FOY / pdb_00006foy
Title : The crystal structure of P.fluorescens Kynurenine 3-monooxygenase (KMO) in complex with competitive inhibitor No. 9
Authors : Levy, C.W.; Leys, D.
Deposited on : 2018-02-08
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	7% 86% 11% .
1	B	460	4% 88% 9% ..

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 14997 atoms, of which 7048 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kynurenine 3-monooxygenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	445	7004	2202	3490	655	639	18	0	10	0
1	B	447	6966	2191	3468	652	637	18	0	4	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	251	SER	CYS	engineered mutation	UNP Q84HF5
A	460	SER	-	expression tag	UNP Q84HF5
B	251	SER	CYS	engineered mutation	UNP Q84HF5
B	460	SER	-	expression tag	UNP Q84HF5

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 84	C 27	H 31	N 9	O 15	P 2	0	0
2	B	1	Total 84	C 27	H 31	N 9	O 15	P 2	0	0

- Molecule 3 is 5-[2,3-bis(chloranyl)phenyl]furan-2-carboxylic acid (CCD ID: E0T) (formula: $C_{11}H_6Cl_2O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	Cl	O	0	0
			16	11	2	3		
3	B	1	Total	C	Cl	O	0	0
			16	11	2	3		

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			24	6	14	4		
4	B	1	Total	C	H	O	0	0
			24	6	14	4		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	402	Total	O	0	0
			402	402		
6	B	376	Total	O	0	0
			376	376		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Kynurenine 3-monooxygenase



• Molecule 1: Kynurenine 3-monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.80Å 52.98Å 135.76Å 90.00° 103.83° 90.00°	Depositor
Resolution (Å)	67.78 – 1.65 67.78 – 1.65	Depositor EDS
% Data completeness (in resolution range)	98.2 (67.78-1.65) 91.0 (67.78-1.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 1.65Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.164 , 0.213 0.166 , 0.215	Depositor DCC
R_{free} test set	1976 reflections (1.70%)	wwPDB-VP
Wilson B-factor (Å ²)	11.8	Xtriage
Anisotropy	0.529	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14997	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.83 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.7054e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: E0T, CA, PGE, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/3616	0.59	0/4909
1	B	0.43	0/3581	0.55	0/4858
All	All	0.46	0/7197	0.57	0/9767

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3514	3490	3475	34	0
1	B	3498	3468	3473	32	0
2	A	53	31	31	0	0
2	B	53	31	31	0	0
3	A	16	0	0	0	0
3	B	16	0	0	0	0
4	A	10	14	14	1	0
4	B	10	14	14	2	0
5	A	1	0	0	0	0
6	A	402	0	0	12	2
6	B	376	0	0	10	0
All	All	7949	7048	7038	66	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:38:ARG:NH2	6:B:601:HOH:O	2.02	0.93
1:A:281:GLN:OE1	6:A:601:HOH:O	1.88	0.91
1:A:156:LYS:HZ1	4:A:503:PGE:H32	1.52	0.74
1:A:271:ASP:OD1	6:A:602:HOH:O	2.08	0.72
1:B:421:GLN:OE1	6:B:602:HOH:O	2.11	0.68
1:A:448:GLU:OE2	1:A:451:ARG:NH2	2.24	0.68
1:B:383:LEU:HD23	1:B:431:VAL:HG13	1.74	0.68
1:A:47:ALA:O	6:A:603:HOH:O	2.12	0.68
1:A:417:MET:O	1:A:421:GLN:HG3	1.93	0.67
1:B:194:GLU:OE2	6:B:603:HOH:O	2.14	0.65
1:A:6:ALA:N	6:A:607:HOH:O	2.31	0.64
1:B:76:GLU:OE2	1:B:112:ARG:NH2	2.35	0.59
1:B:196:GLN:NE2	6:B:609:HOH:O	2.35	0.59
1:A:147:SER:HB3	1:A:153:ARG:HD3	1.85	0.58
1:A:437:LEU:HD12	1:A:440:ILE:HD12	1.87	0.57
1:A:257:ASP:OD2	6:A:604:HOH:O	2.17	0.57
1:A:390:ILE:HD12	1:A:437:LEU:HD21	1.87	0.57
1:B:188:GLN:O	1:B:290:LYS:NZ	2.39	0.56
1:A:244:ALA:O	1:A:245:ALA:HB3	2.06	0.56
1:B:367:GLU:OE2	6:B:604:HOH:O	2.18	0.55
1:B:183:GLU:HG2	1:B:296:LEU:HD22	1.90	0.54
1:B:112:ARG:HG3	6:B:915:HOH:O	2.07	0.54
1:B:290:LYS:HD3	1:B:291:LEU:O	2.07	0.53
1:A:297:THR:OG1	6:A:605:HOH:O	2.19	0.51
1:B:141:ARG:NE	6:B:619:HOH:O	2.42	0.51
1:B:394:ARG:HB3	1:B:446:GLU:OE2	2.11	0.51
1:B:257:ASP:OD1	1:B:257:ASP:C	2.54	0.50
1:B:85:VAL:HG12	1:B:87:VAL:HG23	1.94	0.50
1:B:50:ARG:O	1:B:50:ARG:HG3	2.11	0.49
1:B:282:ASP:O	1:B:285:HIS:O	2.28	0.49
1:A:16:LEU:HD13	1:A:323[A]:MET:HG3	1.93	0.49
1:B:58:GLU:OE2	1:B:410[B]:ARG:NH1	2.47	0.47
1:B:290:LYS:HD3	1:B:291:LEU:N	2.29	0.47
1:B:7[B]:ARG:HG3	1:B:30:TRP:CE2	2.49	0.47
1:B:335:GLU:HG3	6:B:815:HOH:O	2.15	0.46
1:A:153:ARG:O	1:A:154:LEU:HD23	2.17	0.45
1:A:214:ILE:HD12	1:A:222:CYS:O	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:MET:HE3	1:A:223:ILE:HG23	1.98	0.45
1:B:51:SER:OG	1:B:109:ASN:ND2	2.50	0.45
1:A:112:ARG:HG3	6:A:925:HOH:O	2.17	0.45
1:A:424:GLU:HG2	6:A:906:HOH:O	2.17	0.45
1:B:28:ASN:ND2	6:B:608:HOH:O	2.35	0.45
1:B:52:ILE:HD11	1:B:110:ARG:HH21	1.81	0.44
1:A:335:GLU:HG3	6:A:858:HOH:O	2.16	0.44
1:A:192:TYR:HA	1:A:236:LEU:O	2.17	0.44
1:B:71:ARG:NH1	6:B:610:HOH:O	2.38	0.44
1:A:243:PRO:HD3	6:A:734:HOH:O	2.18	0.44
1:A:411:LEU:HD11	1:A:415:GLN:CD	2.43	0.43
1:A:161:LEU:O	1:A:305:ALA:HA	2.19	0.43
1:B:156:LYS:HE3	4:B:503:PGE:H32	2.00	0.43
1:A:244:ALA:O	1:A:245:ALA:CB	2.67	0.43
1:A:200:GLU:H	1:A:200:GLU:CD	2.27	0.43
1:A:52:ILE:HD11	1:A:194[A]:GLU:OE1	2.19	0.43
1:A:71:ARG:NH2	1:A:75:ALA:HB2	2.34	0.43
1:B:144:LEU:HD12	1:B:144:LEU:C	2.43	0.42
1:A:434:HIS:ND1	1:A:439:SER:OG	2.38	0.42
1:A:154:LEU:HD11	6:A:926:HOH:O	2.18	0.42
1:A:373:SER:HA	6:A:787:HOH:O	2.19	0.42
1:B:192:TYR:HA	1:B:236:LEU:O	2.20	0.41
1:B:256:VAL:O	1:B:257:ASP:OD1	2.39	0.41
1:A:224:ALA:HB2	1:A:234:VAL:HG22	2.03	0.41
1:B:285:HIS:O	1:B:286:HIS:HB2	2.21	0.41
1:A:183:GLU:HG2	1:A:296:LEU:HD22	2.02	0.41
1:A:182:GLY:O	1:A:297:THR:HG23	2.21	0.40
1:B:97:TYR:HB3	1:B:408:PHE:CZ	2.56	0.40
1:B:156:LYS:NZ	4:B:503:PGE:H52	2.36	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:641:HOH:O	6:A:786:HOH:O[2_252]	1.90	0.30
6:A:641:HOH:O	6:A:680:HOH:O[2_252]	2.08	0.12

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	451/460 (98%)	441 (98%)	9 (2%)	1 (0%)	43	27
1	B	446/460 (97%)	433 (97%)	12 (3%)	1 (0%)	43	27
All	All	897/920 (98%)	874 (97%)	21 (2%)	2 (0%)	43	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	286	HIS
1	A	246	GLN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/371 (99%)	366 (100%)	2 (0%)	81	72
1	B	366/371 (99%)	363 (99%)	3 (1%)	73	60
All	All	734/742 (99%)	729 (99%)	5 (1%)	76	64

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ARG
1	A	281	GLN
1	B	257	ASP
1	B	290	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	389	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	GLN
1	A	95	GLN
1	A	131	ASN
1	A	206	ASN
1	A	217	HIS
1	A	304	GLN
1	B	321	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	E0T	B	502	-	17,17,17	1.24	1 (5%)	23,24,24	1.48	1 (4%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	E0T	A	502	-	17,17,17	1.49	2 (11%)	23,24,24	1.45	5 (21%)
4	PGE	B	503	-	9,9,9	0.29	0	8,8,8	0.57	0
2	FAD	B	501	-	58,58,58	0.58	1 (1%)	85,89,89	0.43	0
2	FAD	A	501	-	58,58,58	0.63	1 (1%)	85,89,89	0.49	0
4	PGE	A	503	-	9,9,9	0.34	0	8,8,8	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	E0T	B	502	-	-	0/8/8/8	0/2/2/2
3	E0T	A	502	-	-	0/8/8/8	0/2/2/2
4	PGE	B	503	-	-	5/7/7/7	-
2	FAD	B	501	-	-	1/34/50/50	0/6/6/6
2	FAD	A	501	-	-	1/34/50/50	0/6/6/6
4	PGE	A	503	-	-	2/7/7/7	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	E0T	O16-C07	-3.44	1.32	1.38
3	A	502	E0T	O16-C07	-2.77	1.33	1.38
2	B	501	FAD	C10-N1	-2.38	1.28	1.33
3	A	502	E0T	C12-C14	-2.33	1.36	1.39
2	A	501	FAD	P-O3P	-2.25	1.57	1.59

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	E0T	C06-C05-C04	-3.36	103.94	106.94
3	A	502	E0T	C06-C05-C04	-2.47	104.74	106.94
3	A	502	E0T	C14-C12-CL1	-2.42	118.28	120.51
3	A	502	E0T	C11-C12-C14	2.42	123.42	120.52
3	A	502	E0T	C12-C14-CL2	-2.25	117.17	120.02
3	A	502	E0T	C08-C14-CL2	2.10	123.11	120.11

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	503	PGE	O2-C3-C4-O3
4	B	503	PGE	O2-C3-C4-O3
4	B	503	PGE	O1-C1-C2-O2
4	B	503	PGE	C3-C4-O3-C5
4	B	503	PGE	C6-C5-O3-C4
4	B	503	PGE	C1-C2-O2-C3
2	A	501	FAD	C1'-C2'-C3'-O3'
2	B	501	FAD	C1'-C2'-C3'-O3'
4	A	503	PGE	O1-C1-C2-O2


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	503	PGE	2	0
4	A	503	PGE	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand E0T B 502



Ligand E0T A 502







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	445/460 (96%)	-0.13	31 (6%) 22 25	5, 19, 51, 63	9 (2%)
1	B	447/460 (97%)	-0.11	18 (4%) 42 46	7, 22, 48, 65	4 (0%)
All	All	892/920 (96%)	-0.12	49 (5%) 30 34	5, 21, 49, 65	13 (1%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	248	ALA	5.9
1	A	379	PRO	4.9
1	A	245	ALA	4.7
1	A	89	GLY	4.2
1	A	437	LEU	3.9
1	A	381	TYR	3.8
1	A	373	SER	3.7
1	A	48	ARG	3.5
1	B	48	ARG	3.5
1	B	457	SER	3.4
1	B	50	ARG	3.3
1	A	380	THR	3.3
1	B	256	VAL	3.2
1	A	248	ALA	3.1
1	A	154	LEU	3.1
1	B	49	GLY	3.1
1	A	244	ALA	3.0
1	A	285	HIS	3.0
1	A	90	THR	2.8
1	A	382	LEU	2.8
1	B	243	PRO	2.8
1	B	285	HIS	2.8
1	A	246	GLN	2.8
1	A	438	THR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	283	PHE	2.8
1	A	93	ASN	2.7
1	A	99	ARG	2.6
1	B	377	ALA	2.5
1	A	50	ARG	2.5
1	A	432	ALA	2.4
1	A	49	GLY	2.4
1	A	247	PRO	2.4
1	A	91	PRO	2.3
1	B	99	ARG	2.3
1	A	372	MET	2.3
1	A	385	ARG	2.3
1	B	51	SER	2.3
1	A	71	ARG	2.2
1	A	94	LEU	2.2
1	A	243	PRO	2.2
1	B	445	VAL	2.2
1	A	439	SER	2.2
1	B	437	LEU	2.2
1	A	433	ASN	2.2
1	B	397	THR	2.2
1	A	383	LEU	2.1
1	B	290	LYS	2.1
1	B	389	GLN	2.0
1	B	154	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PGE	A	503	10/10	0.84	0.14	37,45,53,53	0
4	PGE	B	503	10/10	0.85	0.14	39,47,52,53	0
3	E0T	A	502	16/16	0.97	0.07	21,24,26,27	0
3	E0T	B	502	16/16	0.99	0.05	15,18,21,22	0
2	FAD	B	501	53/53	0.99	0.04	8,11,23,23	0
2	FAD	A	501	53/53	0.99	0.04	5,9,16,17	0
5	CA	A	504	1/1	0.99	0.23	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around E0T A 502:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around E0T B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.