



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 11:20 PM UTC

PDB ID : 7FO2 / pdb_00007fo2
Title : PanDDA analysis group deposition – Aar2/RNaseH in complex with fragment P07G04 from the F2X-Universal Library
Authors : Barthel, T.; Wollenhaupt, J.; Lima, G.M.A.; Wahl, M.C.; Weiss, M.S.
Deposited on : 2022-08-26
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	21% 83% 9% 8%
2	B	308	23% 78% 17% • •

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9209 atoms, of which 4530 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	237	4068	1287	2060	336	373	12	0	21	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	300	5044	1654	2464	421	485	20	11	17	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	LEU	conflict	UNP P32357
B	167	SER	LYS	conflict	UNP P32357
B	?	-	LEU	deletion	UNP P32357
B	?	-	GLN	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	GLY	deletion	UNP P32357
B	?	-	SER	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ASN	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	170	SER	ASP	conflict	UNP P32357

- Molecule 3 is 3-[(3-bromophenyl)methanesulfonyl]propanoic acid (CCD ID: W0X) (formula: $C_{10}H_{11}BrO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	Br	C	O	S	0	0
			16	1	10	4	1		
3	B	1	Total	Br	C	O	S	0	0
			16	1	10	4	1		

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	H	O	S	0	0
			10	2	6	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	31	Total	O	0	0
			31	31		
5	B	24	Total	O	0	0
			24	24		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pre-mRNA-splicing factor 8



• Molecule 2: A1 cistron-splicing factor AAR2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.38Å 82.26Å 93.88Å 90.00° 108.08° 90.00°	Depositor
Resolution (Å)	44.62 – 2.01 44.62 – 2.01	Depositor EDS
% Data completeness (in resolution range)	98.8 (44.62-2.01) 99.3 (44.62-2.01)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.96 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.241 , 0.294 0.275 , 0.319	Depositor DCC
R_{free} test set	2100 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	47.4	Xtriage
Anisotropy	0.118	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9209	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: W0X, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	2/2149 (0.1%)	0.85	0/2911
2	B	0.98	10/2739 (0.4%)	1.09	14/3699 (0.4%)
All	All	0.91	12/4888 (0.2%)	0.99	14/6610 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	5	PRO	N-CA	16.09	1.65	1.46
2	B	7	THR	C-O	-9.71	1.12	1.24
2	B	70	GLN	C-O	-8.85	1.13	1.24
2	B	4	VAL	C-N	7.68	1.43	1.33
2	B	5	PRO	C-O	-6.83	1.16	1.23
2	B	91	ASN	C-O	-6.42	1.16	1.24
2	B	250	ILE	CG1-CD1	6.17	1.75	1.51
1	A	1940	MET	SD-CE	-5.96	1.64	1.79
2	B	88	LYS	C-O	-5.86	1.17	1.24
2	B	90	GLU	C-O	-5.32	1.17	1.24
1	A	1954	ILE	CG1-CD1	5.18	1.72	1.51
2	B	83	GLU	CD-OE2	-5.12	1.15	1.25

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	VAL	CA-C-N	14.39	135.40	120.14
2	B	4	VAL	C-N-CA	14.39	135.40	120.14
2	B	5	PRO	CB-CA-C	10.91	126.24	110.63
2	B	5	PRO	CA-N-CD	-7.83	101.04	112.00
2	B	6	PHE	CA-C-O	-7.30	112.48	120.36
2	B	86	GLY	CA-C-O	-7.00	113.84	121.05
2	B	7	THR	N-CA-C	-6.90	102.83	111.11
2	B	86	GLY	O-C-N	5.83	127.90	122.13
2	B	70	GLN	CB-CA-C	-5.68	99.43	109.70
2	B	18	ASP	CB-CA-C	-5.42	110.31	116.54
2	B	86	GLY	CA-C-N	5.31	127.84	120.29
2	B	86	GLY	C-N-CA	5.31	127.84	120.29
2	B	106	PRO	N-CA-CB	-5.19	98.94	102.73
2	B	89	PHE	CA-C-O	-5.13	115.61	120.90

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	54[A]	MET	Peptide
2	B	54[B]	MET	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2008	2060	1974	18	0
2	B	2580	2464	2398	41	0
3	B	32	0	0	2	0
4	B	4	6	6	0	0
5	A	31	0	0	2	0
5	B	24	0	0	3	0
All	All	4679	4530	4378	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:250:ILE:CD1	2:B:250:ILE:CG1	1.75	1.59
1:A:2069:VAL:O	5:A:2101:HOH:O	1.96	0.81
2:B:111:ASP:OD2	5:B:501:HOH:O	2.11	0.69
2:B:17:ILE:CD1	2:B:44[B]:ILE:CG1	2.71	0.69
1:A:2069:VAL:O	1:A:2069:VAL:HG12	1.94	0.66
2:B:68:TYR:CD2	3:B:401:W0X:C3	2.79	0.65
2:B:17:ILE:HD12	2:B:44[B]:ILE:HG13	1.80	0.64
2:B:17:ILE:HD13	2:B:44[B]:ILE:HG12	1.82	0.62
2:B:250:ILE:CD1	2:B:250:ILE:CB	2.74	0.62
2:B:77:LEU:HD21	2:B:79:LYS:HE3	1.83	0.61
2:B:17:ILE:HD13	2:B:44[B]:ILE:CG1	2.31	0.60
2:B:17:ILE:CD1	2:B:44[B]:ILE:HG13	2.30	0.60
1:A:1962:ARG:HD3	1:A:1962:ARG:H	1.66	0.60
1:A:1843:LEU:HA	1:A:1849:LYS:HD2	1.87	0.57
2:B:277:GLU:CD	2:B:277:GLU:H	2.12	0.57
2:B:17:ILE:CD1	2:B:44[B]:ILE:HG12	2.34	0.57
2:B:17:ILE:HD12	2:B:44[B]:ILE:CG1	2.35	0.56
2:B:17:ILE:O	2:B:17:ILE:HG23	2.06	0.56
1:A:1848:ILE:H	1:A:1931[A]:LYS:HZ1	1.54	0.56
1:A:2058:LEU:C	1:A:2058:LEU:HD23	2.32	0.55
2:B:68:TYR:CE2	3:B:401:W0X:C3	2.89	0.55
1:A:1878:CYS:SG	1:A:1969:MET:HE3	2.48	0.54
2:B:96:PHE:HB2	2:B:102:MET:HE3	1.91	0.53
1:A:1940:MET:C	1:A:1943:PRO:HD2	2.35	0.52
2:B:225:GLN:O	2:B:229:LEU:HG	2.11	0.50
2:B:190:ARG:HG3	2:B:203[B]:TYR:CE2	2.48	0.49
2:B:91:ASN:C	2:B:91:ASN:ND2	2.70	0.49
2:B:131:ARG:HA	5:B:502:HOH:O	2.12	0.48
2:B:290:ASN:HD21	2:B:317:LEU:HD23	1.78	0.48
2:B:177[B]:ASN:CG	2:B:177[B]:ASN:O	2.57	0.48
1:A:1962:ARG:O	1:A:1962:ARG:HG2	2.15	0.47
2:B:285:ASN:OD1	2:B:288:VAL:HG23	2.17	0.45
2:B:48:HIS:NE2	2:B:56:TYR:OH	2.38	0.45
1:A:2069:VAL:O	1:A:2069:VAL:CG1	2.65	0.44
2:B:37:PRO:HD3	2:B:105:TYR:CD1	2.53	0.44
2:B:44[A]:ILE:O	2:B:44[A]:ILE:HG23	2.17	0.44
2:B:287:ARG:O	2:B:291:ILE:HD13	2.18	0.44
1:A:2023:LYS:HA	1:A:2026:LEU:HD12	2.00	0.44
2:B:59:TRP:CE2	2:B:226:PHE:HE1	2.36	0.43
1:A:2022:ALA:O	1:A:2026:LEU:HD12	2.18	0.43
2:B:147:VAL:HG23	2:B:177[A]:ASN:HA	2.00	0.43
2:B:19:GLN:OE1	2:B:19:GLN:N	2.42	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1835:MET:HE3	1:A:1835:MET:HB3	1.90	0.41
1:A:2069:VAL:HG12	5:A:2101:HOH:O	2.19	0.41
2:B:1:MET:HB3	2:B:35:ASP:HA	2.03	0.41
1:A:2015:LEU:HD23	1:A:2022:ALA:HB3	2.03	0.41
2:B:290:ASN:ND2	2:B:317:LEU:HD23	2.35	0.41
1:A:1993:ASP:HB3	1:A:2038:HIS:CD2	2.57	0.40
2:B:80:MET:C	2:B:81:MET:HG3	2.46	0.40
2:B:17:ILE:O	2:B:17:ILE:CG2	2.69	0.40
2:B:40:HIS:HD2	5:B:520:HOH:O	2.05	0.40
1:A:2047:GLN:O	1:A:2051:ILE:HG12	2.22	0.40
2:B:51:ASN:OD1	2:B:53:SER:HB2	2.22	0.40
2:B:190:ARG:HG3	2:B:203[B]:TYR:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/258 (100%)	250 (97%)	8 (3%)	0	100	100
2	B	315/308 (102%)	298 (95%)	17 (5%)	0	100	100
All	All	573/566 (101%)	548 (96%)	25 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/233 (102%)	235 (99%)	2 (1%)	73	80
2	B	294/284 (104%)	283 (96%)	11 (4%)	30	30
All	All	531/517 (103%)	518 (98%)	13 (2%)	40	47

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2027	LEU
1	A	2033	THR
2	B	5	PRO
2	B	7	THR
2	B	77	LEU
2	B	91	ASN
2	B	101	MET
2	B	106	PRO
2	B	107	LYS
2	B	258	LYS
2	B	295	SER
2	B	316	LEU
2	B	317	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1907	GLN
1	A	1947	HIS
2	B	2	ASN
2	B	47	GLN
2	B	91	ASN
2	B	94	HIS
2	B	150	ASN
2	B	259	HIS
2	B	290	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	DMS	B	403	-	3,3,3	0.26	0	3,3,3	0.44	0
3	W0X	B	402	-	16,16,16	5.50	4 (25%)	20,22,22	2.27	7 (35%)
3	W0X	B	401	-	16,16,16	5.90	7 (43%)	20,22,22	1.92	4 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	W0X	B	402	-	-	3/11/11/11	0/1/1/1
3	W0X	B	401	-	-	3/11/11/11	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	W0X	C7-S	-21.08	1.55	1.78
3	B	401	W0X	C7-S	-20.41	1.56	1.78
3	B	401	W0X	O1-S	-8.89	1.27	1.44
3	B	401	W0X	O-S	-4.70	1.35	1.44
3	B	402	W0X	C6-S	-3.79	1.75	1.79

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	W0X	C6-C2	3.41	1.57	1.51
3	B	401	W0X	C8-C7	2.98	1.55	1.52
3	B	401	W0X	C3-C4	-2.32	1.33	1.38
3	B	402	W0X	O-S	-2.32	1.39	1.44
3	B	401	W0X	O2-C9	-2.06	1.24	1.30
3	B	402	W0X	C1-C2	-2.02	1.34	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	W0X	O-S-O1	5.42	124.90	117.55
3	B	401	W0X	O1-S-C7	4.83	115.72	108.51
3	B	402	W0X	O-S-C6	-4.31	101.45	108.04
3	B	401	W0X	O1-S-C6	-4.12	101.74	108.04
3	B	402	W0X	O1-S-C7	-4.06	102.46	108.51
3	B	402	W0X	O3-C9-C8	-2.85	114.06	123.09
3	B	402	W0X	C6-C2-C1	-2.71	117.29	120.56
3	B	401	W0X	O-S-O1	2.70	121.21	117.55
3	B	402	W0X	C6-S-C7	2.39	112.39	105.36
3	B	401	W0X	O3-C9-C8	-2.05	116.59	123.09
3	B	402	W0X	BR-C4-C5	-2.03	116.53	119.30

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	402	W0X	S-C7-C8-C9
3	B	402	W0X	C7-C8-C9-O3
3	B	402	W0X	C7-C8-C9-O2
3	B	401	W0X	C7-C8-C9-O3
3	B	401	W0X	C7-C8-C9-O2
3	B	401	W0X	C1-C2-C6-S


There are no ring outliers.


1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	W0X	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/258 (91%)	1.36	54 (22%)  	26, 73, 117, 192	12 (5%)
2	B	300/308 (97%)	1.39	70 (23%)  	23, 73, 124, 167	9 (3%)
All	All	537/566 (94%)	1.38	124 (23%)  	23, 73, 123, 192	21 (3%)

All (124) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	106	PRO	5.6
1	A	1979[A]	MET	5.6
2	B	152	LEU	5.2
1	A	1840	TYR	4.9
1	A	1898[A]	VAL	4.7
1	A	2069	VAL	4.6
2	B	54[A]	MET	4.6
1	A	1834	ALA	4.3
1	A	2030	PRO	4.2
2	B	1	MET	4.2
2	B	20	TYR	4.1
1	A	1835	MET	4.1
1	A	1963	LEU	4.0
2	B	22	PHE	4.0
2	B	288	VAL	4.0
2	B	297	PHE	3.9
1	A	2027	LEU	3.9
1	A	2055	MET	3.9
1	A	1961[A]	LEU	3.8
2	B	199	LEU	3.8
1	A	1967	ALA	3.7
2	B	284	LEU	3.7
1	A	2017[A]	THR	3.6
2	B	279	TYR	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	170	SER	3.5
1	A	1991	ILE	3.4
2	B	7	THR	3.4
2	B	174	HIS	3.4
2	B	108	ILE	3.4
2	B	275	LEU	3.4
2	B	203[A]	TYR	3.3
2	B	104	SER	3.3
2	B	195	MET	3.2
2	B	53	SER	3.2
2	B	264	LEU	3.1
2	B	124	ASP	3.1
2	B	313	TYR	3.1
2	B	21	SER	3.1
2	B	107	LYS	3.1
1	A	1833	GLY	3.1
1	A	2005	PHE	3.1
1	A	1999	ILE	3.1
2	B	38	ILE	3.0
1	A	2056[A]	ARG	3.0
2	B	302	LEU	3.0
2	B	282	ILE	2.9
2	B	289	TRP	2.9
2	B	181	ILE	2.9
1	A	1862	VAL	2.8
2	B	232	MET	2.8
2	B	101	MET	2.8
1	A	1863	HIS	2.8
1	A	1940	MET	2.8
1	A	2031	THR	2.8
2	B	308	ILE	2.8
1	A	1965	PHE	2.7
2	B	294	TYR	2.7
1	A	2006	SER	2.7
1	A	2061	THR	2.7
2	B	316	LEU	2.7
1	A	2038	HIS	2.7
2	B	256	VAL	2.6
2	B	292	CYS	2.6
2	B	171	ASP	2.6
1	A	1860	VAL	2.6
2	B	19	GLN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2033	THR	2.5
1	A	2063	TYR	2.5
1	A	1836	ASN	2.5
2	B	5	PRO	2.5
2	B	120	PHE	2.5
2	B	263	LYS	2.5
2	B	86	GLY	2.5
2	B	77	LEU	2.4
1	A	2034	ILE	2.4
2	B	92	ILE	2.4
1	A	1866	PHE	2.4
1	A	2066	LYS	2.4
2	B	3	THR	2.4
2	B	10	PRO	2.4
2	B	70	GLN	2.3
2	B	283	LEU	2.3
2	B	15	ILE	2.3
2	B	259	HIS	2.3
1	A	1878	CYS	2.3
1	A	2067	TYR	2.3
1	A	1844	PHE	2.3
2	B	110	GLU	2.3
2	B	276	PRO	2.3
2	B	105	TYR	2.3
1	A	2053[A]	SER	2.3
1	A	1951	PHE	2.3
1	A	2032	ILE	2.3
1	A	1934	ILE	2.2
1	A	1946	VAL	2.2
2	B	246	MET	2.2
1	A	2037	TYR	2.2
2	B	96	PHE	2.2
2	B	131	ARG	2.2
1	A	1864	LYS	2.2
2	B	132	LYS	2.2
2	B	189	ILE	2.2
2	B	64[A]	MET	2.2
2	B	172	PRO	2.2
2	B	109	ASP	2.2
1	A	1843	LEU	2.2
1	A	2036	SER	2.2
2	B	213	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	193	HIS	2.1
1	A	1838	SER	2.1
1	A	2068	ASN	2.1
2	B	150	ASN	2.1
2	B	211[A]	GLN	2.1
1	A	1841	ALA	2.1
1	A	1950	ASP	2.1
1	A	1998	ARG	2.1
2	B	237	TYR	2.1
1	A	1837	SER	2.1
2	B	100	GLN	2.0
1	A	2023	LYS	2.0
1	A	1846	ASN	2.0
2	B	40	HIS	2.0
2	B	37	PRO	2.0
1	A	1901	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	DMS	B	403	4/4	0.77	0.33	20,20,20,20	0
3	W0X	B	402	16/16	0.91	0.22	20,20,20,20	16
3	W0X	B	401	16/16	0.91	0.21	20,20,20,20	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around W0X B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around W0X B 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.