



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:20 PM UTC

PDB ID : 9FOS / pdb_00009fos
Title : The structure of ornithine decarboxylase from Leishmania infantum in complex with PLP and DFMO
Authors : Fiorillo, A.; Ilari, A.; Bibi, A.
Deposited on : 2024-06-12
Resolution : 3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	727	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3476 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ornithine decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	440	Total	C	N	O	S	0	0	0
			3454	2203	583	647	21			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP E9AGB5
A	-18	GLY	-	expression tag	UNP E9AGB5
A	-17	SER	-	expression tag	UNP E9AGB5
A	-16	SER	-	expression tag	UNP E9AGB5
A	-15	HIS	-	expression tag	UNP E9AGB5
A	-14	HIS	-	expression tag	UNP E9AGB5
A	-13	HIS	-	expression tag	UNP E9AGB5
A	-12	HIS	-	expression tag	UNP E9AGB5
A	-11	HIS	-	expression tag	UNP E9AGB5
A	-10	HIS	-	expression tag	UNP E9AGB5
A	-9	SER	-	expression tag	UNP E9AGB5
A	-8	SER	-	expression tag	UNP E9AGB5
A	-7	GLY	-	expression tag	UNP E9AGB5
A	-6	LEU	-	expression tag	UNP E9AGB5
A	-5	VAL	-	expression tag	UNP E9AGB5
A	-4	PRO	-	expression tag	UNP E9AGB5
A	-3	ARG	-	expression tag	UNP E9AGB5
A	-2	GLY	-	expression tag	UNP E9AGB5
A	-1	SER	-	expression tag	UNP E9AGB5
A	0	HIS	-	expression tag	UNP E9AGB5

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is ALPHA-DIFLUOROMETHYLORNITHINE (CCD ID: DMO) (formula: $C_6H_{12}F_2N_2O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			7	5	2		

4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	117.87Å 117.87Å 82.17Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.61 – 3.30 38.61 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.61-3.30) 99.7 (38.61-3.30)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.43 (at 3.32Å)	Xtriage
Refinement program	REFMAC 5.8.0425, REFMAC 5.8.0425	Depositor
R, R_{free}	0.193 , 0.240 0.199 , 0.245	Depositor DCC
R_{free} test set	503 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	104.8	Xtriage
Anisotropy	0.482	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 83.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.042 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3476	wwPDB-VP
Average B, all atoms (Å ²)	125.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMO, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	4/3534 (0.1%)	1.72	61/4797 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	502	TYR	C-O	7.90	1.33	1.24
1	A	258	PRO	C-O	5.93	1.30	1.23
1	A	330	ILE	CB-CG1	-5.76	1.42	1.53
1	A	375	ARG	NE-CZ	5.05	1.38	1.33

All (61) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	211	THR	CA-CB-OG1	11.06	126.20	109.60
1	A	353	THR	CA-CB-OG1	-9.62	95.17	109.60
1	A	670	PHE	CB-CA-C	9.20	125.53	111.18
1	A	350	THR	CA-CB-OG1	-8.40	97.00	109.60
1	A	666	ALA	O-C-N	7.62	129.92	122.07
1	A	635	ASP	CA-CB-CG	7.26	119.86	112.60
1	A	486	GLU	CB-CA-C	6.98	121.72	110.96
1	A	285	PHE	CB-CA-C	-6.95	98.77	109.89
1	A	412	VAL	CB-CA-C	-6.86	104.86	112.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	ARG	N-CA-CB	-6.69	100.31	110.01
1	A	218	ARG	CA-CB-CG	6.48	127.06	114.10
1	A	653	ASP	CA-CB-CG	6.39	118.99	112.60
1	A	350	THR	OG1-CB-CG2	6.36	122.01	109.30
1	A	556	PHE	CA-CB-CG	-6.14	107.66	113.80
1	A	306	PHE	CA-CB-CG	-6.10	107.70	113.80
1	A	377	LYS	N-CA-CB	-6.08	101.67	110.36
1	A	502	TYR	N-CA-CB	-6.07	100.60	109.82
1	A	492	ASP	CA-CB-CG	6.04	118.64	112.60
1	A	502	TYR	CB-CA-C	6.00	120.45	110.92
1	A	327	ASP	CA-CB-CG	5.96	118.56	112.60
1	A	677	ARG	CD-NE-CZ	5.95	132.73	124.40
1	A	155	GLY	CA-C-O	-5.90	116.42	122.26
1	A	409	ASN	CB-CA-C	5.86	119.32	111.82
1	A	457	GLY	O-C-N	5.84	130.30	122.70
1	A	158	THR	OG1-CB-CG2	5.83	120.97	109.30
1	A	519	THR	CA-CB-OG1	-5.81	100.88	109.60
1	A	135	PHE	O-C-N	5.80	128.27	122.12
1	A	638	ASP	CA-CB-CG	5.76	118.36	112.60
1	A	681	TRP	N-CA-C	5.75	118.06	109.25
1	A	160	GLU	N-CA-C	-5.75	104.39	111.40
1	A	316	MET	CG-SD-CE	5.75	113.55	100.90
1	A	152	PRO	N-CA-CB	5.70	108.11	103.32
1	A	646	PHE	CA-CB-CG	5.70	119.50	113.80
1	A	437	TYR	N-CA-C	-5.67	104.79	110.97
1	A	138	GLU	CB-CA-C	5.62	118.75	109.70
1	A	162	ARG	NE-CZ-NH1	-5.59	115.91	121.50
1	A	133	ASP	O-C-N	5.55	128.02	122.08
1	A	540	GLU	CB-CG-CD	5.48	121.91	112.60
1	A	449	LYS	CB-CG-CD	5.43	123.78	111.30
1	A	340	ASP	CA-CB-CG	5.39	117.99	112.60
1	A	484	LEU	O-C-N	5.39	128.29	122.15
1	A	562	ASP	CA-CB-CG	5.37	117.97	112.60
1	A	389	THR	OG1-CB-CG2	5.34	119.98	109.30
1	A	564	ALA	CA-C-O	-5.30	115.78	121.56
1	A	687	LEU	N-CA-CB	-5.29	101.54	110.49
1	A	257	ASP	CA-CB-CG	5.29	117.89	112.60
1	A	639	CYS	CB-CA-C	-5.25	101.74	110.14
1	A	553	TYR	OH-CZ-CE2	5.24	135.63	119.90
1	A	150	PRO	N-CA-CB	5.24	108.75	103.25
1	A	136	PHE	O-C-N	5.20	129.30	122.33
1	A	569	LEU	CA-C-O	-5.19	114.61	120.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	173	TYR	N-CA-C	-5.17	106.29	112.59
1	A	434	ARG	CB-CA-C	5.17	119.07	110.81
1	A	130	ASP	CA-CB-CG	5.15	117.75	112.60
1	A	644	GLN	N-CA-CB	5.11	119.71	110.72
1	A	365	ARG	CA-C-O	-5.10	115.65	120.90
1	A	411	THR	CA-C-O	-5.06	115.66	120.92
1	A	670	PHE	N-CA-CB	-5.05	104.21	111.54
1	A	386	SER	N-CA-C	5.02	116.49	108.41
1	A	218	ARG	CB-CA-C	5.01	118.75	110.88
1	A	372	ALA	N-CA-C	5.01	117.49	110.23

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	162	ARG	Sidechain
1	A	406	ARG	Sidechain
1	A	451	THR	Peptide
1	A	470	ASN	Peptide
1	A	489	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3454	0	3358	70	0
2	A	15	0	7	0	0
3	A	7	0	8	0	0
All	All	3476	0	3373	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:GLN:HA	1:A:141:GLN:OE1	1.79	0.80
1:A:451:THR:HB	1:A:452:ILE:HG13	1.66	0.76
1:A:435:ASP:O	1:A:439:VAL:HG23	1.90	0.72
1:A:521:ARG:HH21	1:A:540:GLU:HA	1.61	0.66
1:A:138:GLU:HB2	1:A:148:PHE:CE2	2.33	0.64
1:A:278:LEU:HB3	1:A:281:VAL:HG23	1.81	0.62
1:A:143:VAL:O	1:A:146:LEU:HB2	2.00	0.62
1:A:482:PRO:O	1:A:485:ALA:HB3	1.99	0.62
1:A:517:SER:HB3	1:A:651:LEU:HD11	1.82	0.62
1:A:350:THR:HG21	1:A:371:HIS:CE1	2.36	0.61
1:A:413:CYS:SG	1:A:452:ILE:HD12	2.41	0.60
1:A:645:PRO:O	1:A:646:PHE:HB2	2.01	0.60
1:A:332:ALA:O	1:A:334:PRO:HD3	2.01	0.60
1:A:313:GLU:HA	1:A:316:MET:HE3	1.83	0.59
1:A:259:PHE:HB3	1:A:678:ARG:HB3	1.84	0.59
1:A:288:LYS:NZ	1:A:309:ALA:HB2	2.19	0.57
1:A:429:TYR:O	1:A:433:VAL:HG23	2.04	0.57
1:A:395:LEU:O	1:A:398:VAL:HG23	2.05	0.57
1:A:180:PRO:HA	1:A:218:ARG:HH12	1.69	0.56
1:A:485:ALA:O	1:A:489:GLY:HA3	2.06	0.55
1:A:521:ARG:NH2	1:A:540:GLU:HA	2.20	0.55
1:A:260:TYR:CE1	1:A:510:LEU:HB2	2.42	0.53
1:A:158:THR:HG23	1:A:161:GLU:OE1	2.09	0.53
1:A:336:LYS:HE2	1:A:341:LEU:HD21	1.88	0.53
1:A:350:THR:HG23	1:A:369:SER:O	2.10	0.52
1:A:412:VAL:HG12	1:A:412:VAL:O	2.09	0.52
1:A:176:VAL:HG21	1:A:178:ARG:NH2	2.25	0.51
1:A:264:LEU:HD22	1:A:298:VAL:HG21	1.93	0.51
1:A:522:LEU:HB2	1:A:541:PRO:HB2	1.92	0.51
1:A:284:TYR:O	1:A:498:GLU:HA	2.10	0.51
1:A:268:VAL:HG22	1:A:302:LEU:HD11	1.92	0.51
1:A:480:ILE:HG22	1:A:484:LEU:CD1	2.42	0.50
1:A:493:VAL:HG23	1:A:493:VAL:O	2.12	0.49
1:A:543:GLU:OE1	1:A:643:LYS:HE2	2.11	0.49
1:A:452:ILE:HG22	1:A:453:LEU:N	2.27	0.49
1:A:540:GLU:N	1:A:541:PRO:HD3	2.27	0.49
1:A:481:ARG:N	1:A:482:PRO:CD	2.76	0.48
1:A:307:ASP:C	1:A:307:ASP:OD1	2.57	0.48
1:A:154:TYR:OH	1:A:275:ARG:HA	2.13	0.48
1:A:550:ASP:OD1	1:A:551:GLY:N	2.44	0.48
1:A:330:ILE:HD13	1:A:330:ILE:N	2.29	0.47
1:A:484:LEU:HD22	1:A:495:ILE:HD13	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:315:HIS:HE1	1:A:343:GLU:HG2	1.79	0.47
1:A:345:GLN:OE1	1:A:367:MET:HA	2.15	0.47
1:A:440:PHE:CE2	1:A:450:CYS:HB2	2.50	0.47
1:A:439:VAL:O	1:A:440:PHE:C	2.57	0.46
1:A:274:TRP:CD1	1:A:283:PRO:HG3	2.50	0.46
1:A:281:VAL:HG13	1:A:495:ILE:HG22	1.97	0.46
1:A:371:HIS:HB3	1:A:413:CYS:SG	2.56	0.46
1:A:315:HIS:CE1	1:A:343:GLU:HG2	2.51	0.45
1:A:637:MET:HE3	1:A:637:MET:HB2	1.68	0.45
1:A:409:ASN:OD1	1:A:409:ASN:O	2.34	0.45
1:A:149:SER:HB2	1:A:150:PRO:HD2	1.97	0.45
1:A:326:PRO:HB2	1:A:348:GLY:O	2.17	0.44
1:A:259:PHE:CB	1:A:678:ARG:HB3	2.48	0.44
1:A:176:VAL:HG22	1:A:176:VAL:O	2.18	0.44
1:A:350:THR:HG22	1:A:371:HIS:CG	2.52	0.44
1:A:354:VAL:HG23	1:A:359:GLU:HB3	2.00	0.43
1:A:481:ARG:HB2	1:A:482:PRO:HD3	2.01	0.43
1:A:176:VAL:HG21	1:A:178:ARG:CZ	2.50	0.42
1:A:177:THR:CG2	1:A:179:LEU:HD21	2.48	0.42
1:A:313:GLU:HA	1:A:316:MET:CE	2.50	0.42
1:A:263:ASP:O	1:A:265:GLY:N	2.53	0.42
1:A:289:SER:O	1:A:290:ASN:HB2	2.20	0.42
1:A:501:ARG:HA	1:A:662:SER:OG	2.19	0.42
1:A:519:THR:O	1:A:520:LEU:HD23	2.19	0.41
1:A:628:THR:HA	1:A:641:LEU:O	2.20	0.41
1:A:540:GLU:N	1:A:541:PRO:CD	2.84	0.41
1:A:646:PHE:CG	1:A:647:PRO:HD2	2.56	0.40
1:A:315:HIS:NE2	1:A:343:GLU:OE2	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	428/727 (59%)	390 (91%)	36 (8%)	2 (0%)	24 55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	490	GLY
1	A	321	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	372/592 (63%)	356 (96%)	16 (4%)	26 54

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	ASP
1	A	176	VAL
1	A	281	VAL
1	A	316	MET
1	A	350	THR
1	A	369	SER
1	A	381	SER
1	A	382	LYS
1	A	421	SER
1	A	426	GLN
1	A	449	LYS
1	A	451	THR
1	A	523	SER
1	A	552	LEU
1	A	672	ASN
1	A	686	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	225	GLN
1	A	441	GLN
1	A	513	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PLP	A	801	3	15,15,16	2.58	6 (40%)	21,22,23	3.34	9 (42%)
3	DMO	A	802	1,2	6,6,11	0.71	0	6,6,15	2.88	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	801	3	-	0/6/6/8	0/1/1/1
3	DMO	A	802	1,2	-	2/4/4/17	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	PLP	P-O4P	6.61	1.81	1.60
2	A	801	PLP	O3-C3	3.58	1.45	1.36
2	A	801	PLP	C5-C4	3.50	1.44	1.40
2	A	801	PLP	C4A-C4	-2.46	1.46	1.51
2	A	801	PLP	C3-C4	2.46	1.44	1.40
2	A	801	PLP	P-O2P	-2.43	1.45	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	PLP	O4P-C5A-C5	10.56	129.14	109.36
3	A	802	DMO	CBO-CAO-NO	-6.11	90.92	108.68
2	A	801	PLP	C6-C5-C4	-6.07	113.12	118.10
2	A	801	PLP	O3P-P-O4P	4.42	118.19	106.67
2	A	801	PLP	C3-C4-C5	3.96	123.33	118.59
2	A	801	PLP	O3-C3-C4	3.12	126.24	118.10
3	A	802	DMO	CO-CAO-CBO	3.00	120.68	111.99
2	A	801	PLP	O4P-P-O1P	2.79	113.99	106.44
2	A	801	PLP	O2P-P-O1P	-2.25	102.09	110.83
2	A	801	PLP	C6-N1-C2	2.19	123.17	119.20
2	A	801	PLP	C4-C3-C2	-2.08	116.78	119.89

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	802	DMO	CAO-CBO-CGO-CDO
3	A	802	DMO	NEO-CDO-CGO-CBO

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	440/727 (60%)	-0.65	0 100 100	68, 124, 166, 203	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PLP	A	801	15/16	0.92	0.10	77,125,177,182	0
3	DMO	A	802	7/12	0.97	0.10	78,92,112,112	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PLP A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DMO A 802:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.