



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:28 PM UTC

PDB ID : 1FP5 / pdb_00001fp5
Title : CRYSTAL STRUCTURE ANALYSIS OF THE HUMAN IGE-FC
CEPSILON3-CEPSILON4 FRAGMENT.
Authors : Wurzburg, B.A.; Garman, S.C.; Jardetzky, T.S.
Deposited on : 2000-08-30
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	13% 55% 32% 7% 6%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1763 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGE HEAVY CHAIN EPSILON-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	208	Total	C	N	O	S	0	0	0
			1618	1015	294	303	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	326	ALA	-	cloning artifact	UNP P01854
A	327	ASP	-	cloning artifact	UNP P01854
A	328	PRO	-	cloning artifact	UNP P01854
A	329	CYS	-	cloning artifact	UNP P01854


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	145	Total	O	0	0
			145	145		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: IGE HEAVY CHAIN EPSILON-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	105.60Å 105.60Å 47.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30 30.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-2.30) 99.7 (30.00-2.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.83 (at 2.31Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.242 , 0.270 0.242 , 0.271	Depositor DCC
R_{free} test set	1269 reflections (10.30%)	wwPDB-VP
Wilson B-factor (Å ²)	45.8	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1763	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/1660	1.17	21/2266 (0.9%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	456	SER	N-CA-C	13.36	130.23	109.52
1	A	458	ASP	N-CA-C	11.83	123.45	108.19
1	A	457	ARG	N-CA-C	-11.24	86.85	110.80
1	A	367	LYS	N-CA-C	9.19	123.20	108.41
1	A	366	SER	N-CA-C	-8.10	96.20	109.40
1	A	455	GLY	N-CA-C	-7.56	95.26	113.18
1	A	424	HIS	N-CA-C	7.55	121.74	112.23
1	A	353	SER	CA-C-N	7.25	127.23	119.76
1	A	353	SER	C-N-CA	7.25	127.23	119.76
1	A	467	GLN	N-CA-C	6.85	118.19	108.74
1	A	368	GLY	N-CA-C	-6.82	97.81	110.83
1	A	456	SER	CA-C-N	5.73	132.49	121.54
1	A	456	SER	C-N-CA	5.73	132.49	121.54
1	A	534	SER	N-CA-C	5.72	119.46	111.90
1	A	425	LEU	N-CA-C	5.39	116.68	109.83
1	A	453	TRP	CA-C-N	5.33	126.50	119.84
1	A	453	TRP	C-N-CA	5.33	126.50	119.84
1	A	396	THR	N-CA-C	5.15	116.45	107.49
1	A	462	LEU	N-CA-C	-5.12	101.53	109.72
1	A	529	GLU	N-CA-C	5.01	117.12	111.11
1	A	472	GLU	N-CA-C	5.00	119.01	113.01

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1618	0	1578	78	0
2	A	145	0	0	9	0
All	All	1763	0	1578	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:452:GLU:HG2	1:A:460:ARG:HG2	1.53	0.90
1:A:458:ASP:HA	2:A:132:HOH:O	1.79	0.83
1:A:377:ALA:HB3	1:A:415:THR:HG23	1.62	0.81
1:A:366:SER:OG	1:A:424:HIS:CE1	2.36	0.78
1:A:415:THR:HB	1:A:434:THR:CG2	2.16	0.76
1:A:366:SER:CB	1:A:424:HIS:HE1	2.00	0.74
1:A:348:LEU:HA	1:A:354:PRO:HG3	1.70	0.74
1:A:513:ARG:NH1	1:A:517:GLU:HG2	2.04	0.71
1:A:415:THR:HA	1:A:434:THR:HG22	1.72	0.71
1:A:366:SER:OG	1:A:424:HIS:HE1	1.73	0.71
1:A:512:THR:OG1	1:A:515:GLU:HG2	1.90	0.69
1:A:460:ARG:CZ	1:A:513:ARG:HG2	2.24	0.67
1:A:361:VAL:HG13	1:A:396:THR:HG23	1.77	0.66
1:A:532:SER:HB2	2:A:74:HOH:O	1.96	0.66
1:A:415:THR:HB	1:A:434:THR:HG22	1.77	0.65
1:A:502:GLY:HA2	2:A:130:HOH:O	1.96	0.63
1:A:372:LEU:HD22	1:A:420:VAL:HG22	1.79	0.63
1:A:480:HIS:O	1:A:481:ASN:HB2	1.98	0.63
1:A:377:ALA:CB	1:A:415:THR:HG23	2.29	0.62
1:A:393:ARG:HD3	2:A:13:HOH:O	2.01	0.61
1:A:499:LYS:O	1:A:501:SER:N	2.30	0.61
1:A:511:VAL:HB	1:A:515:GLU:HG3	1.83	0.60
1:A:440:ARG:HD3	1:A:471:PRO:HD3	1.82	0.60
1:A:392:GLN:C	1:A:394:ASN:H	2.10	0.59
1:A:513:ARG:HD3	1:A:513:ARG:O	2.01	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:422:HIS:CG	1:A:423:PRO:HD2	2.37	0.59
1:A:513:ARG:HH11	1:A:517:GLU:HG2	1.68	0.59
1:A:415:THR:HB	1:A:434:THR:HG21	1.85	0.59
1:A:366:SER:HB2	1:A:424:HIS:HE1	1.66	0.59
1:A:458:ASP:O	1:A:459:LYS:HG2	2.02	0.58
1:A:349:PHE:CG	1:A:535:GLN:HG3	2.38	0.58
1:A:394:ASN:HB3	2:A:91:HOH:O	2.03	0.57
1:A:440:ARG:NH1	1:A:471:PRO:HG2	2.19	0.57
1:A:427:ARG:HG3	2:A:140:HOH:O	2.05	0.57
1:A:478:TRP:O	1:A:479:LEU:HD23	2.05	0.57
1:A:366:SER:CB	1:A:424:HIS:CE1	2.85	0.56
1:A:415:THR:CB	1:A:434:THR:HG22	2.35	0.55
1:A:405:VAL:HG22	1:A:416:TYR:CZ	2.42	0.54
1:A:466:ILE:HB	1:A:505:VAL:HG12	1.90	0.53
1:A:422:HIS:ND1	1:A:423:PRO:HD2	2.23	0.53
1:A:415:THR:CA	1:A:434:THR:HG22	2.38	0.52
1:A:426:PRO:HG2	1:A:427:ARG:H	1.76	0.51
1:A:441:ALA:HB3	1:A:470:MET:HG2	1.92	0.51
1:A:535:GLN:HG2	2:A:21:HOH:O	2.10	0.50
1:A:534:SER:O	1:A:536:THR:HG22	2.10	0.50
1:A:460:ARG:HD3	1:A:516:TRP:CG	2.47	0.50
1:A:392:GLN:C	1:A:394:ASN:N	2.70	0.50
1:A:376:ARG:HG3	1:A:382:VAL:HG12	1.93	0.50
1:A:531:ALA:O	1:A:535:GLN:HA	2.11	0.50
1:A:392:GLN:O	1:A:394:ASN:N	2.45	0.50
1:A:420:VAL:HB	1:A:429:LEU:HB2	1.92	0.50
1:A:440:ARG:HB3	1:A:530:ALA:HB2	1.94	0.49
1:A:508:ARG:HH12	1:A:510:GLU:CD	2.22	0.48
1:A:525:ARG:HD3	1:A:538:GLN:OE1	2.14	0.47
1:A:452:GLU:HG2	1:A:460:ARG:CG	2.35	0.47
1:A:459:LYS:HD3	1:A:459:LYS:HA	1.65	0.47
1:A:363:LEU:HA	1:A:395:GLY:O	2.15	0.46
1:A:370:VAL:HG22	1:A:422:HIS:ND1	2.31	0.46
1:A:452:GLU:CG	1:A:460:ARG:HG2	2.34	0.45
1:A:336:VAL:HA	1:A:361:VAL:O	2.17	0.45
1:A:528:HIS:O	1:A:535:GLN:HB3	2.16	0.45
1:A:513:ARG:HD3	1:A:513:ARG:C	2.42	0.45
1:A:360:VAL:HG21	1:A:372:LEU:HD21	1.98	0.45
1:A:470:MET:HA	1:A:471:PRO:C	2.42	0.45
1:A:453:TRP:HA	1:A:454:PRO:HD2	1.77	0.45
1:A:372:LEU:CD2	1:A:420:VAL:HG22	2.47	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:372:LEU:HD12	1:A:401:SER:HB2	2.01	0.43
1:A:460:ARG:HB3	1:A:516:TRP:CD1	2.54	0.43
1:A:513:ARG:HH11	1:A:517:GLU:CG	2.31	0.43
1:A:516:TRP:HZ2	2:A:60:HOH:O	2.02	0.42
1:A:370:VAL:HG22	1:A:422:HIS:CE1	2.54	0.42
1:A:501:SER:OG	1:A:502:GLY:N	2.52	0.42
1:A:460:ARG:NH2	1:A:513:ARG:HG2	2.35	0.41
1:A:365:PRO:HB3	2:A:61:HOH:O	2.21	0.41
1:A:515:GLU:HB2	1:A:522:PHE:CZ	2.55	0.41
1:A:531:ALA:HB1	1:A:537:VAL:HG23	2.02	0.41
1:A:535:GLN:CD	1:A:535:GLN:N	2.79	0.41
1:A:349:PHE:CD2	1:A:535:GLN:HG3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	206/222 (93%)	189 (92%)	9 (4%)	8 (4%)	2 1

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	454	PRO
1	A	393	ARG
1	A	459	LYS
1	A	500	GLY
1	A	395	GLY
1	A	481	ASN
1	A	426	PRO
1	A	488	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	178/195 (91%)	172 (97%)	6 (3%)	32 49

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	351	ARG
1	A	405	VAL
1	A	415	THR
1	A	482	GLU
1	A	520	ASP
1	A	535	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	383	ASN
1	A	394	ASN
1	A	424	HIS
1	A	468	ASN
1	A	484	GLN
1	A	535	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	208/222 (93%)	0.86	29 (13%) 6 7	22, 49, 97, 111	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	424	HIS	4.3
1	A	456	SER	4.1
1	A	397	LEU	4.1
1	A	453	TRP	4.0
1	A	395	GLY	3.7
1	A	393	ARG	3.7
1	A	458	ASP	3.6
1	A	396	THR	3.6
1	A	394	ASN	3.4
1	A	501	SER	3.3
1	A	425	LEU	3.3
1	A	542	SER	3.1
1	A	369	THR	3.1
1	A	423	PRO	3.1
1	A	543	VAL	3.0
1	A	457	ARG	2.9
1	A	365	PRO	2.9
1	A	422	HIS	2.8
1	A	428	ALA	2.8
1	A	473	ASP	2.7
1	A	426	PRO	2.6
1	A	391	LYS	2.5
1	A	427	ARG	2.4
1	A	533	PRO	2.4
1	A	363	LEU	2.2
1	A	520	ASP	2.2
1	A	455	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	368	GLY	2.1
1	A	487	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.