



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:22 PM UTC

PDB ID : 8FPQ / pdb_00008fpq
Title : PCSK9 in complex with an inhibitor
Authors : Xu, M.; Chopra, R.
Deposited on : 2023-01-05
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	92	95% 5%
2	B	530	8% 71% 17% • 11%
3	L	15	40% 53% 7%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4427 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	92	Total	C	N	O	S	0	1	0
			748	479	136	131	2			

- Molecule 2 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	472	Total	C	N	O	S	0	0	0
			3508	2163	646	668	31			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	474	ILE	VAL	conflict	UNP Q8NBP7

- Molecule 3 is a protein called MCR-PHE-VAL-PRO-THR-THR-BIF-MAA-BIF-MAA-GLU-ALA-PRO-ALA-NEH inhibitor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	15	Total	C	N	O	S	0	0	0
			118	85	14	18	1			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	13	Total	O	0	0
			13	13		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	38	Total 38	O 38	0	0
5	L	1	Total 1	O 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proprotein convertase subtilisin/kexin type 9



- Molecule 2: Proprotein convertase subtilisin/kexin type 9



- Molecule 3: MCR-PHE-VAL-PRO-THR-THR-BIF-MAA-BIF-MAA-GLU-ALA-PRO-ALA-NEH inhibitor



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.41Å 70.37Å 150.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.67 – 2.50 57.67 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (57.67-2.50) 99.8 (57.67-2.50)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1	Depositor
R, R_{free}	0.188 , 0.234 0.194 , 0.233	Depositor DCC
R_{free} test set	1215 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	37.0	Xtriage
Anisotropy	0.258	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4427	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MCR, NEH, BIF, MAA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/768	0.95	0/1037
2	B	0.71	0/3566	1.11	16/4836 (0.3%)
3	L	1.20	0/66	1.54	0/90
All	All	0.71	0/4400	1.09	16/5963 (0.3%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	336	VAL	N-CA-C	7.58	118.81	107.75
2	B	519	GLY	N-CA-C	7.25	119.53	111.85
2	B	539	ALA	CA-C-N	-6.64	113.54	120.38
2	B	539	ALA	C-N-CA	-6.64	113.54	120.38
2	B	620	GLU	N-CA-C	6.33	119.86	111.75
2	B	426	GLU	CA-CB-CG	-6.09	101.91	114.10
2	B	330	ALA	N-CA-C	5.69	119.54	109.58
2	B	221	SER	N-CA-C	-5.65	105.88	112.89
2	B	467	PRO	N-CA-C	5.54	121.51	114.92
2	B	436	LEU	N-CA-C	5.49	118.02	111.71
2	B	154	ILE	N-CA-C	-5.48	101.48	107.73
2	B	474	ILE	N-CA-C	5.46	115.72	107.75
2	B	163	PRO	CA-C-N	5.40	125.07	119.56
2	B	163	PRO	C-N-CA	5.40	125.07	119.56
2	B	431	GLU	N-CA-C	5.33	117.50	111.11
2	B	313	THR	N-CA-C	5.21	117.59	109.52

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	748	0	763	5	0
2	B	3508	0	3425	54	0
3	L	118	0	106	7	0
4	B	1	0	0	0	0
5	A	13	0	0	0	0
5	B	38	0	0	2	0
5	L	1	0	0	0	0
All	All	4427	0	4294	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:641:THR:HG1	2:B:671:ALA:N	1.62	0.96
2:B:515:PHE:C	2:B:517:GLY:HA3	2.04	0.82
2:B:340:ASN:HD21	2:B:344:GLN:HE21	1.36	0.73
2:B:638:LEU:HB2	2:B:673:THR:HG23	1.73	0.70
2:B:199:ARG:NH1	2:B:243:LYS:O	2.24	0.70
2:B:584:GLN:HB2	2:B:587:GLN:HG2	1.74	0.70
2:B:490:SER:HB2	2:B:520:VAL:HG12	1.78	0.65
1:A:118:LEU:O	2:B:303:ARG:NE	2.30	0.64
2:B:332:GLU:O	2:B:408:LEU:HD21	1.99	0.62
1:A:104:ARG:HH21	1:A:105[B]:ARG:NH2	1.98	0.62
2:B:495:ARG:HH11	2:B:644:VAL:HG23	1.63	0.62
2:B:317:ASN:HB3	3:L:9:BIF:H11	1.83	0.59
2:B:455:LEU:HD13	2:B:529:LEU:HD13	1.84	0.58
2:B:529:LEU:HG	2:B:532:ALA:HB2	1.85	0.58
2:B:352:GLY:C	3:L:9:BIF:H10	2.31	0.56
2:B:330:ALA:O	2:B:333:VAL:HG22	2.05	0.56
2:B:494:LYS:HD3	2:B:517:GLY:N	2.22	0.55
3:L:14:ALA:O	3:L:15:NEH:HB3	2.07	0.55
2:B:566:TRP:HD1	2:B:568:VAL:O	1.90	0.54
2:B:558:VAL:HG21	2:B:606:LEU:HD23	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:515:PHE:C	2:B:517:GLY:CA	2.79	0.53
2:B:641:THR:OG1	2:B:671:ALA:N	2.34	0.53
2:B:610:VAL:HG22	2:B:677:ILE:HG12	1.90	0.53
2:B:612:GLU:HG2	2:B:675:VAL:HG22	1.89	0.52
2:B:196:ILE:O	2:B:199:ARG:HG2	2.09	0.52
2:B:584:GLN:HB2	2:B:587:GLN:CG	2.39	0.52
2:B:529:LEU:O	2:B:532:ALA:HB3	2.10	0.52
2:B:468:THR:O	2:B:513:ASN:ND2	2.43	0.52
2:B:541:PRO:HA	2:B:596:ILE:HD12	1.92	0.51
2:B:588:CYS:CB	2:B:596:ILE:HD13	2.41	0.51
2:B:468:THR:HG22	2:B:470:MET:H	1.76	0.51
2:B:317:ASN:HB3	3:L:9:BIF:C11	2.41	0.50
2:B:340:ASN:ND2	2:B:344:GLN:HE21	2.07	0.49
2:B:584:GLN:N	2:B:585:PRO:HD3	2.27	0.49
2:B:476:ARG:HG2	2:B:508:VAL:HG22	1.95	0.49
2:B:340:ASN:HD21	2:B:344:GLN:HG2	1.78	0.49
2:B:185:LEU:HD11	2:B:271:ILE:HD11	1.95	0.48
2:B:186:ASP:OD2	2:B:230:LEU:HD12	2.13	0.48
2:B:465:SER:HB3	2:B:473:ALA:HB2	1.96	0.47
2:B:340:ASN:HD21	2:B:344:GLN:NE2	2.09	0.47
2:B:588:CYS:HB2	2:B:596:ILE:HD13	1.95	0.47
2:B:340:ASN:ND2	2:B:344:GLN:HG2	2.30	0.47
2:B:313:THR:HG21	2:B:328:ALA:HA	1.96	0.47
2:B:476:ARG:HD3	2:B:506:LYS:HZ1	1.80	0.46
2:B:591:HIS:CE1	2:B:593:GLU:HB3	2.51	0.46
3:L:4:PRO:HA	3:L:9:BIF:O	2.16	0.45
3:L:7:BIF:HA	3:L:8:MAA:HA	1.75	0.44
2:B:591:HIS:CE1	2:B:594:ALA:HB2	2.52	0.44
2:B:258:LYS:HE3	5:B:818:HOH:O	2.17	0.43
2:B:622:VAL:HB	2:B:656:VAL:CG2	2.49	0.43
1:A:118:LEU:HD23	1:A:118:LEU:HA	1.72	0.43
5:B:806:HOH:O	3:L:6:THR:HG22	2.18	0.42
2:B:201:MET:CE	2:B:248:ARG:HH21	2.32	0.42
2:B:522:ALA:C	2:B:523:ILE:HG13	2.45	0.41
1:A:101:GLN:HG2	1:A:104:ARG:NH2	2.35	0.41
2:B:559:LEU:HD21	2:B:562:CYS:SG	2.61	0.41
1:A:101:GLN:HB3	1:A:133:LEU:HD21	2.03	0.40
2:B:338:ALA:HB1	2:B:365:GLY:HA3	2.04	0.40
2:B:178:SER:C	2:B:180:VAL:H	2.30	0.40
2:B:156:TRP:CH2	2:B:364:PRO:HB3	2.57	0.40
2:B:591:HIS:O	2:B:593:GLU:N	2.55	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:613:HIS:CE1	2:B:615:ILE:HD11	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	91/92 (99%)	87 (96%)	4 (4%)	0	100	100
2	B	452/530 (85%)	435 (96%)	14 (3%)	3 (1%)	18	34
3	L	9/15 (60%)	9 (100%)	0	0	100	100
All	All	552/637 (87%)	531 (96%)	18 (3%)	3 (0%)	24	43

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	534	CYS
2	B	427	ALA
2	B	280	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/79 (101%)	80 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	380/422 (90%)	369 (97%)	11 (3%)	37	65
3	L	7/7 (100%)	5 (71%)	2 (29%)	0	0
All	All	467/508 (92%)	454 (97%)	13 (3%)	38	66

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	153	SER
2	B	190	GLN
2	B	192	ASP
2	B	283	LEU
2	B	468	THR
2	B	508	VAL
2	B	549	ARG
2	B	550	VAL
2	B	567	GLU
2	B	570	ASP
2	B	673	THR
3	L	5	THR
3	L	6	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	190	GLN
2	B	344	GLN
2	B	417	HIS
2	B	513	ASN
2	B	591	HIS
2	B	602	HIS
2	B	621	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BIF	L	7	3	17,18,19	0.52	0	18,23,25	0.75	0
3	MAA	L	10	3	4,5,6	0.81	0	2,5,7	2.32	1 (50%)
3	MAA	L	8	3	4,5,6	0.67	0	2,5,7	0.92	0
3	BIF	L	9	3	17,18,19	0.52	0	18,23,25	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BIF	L	7	3	-	0/9/10/12	0/2/2/2
3	MAA	L	10	3	-	1/2/4/6	-
3	MAA	L	8	3	-	0/2/4/6	-
3	BIF	L	9	3	-	2/9/10/12	0/2/2/2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	10	MAA	CB-CA-N	3.02	117.23	110.20

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	L	10	MAA	CB-CA-N-CM
3	L	9	BIF	CA-CB-CG-CD2
3	L	9	BIF	CA-CB-CG-CD1

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	L	7	BIF	1	0
3	L	8	MAA	1	0
3	L	9	BIF	4	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	92/92 (100%)	-0.13	1 (1%) 78 75	21, 32, 48, 63	1 (1%)
2	B	472/530 (89%)	0.31	45 (9%) 14 12	18, 35, 75, 106	0
3	L	9/15 (60%)	-0.21	0 100 100	26, 28, 35, 47	0
All	All	573/637 (89%)	0.24	46 (8%) 18 16	18, 34, 73, 106	1 (0%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	515	PHE	6.2
2	B	548	THR	5.4
2	B	553	HIS	4.9
2	B	671	ALA	4.5
2	B	530	PRO	3.9
2	B	615	ILE	3.6
2	B	470	MET	3.4
2	B	533	ASN	3.4
2	B	568	VAL	3.4
2	B	602	HIS	3.4
2	B	585	PRO	3.3
2	B	556	GLY	3.3
2	B	220	ALA	3.2
2	B	532	ALA	3.2
2	B	604	PRO	3.2
2	B	639	PRO	3.2
2	B	641	THR	3.0
2	B	591	HIS	3.0
2	B	468	THR	2.9
2	B	517	GLY	2.9
2	B	570	ASP	2.6
2	B	640	GLY	2.6
2	B	494	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	505	GLY	2.6
2	B	534	CYS	2.6
2	B	642	SER	2.5
2	B	212	ASP	2.5
2	B	447	SER	2.4
2	B	605	GLY	2.4
2	B	192	ASP	2.4
2	B	531	GLN	2.4
2	B	514	ALA	2.3
2	B	469	ARG	2.3
2	B	643	HIS	2.3
2	B	153	SER	2.3
2	B	492	SER	2.2
2	B	244	GLY	2.1
2	B	567	GLU	2.1
2	B	331	PRO	2.1
2	B	506	LYS	2.1
2	B	584	GLN	2.1
2	B	594	ALA	2.1
1	A	61	THR	2.1
2	B	221	SER	2.1
2	B	619	GLN	2.0
2	B	603	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAA	L	8	6/7	0.90	0.11	24,25,27,30	0
3	BIF	L	9	17/18	0.93	0.08	19,23,28,30	0
3	MAA	L	10	6/7	0.94	0.08	28,30,32,33	0
3	BIF	L	7	17/18	0.96	0.07	18,23,27,27	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CA	B	701	1/1	0.93	0.10	59,59,59,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.