



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 12:31 AM UTC

PDB ID : 2FQZ / pdb_00002fqz
Title : Metal-depleted Ecl18kI in complex with uncleaved DNA
Authors : Bochtler, M.; Szczepanowski, R.H.; Tamulaitis, G.; Grazulis, S.; Czapinska, H.; Manakova, E.; Siksny, V.
Deposited on : 2006-01-18
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	9	 33% 67%
1	G	9	 22% 78%
2	F	9	 11% 89%
2	H	9	 22% 78%
3	A	305	 31% 70% 23% • 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	B	305	12% 70% 24%
3	C	305	7% 71% 22%
3	D	305	3% 72% 21%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11645 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA STRAND 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	9	Total	C	N	O	P	0	9	0
			364	172	74	102	16			
1	G	9	Total	C	N	O	P	0	9	0
			364	172	74	102	16			

- Molecule 2 is a DNA chain called DNA STRAND 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	9	Total	C	N	O	P	0	9	0
			362	172	68	106	16			
2	H	9	Total	C	N	O	P	0	9	0
			362	172	68	106	16			

- Molecule 3 is a protein called R.Ecl18kI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	290	Total	C	N	O	S	21	0	0
			2393	1522	396	462	13			
3	B	293	Total	C	N	O	S	15	0	0
			2421	1541	399	468	13			
3	C	292	Total	C	N	O	S	19	0	0
			2412	1537	398	464	13			
3	D	292	Total	C	N	O	S	11	0	0
			2412	1537	398	464	13			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	277	GLN	ARG	engineered mutation	UNP O87963
B	277	GLN	ARG	engineered mutation	UNP O87963
C	277	GLN	ARG	engineered mutation	UNP O87963
D	277	GLN	ARG	engineered mutation	UNP O87963

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	16	Total 16	O 16	0	0
4	F	15	Total 15	O 15	0	0
4	G	24	Total 24	O 24	0	0
4	H	20	Total 20	O 20	0	0
4	A	83	Total 83	O 83	0	0
4	B	133	Total 133	O 133	0	0
4	C	135	Total 135	O 135	0	0
4	D	129	Total 129	O 129	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA STRAND 1



• Molecule 1: DNA STRAND 1




• Molecule 2: DNA STRAND 2



• Molecule 2: DNA STRAND 2




• Molecule 3: R.Ecl18kI








• Molecule 3: R.Ecl18kI



• Molecule 3: R.Ecl18kI



• Molecule 3: R.Ecl18kI



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	77.37Å 95.52Å 190.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 20.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	92.0 (20.00-2.00) 91.9 (20.00-2.00)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.99 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.210 , 0.260 0.210 , 0.253	Depositor DCC
R_{free} test set	4666 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtriage
Anisotropy	0.317	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	11645	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	E	0.23	0/408	0.60	0/626
1	G	0.26	0/408	0.62	0/626
2	F	0.25	0/404	0.61	0/620
2	H	0.26	0/404	0.61	0/620
3	A	0.60	0/2440	0.93	0/3293
3	B	0.68	0/2469	0.94	0/3331
3	C	0.71	0/2460	0.96	0/3320
3	D	0.69	0/2460	0.94	0/3320
All	All	0.63	0/11453	0.90	0/15756

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	B	241	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	364	0	193	13	0
1	G	364	0	195	10	0
2	F	362	0	194	15	0
2	H	362	0	196	10	0
3	A	2393	0	2360	60	0
3	B	2421	0	2384	64	0
3	C	2412	0	2380	59	0
3	D	2412	0	2380	61	0
4	A	83	0	0	10	0
4	B	133	0	0	9	0
4	C	135	0	0	9	0
4	D	129	0	0	11	0
4	E	16	0	0	1	0
4	F	15	0	0	1	0
4	G	24	0	0	1	0
4	H	20	0	0	2	0
All	All	11645	0	10282	257	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 257 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:16:HOH:O	3:D:157:LYS:HG3	1.35	1.26
1:G:-3[A]:DG:H5"	3:C:157:LYS:HB2	1.21	1.20
3:D:160:ASP:HB3	4:D:398:HOH:O	1.52	1.09
2:H:-3[B]:DC:H5"	3:C:157:LYS:HB2	1.41	1.02
3:B:155:ILE:HD11	3:B:169:TYR:CE2	1.97	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	286/305 (94%)	275 (96%)	11 (4%)	0	100	100
3	B	289/305 (95%)	278 (96%)	10 (4%)	1 (0%)	36	35
3	C	288/305 (94%)	274 (95%)	13 (4%)	1 (0%)	36	35
3	D	288/305 (94%)	283 (98%)	4 (1%)	1 (0%)	36	35
All	All	1151/1220 (94%)	1110 (96%)	38 (3%)	3 (0%)	36	35

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	172	ASN
3	B	85	ILE
3	C	101	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	269/283 (95%)	253 (94%)	16 (6%)	18	14
3	B	272/283 (96%)	252 (93%)	20 (7%)	13	9
3	C	271/283 (96%)	260 (96%)	11 (4%)	27	26
3	D	271/283 (96%)	256 (94%)	15 (6%)	19	17
All	All	1083/1132 (96%)	1021 (94%)	62 (6%)	18	15

5 of 62 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	B	266	SER
3	D	174	ARG
3	B	301	SER
3	D	158	LEU
3	D	252	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	20	HIS
3	D	47	ASN
3	D	277	GLN
3	D	225	ASN
3	B	142	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	9/9 (100%)	0.10	0 100 100	16, 18, 21, 25	9 (100%)
1	G	9/9 (100%)	-0.05	0 100 100	13, 14, 18, 21	9 (100%)
2	F	9/9 (100%)	0.23	0 100 100	14, 16, 23, 24	9 (100%)
2	H	9/9 (100%)	-0.16	0 100 100	14, 15, 18, 20	9 (100%)
3	A	290/305 (95%)	1.57	95 (32%) 1 1	24, 43, 65, 69	5 (1%)
3	B	293/305 (96%)	0.84	37 (12%) 8 7	15, 34, 53, 65	3 (1%)
3	C	292/305 (95%)	0.62	21 (7%) 21 20	22, 33, 53, 66	5 (1%)
3	D	292/305 (95%)	0.55	10 (3%) 48 47	21, 34, 51, 58	3 (1%)
All	All	1203/1256 (95%)	0.87	163 (13%) 7 6	13, 35, 57, 69	52 (4%)

The worst 5 of 163 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	B	145	ILE	6.0
3	A	135	ALA	5.0
3	C	78	TYR	4.8
3	A	4	LEU	4.7
3	A	67	PHE	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.