



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:42 AM UTC

PDB ID : 1FRB / pdb_00001frb
Title : FR-1 PROTEIN/NADPH/ZOPOLRESTAT COMPLEX
Authors : Wilson, D.K.; Quiocho, F.A.
Deposited on : 1995-08-08
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	315	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3060 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FR-1 PROTEIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	0	0	0
			2535	1627	424	474	10			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is 3,4-DIHYDRO-4-OXO-3-((5-TRIFLUOROMETHYL-2-BENZOTHAZOLYL)METHYL)-1-PHTHALAZINE ACETIC ACID (CCD ID: ZST) (formula: $C_{19}H_{12}F_3N_3O_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	S	
			29	19	3	3	3	1	
									0
									0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	448	Total	O		
			448	448	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: FR-1 PROTEIN

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.45Å 65.38Å 90.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.70	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-1.70)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.158 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3060	wwPDB-VP
Average B, all atoms (Å ²)	9.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, ZST

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	10/2595 (0.4%)	1.18	2/3518 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	25

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	177	HIS	CD2-NE2	-6.11	1.31	1.37
1	A	163	HIS	CD2-NE2	-5.73	1.31	1.37
1	A	240	HIS	CD2-NE2	-5.66	1.31	1.37
1	A	110	HIS	CD2-NE2	-5.53	1.31	1.37
1	A	252	HIS	CD2-NE2	-5.53	1.31	1.37
1	A	200	HIS	CD2-NE2	-5.45	1.31	1.37
1	A	41	HIS	CD2-NE2	-5.43	1.31	1.37
1	A	187	HIS	CD2-NE2	-5.34	1.31	1.37
1	A	163	HIS	CG-ND1	-5.22	1.32	1.38
1	A	240	HIS	CG-ND1	-5.07	1.32	1.38

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	109	ILE	N-CA-C	-5.61	101.44	108.84
1	A	136	THR	N-CA-C	5.40	117.49	110.53

There are no chirality outliers.

All (25) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	118	GLY	Mainchain
1	A	120	GLU	Mainchain
1	A	122	PHE	Mainchain
1	A	123	PRO	Mainchain
1	A	129	ARG	Mainchain
1	A	135	THR	Mainchain
1	A	173	PRO	Mainchain
1	A	178	LYS	Mainchain
1	A	182	ASN	Mainchain
1	A	209	TYR	Sidechain
1	A	219	SER	Mainchain
1	A	220	ALA	Mainchain
1	A	221	LYS	Mainchain
1	A	223	GLU	Mainchain
1	A	24	PRO	Mainchain
1	A	25	ASN	Mainchain
1	A	277	ASP	Mainchain
1	A	279	GLN	Mainchain
1	A	300	LEU	Mainchain
1	A	303	THR	Mainchain
1	A	306	MET	Mainchain
1	A	4	VAL	Mainchain
1	A	45	ALA	Mainchain
1	A	68	GLN	Mainchain
1	A	84	GLU	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2535	0	2547	30	0
2	A	48	0	26	4	0
3	A	29	0	11	1	0
4	A	448	0	0	4	0
All	All	3060	0	2584	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:134:LYS:H	1:A:134:LYS:HE2	1.59	0.67
1:A:187:HIS:HD2	1:A:189:TYR:H	1.46	0.62
1:A:239:LYS:HE3	1:A:284:GLU:OE2	2.00	0.61
1:A:79:TRP:CD2	1:A:80:PRO:HD2	2.35	0.61
1:A:23:PRO:HG2	1:A:26:GLN:NE2	2.17	0.60
2:A:350:NDP:H42N	3:A:351:ZST:H171	1.88	0.55
1:A:239:LYS:HE2	4:A:824:HOH:O	2.10	0.52
1:A:8:THR:O	1:A:9:LYS:HB2	2.09	0.52
1:A:135:THR:CG2	4:A:782:HOH:O	2.57	0.52
1:A:209:TYR:CZ	2:A:350:NDP:H41N	2.45	0.52
1:A:85:LYS:HG2	4:A:771:HOH:O	2.10	0.51
1:A:126:ASP:OD2	1:A:127:GLN:NE2	2.45	0.50
1:A:79:TRP:CG	1:A:80:PRO:HD2	2.47	0.49
1:A:183:GLN:OE1	2:A:350:NDP:H2N	2.13	0.48
1:A:23:PRO:HG2	1:A:26:GLN:HE21	1.79	0.46
1:A:209:TYR:CE1	2:A:350:NDP:H41N	2.50	0.46
1:A:84:GLU:HG3	4:A:619:HOH:O	2.15	0.45
1:A:301:PRO:HD2	1:A:302:GLU:OE2	2.16	0.45
1:A:80:PRO:HG3	1:A:111:TRP:HB2	1.99	0.45
1:A:20:TRP:CE3	1:A:21:LYS:HE3	2.53	0.43
1:A:185:GLU:OE2	1:A:187:HIS:HE1	2.01	0.43
1:A:20:TRP:O	1:A:21:LYS:HB2	2.19	0.43
1:A:178:LYS:HE3	1:A:178:LYS:HB3	1.58	0.42
1:A:129:ARG:HH11	1:A:129:ARG:HD3	1.73	0.42
1:A:262:LYS:C	1:A:262:LYS:HD3	2.44	0.41
1:A:185:GLU:HB2	1:A:209:TYR:CE1	2.55	0.41
1:A:187:HIS:HB2	1:A:188:PRO:HD2	2.02	0.41
1:A:217:ARG:HG2	1:A:220:ALA:HB2	2.02	0.41
1:A:221:LYS:H	1:A:221:LYS:HG2	1.52	0.41
1:A:213:GLY:O	1:A:214:SER:C	2.63	0.41
1:A:221:LYS:HB2	1:A:223:GLU:OE2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/315 (99%)	304 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/281 (100%)	272 (97%)	9 (3%)	34	17

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	GLU
1	A	95	THR
1	A	119	LYS
1	A	134	LYS
1	A	135	THR
1	A	221	LYS
1	A	229	GLU
1	A	274	GLN
1	A	302	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	116	GLN
1	A	127	GLN
1	A	150	GLN
1	A	187	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	A	350	-	51,52,52	1.69	6 (11%)	71,80,80	1.10	5 (7%)
3	ZST	A	351	-	32,32,32	0.69	0	43,48,48	1.11	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	350	-	-	6/34/77/77	0/5/5/5
3	ZST	A	351	-	-	0/14/14/14	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	350	NDP	C4N-C3N	-6.87	1.37	1.50
2	A	350	NDP	C4N-C5N	-5.05	1.35	1.49
2	A	350	NDP	PN-O3	4.04	1.63	1.59
2	A	350	NDP	PA-O3	3.64	1.63	1.59
2	A	350	NDP	C2N-C3N	3.32	1.44	1.35
2	A	350	NDP	P2B-O2B	2.28	1.63	1.59

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	351	ZST	C10-C9-N1	-4.61	108.47	112.10
2	A	350	NDP	O2A-PA-O3	3.73	117.34	107.27
2	A	350	NDP	C3N-C2N-N1N	-3.67	117.81	123.20
2	A	350	NDP	C1D-N1N-C2N	-2.99	116.21	121.14
2	A	350	NDP	O7N-C7N-N7N	-2.40	117.51	122.89
2	A	350	NDP	P2B-O2B-C2B	-2.03	118.00	123.43
3	A	351	ZST	O3-C18-C17	2.01	120.75	114.51

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	350	NDP	O4D-C1D-N1N-C2N
2	A	350	NDP	PA-O3-PN-O5D
2	A	350	NDP	C4D-C5D-O5D-PN
2	A	350	NDP	PN-O3-PA-O1A
2	A	350	NDP	C2B-O2B-P2B-O2X
2	A	350	NDP	PA-O3-PN-O2N


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	350	NDP	4	0
3	A	351	ZST	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.