



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:32 PM UTC

PDB ID : 3FRO / pdb_00003fro
Title : Crystal structure of Pyrococcus abyssi glycogen synthase with open and closed conformations
Authors : Diaz, A.; Guinovart, J.J.; Fita, I.; Ferrer, J.C.
Deposited on : 2009-01-08
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	439	4% 84% 15% .
1	B	439	7% 86% 13%
1	C	439	2% 90% 10% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10635 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GlgA glycogen synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	439	Total	C	N	O	S	0	2	0
			3502	2268	589	634	11			
1	B	439	Total	C	N	O	S	0	1	0
			3494	2263	588	633	10			
1	C	439	Total	C	N	O	S	0	1	0
			3494	2263	588	633	10			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ARG	-	expression tag	UNP Q9V2J8
A	0	HIS	-	expression tag	UNP Q9V2J8
B	-1	ARG	-	expression tag	UNP Q9V2J8
B	0	HIS	-	expression tag	UNP Q9V2J8
C	-1	ARG	-	expression tag	UNP Q9V2J8
C	0	HIS	-	expression tag	UNP Q9V2J8

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			8	4	1	3		
2	A	1	Total	C	N	O	0	0
			8	4	1	3		
2	B	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O P	0	0
			5	4 1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is 1,5-anhydro-D-fructose (CCD ID: NHF) (formula: C₆H₁₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			11	6	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	40	Total	O	0	0
			40	40		
5	B	30	Total	O	0	0
			30	30		
5	C	25	Total	O	0	0
			25	25		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GlgA glycogen synthase




- Molecule 1: GlgA glycogen synthase



- Molecule 1: GlgA glycogen synthase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.00Å 139.91Å 159.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.91 – 2.50 24.91 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.4 (24.91-2.50) 99.3 (24.91-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.214 , 0.255 0.218 , 0.256	Depositor DCC
R_{free} test set	2950 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	41.8	Xtriage
Anisotropy	0.658	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 54.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	10635	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NHF, PO4, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/3584	0.67	0/4832
1	B	0.41	0/3576	0.67	1/4822 (0.0%)
1	C	0.41	0/3576	0.66	0/4822
All	All	0.41	0/10736	0.67	1/14476 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	118	GLU	N-CA-C	5.05	114.11	108.25

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3502	0	3504	45	0
1	B	3494	0	3496	38	0
1	C	3494	0	3496	24	0
2	A	16	0	24	0	0
2	B	8	0	12	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	5	0	0	0	0
4	B	11	0	10	0	0
5	A	40	0	0	0	0
5	B	30	0	0	0	0
5	C	25	0	0	0	0
All	All	10635	0	10542	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253:MET:HE3	1:A:289:ILE:HD12	1.56	0.86
1:C:281:PHE:HA	1:C:284:MET:HE2	1.62	0.81
1:A:250:VAL:HG21	1:A:391:LEU:HD22	1.63	0.79
1:A:150:ILE:HG21	1:A:153:LEU:HD13	1.73	0.70
1:C:150:ILE:HG21	1:C:153:LEU:CD1	2.22	0.69
1:B:269:LEU:HD22	1:B:303:LEU:HD12	1.75	0.69
1:A:433:PHE:HB2	1:A:436:ILE:HD13	1.77	0.67
1:B:287:ILE:HG22	1:B:289:ILE:HD11	1.76	0.67
1:A:323:ARG:HG2	1:A:351:LEU:HD11	1.79	0.64
1:A:29:LEU:HD12	1:A:418:ALA:HB1	1.80	0.63
1:C:346:LEU:HD13	1:C:366:ILE:HD13	1.79	0.63
1:A:275:LEU:HD13	1:A:284:MET:HE1	1.81	0.62
1:C:51:ILE:HD13	1:C:65:VAL:HG21	1.82	0.61
1:A:253:MET:HE3	1:A:289:ILE:CD1	2.30	0.61
1:B:253:MET:HE2	1:B:289:ILE:HD13	1.83	0.60
1:A:253:MET:HE2	1:A:255:ILE:HB	1.84	0.59
1:A:393:LEU:C	1:A:395:ARG:H	2.10	0.59
1:B:407:LYS:HA	1:B:410:MET:HE3	1.85	0.59
1:B:246:MET:HE2	1:B:285:ARG:HD2	1.85	0.59
1:C:289:ILE:HD13	1:C:317:LEU:HD22	1.85	0.59
1:B:56:VAL:HG12	1:B:57:PHE:CD1	2.38	0.58
1:C:129:TRP:HA	1:C:132:VAL:HG23	1.83	0.58
1:C:312:VAL:C	1:C:313:ILE:HD12	2.28	0.58
1:A:29:LEU:CD1	1:A:418:ALA:HB1	2.33	0.58
1:C:150:ILE:HG21	1:C:153:LEU:HD12	1.86	0.58
1:A:433:PHE:CB	1:A:436:ILE:HD13	2.34	0.57
1:A:51:ILE:HD13	1:A:65:VAL:HG21	1.86	0.57
1:B:284:MET:HE1	1:B:387:ILE:HG21	1.86	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:37:LEU:HD21	1:C:74:ARG:NH1	2.19	0.56
1:B:129:TRP:HA	1:B:132:VAL:HG23	1.88	0.56
1:B:287:ILE:HG22	1:B:289:ILE:CD1	2.36	0.56
1:B:250:VAL:HG11	1:B:391:LEU:HD13	1.87	0.56
1:B:250:VAL:HG11	1:B:391:LEU:CD1	2.36	0.56
1:B:253:MET:CE	1:B:255:ILE:HD11	2.35	0.56
1:C:56:VAL:HG12	1:C:57:PHE:CD1	2.42	0.54
1:A:275:LEU:HD13	1:A:284:MET:CE	2.38	0.54
1:A:332:VAL:HG23	1:A:353:ALA:HB1	1.90	0.53
1:C:240:LEU:HD22	1:C:328:SER:HB2	1.90	0.53
1:B:13:VAL:HG22	1:B:45:ARG:HG3	1.90	0.53
1:A:326:TYR:HB2	1:A:351:LEU:HD13	1.89	0.53
1:A:333:ILE:HG22	1:A:335:PRO:HD3	1.91	0.52
1:C:219:ILE:HD12	1:C:346:LEU:HB3	1.90	0.52
1:A:371:THR:HG21	1:A:408:ARG:HD3	1.91	0.51
1:B:209:GLU:HG3	1:B:434:ASP:HB2	1.91	0.51
1:C:54:ILE:HG23	1:C:109:LEU:HD22	1.92	0.51
1:C:240:LEU:HD21	1:C:325:LEU:HD23	1.93	0.50
1:B:318:SER:O	1:B:322:VAL:HG23	2.11	0.50
1:B:176:ASP:C	1:B:177:ILE:HD12	2.37	0.50
1:B:82:LEU:HD12	1:B:98:ARG:HG3	1.93	0.49
1:C:140:LYS:NZ	1:C:170:GLU:OE2	2.43	0.49
1:C:250:VAL:HG21	1:C:391:LEU:HD12	1.95	0.49
1:B:252:PHE:CE1	1:B:284:MET:HE3	2.48	0.49
1:C:29:LEU:CD1	1:C:418:ALA:HB1	2.43	0.49
1:A:115:LEU:HD21	1:A:120:LEU:HA	1.96	0.48
1:C:394:SER:HA	1:C:398:LEU:HD11	1.96	0.48
1:A:275:LEU:CD2	1:A:388:LEU:HD13	2.44	0.48
1:C:120:LEU:HD12	1:C:121:PRO:HD2	1.96	0.47
1:C:51:ILE:HD13	1:C:65:VAL:CG2	2.44	0.47
1:A:54:ILE:HG23	1:A:109:LEU:HD22	1.96	0.46
1:B:357:ALA:HB1	1:B:363:LEU:HD13	1.96	0.46
1:C:289:ILE:HD13	1:C:317:LEU:CD2	2.46	0.46
1:A:348:ALA:HB1	1:A:353:ALA:HB3	1.98	0.45
1:B:54:ILE:HG23	1:B:109:LEU:HD22	1.98	0.45
1:B:120:LEU:HD12	1:B:121:PRO:HD2	1.97	0.45
1:A:357:ALA:HB3	1:A:367:ILE:HD12	1.97	0.45
1:B:322:VAL:HG12	1:B:326:TYR:CE1	2.52	0.45
1:B:287:ILE:CG2	1:B:289:ILE:HD11	2.45	0.44
1:A:150:ILE:CG2	1:A:153:LEU:HD13	2.46	0.44
1:A:253:MET:CE	1:A:289:ILE:HD12	2.38	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:TRP:HA	1:A:132:VAL:HG23	2.00	0.44
1:A:395:ARG:O	1:A:396:SER:CB	2.66	0.44
1:B:326:TYR:HB2	1:B:351:LEU:HD13	2.00	0.44
1:A:250:VAL:O	1:A:250:VAL:HG23	2.19	0.43
1:A:89:TYR:HA	1:A:96:LEU:HD13	2.00	0.43
1:A:255:ILE:HG21	1:A:326:TYR:OH	2.19	0.43
1:A:257:ARG:HG2	1:A:259:ASP:OD1	2.18	0.43
1:C:1:MET:HE1	1:C:425:TYR:HB3	2.01	0.43
1:A:50:GLU:HB2	1:A:62:GLN:HE21	1.84	0.42
1:B:115:LEU:HD21	1:B:120:LEU:HA	2.00	0.42
1:A:212:ILE:HG23	1:A:432:ALA:HB3	2.00	0.42
1:B:291:LYS:CB	1:B:314:THR:HG23	2.50	0.42
1:C:63:VAL:HG21	1:C:109:LEU:HD23	2.00	0.42
1:C:269:LEU:HD22	1:C:303:LEU:HD12	2.01	0.42
1:B:333:ILE:HG22	1:B:335:PRO:HD3	2.02	0.42
1:A:176:ASP:C	1:A:177:ILE:HD12	2.44	0.42
1:A:393:LEU:C	1:A:395:ARG:N	2.77	0.42
1:B:291:LYS:HB3	1:B:314:THR:HG23	2.01	0.42
1:A:284:MET:HE3	1:A:286:PHE:HZ	1.85	0.42
1:B:246:MET:HE3	1:B:308:GLY:O	2.20	0.41
1:A:123:VAL:HG11	1:A:425:TYR:CG	2.55	0.41
1:B:240:LEU:HD21	1:B:325:LEU:HD23	2.01	0.41
1:B:253:MET:CE	1:B:325:LEU:HD13	2.50	0.41
1:B:398:LEU:HD23	1:B:398:LEU:O	2.20	0.41
1:A:46:PHE:CE1	1:A:77:ARG:HD3	2.55	0.41
1:A:371:THR:HG21	1:A:408:ARG:CD	2.51	0.41
1:B:82:LEU:CD1	1:B:98:ARG:HG3	2.50	0.41
1:B:41:PRO:HB3	1:B:83:LEU:HD22	2.03	0.41
1:B:357:ALA:HB3	1:B:367:ILE:HD12	2.02	0.41
1:A:6:LEU:HD12	1:A:126:PHE:HB3	2.03	0.41
1:B:323:ARG:HG3	1:B:351:LEU:HD11	2.01	0.41
1:A:255:ILE:HG23	1:A:255:ILE:O	2.21	0.40
1:A:375:VAL:HG12	1:A:376:LYS:N	2.37	0.40
1:B:313:ILE:HD12	1:B:313:ILE:N	2.36	0.40
1:A:250:VAL:HG21	1:A:391:LEU:CD2	2.43	0.40
1:A:57:PHE:CZ	1:A:98:ARG:HG2	2.56	0.40
1:A:418:ALA:O	1:A:422:VAL:HG23	2.20	0.40
1:B:423:LYS:HB3	1:B:428:SER:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	439/439 (100%)	422 (96%)	15 (3%)	2 (0%)	24	43
1	B	438/439 (100%)	419 (96%)	18 (4%)	1 (0%)	43	63
1	C	438/439 (100%)	425 (97%)	13 (3%)	0	100	100
All	All	1315/1317 (100%)	1266 (96%)	46 (4%)	3 (0%)	43	63

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	394	SER
1	A	396	SER
1	B	128	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/367 (100%)	354 (96%)	15 (4%)	27	53
1	B	368/367 (100%)	360 (98%)	8 (2%)	45	73
1	C	368/367 (100%)	357 (97%)	11 (3%)	36	64
All	All	1105/1101 (100%)	1071 (97%)	34 (3%)	35	62

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	LEU
1	A	29	LEU
1	A	153	LEU
1	A	209	GLU
1	A	247	ASP
1	A	248	GLU
1	A	260	ARG
1	A	280	GLU
1	A	282	GLN
1	A	296	LEU
1	A	369	ASN
1	A	388	LEU
1	A	395	ARG
1	A	398	LEU
1	A	431	ARG
1	B	11	LEU
1	B	29	LEU
1	B	37	LEU
1	B	49	GLU
1	B	260	ARG
1	B	398	LEU
1	B	423	LYS
1	B	431	ARG
1	C	-1	ARG
1	C	11	LEU
1	C	29	LEU
1	C	153	LEU
1	C	189	ILE
1	C	255	ILE
1	C	270	LYS
1	C	317	LEU
1	C	346	LEU
1	C	395	ARG
1	C	431	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	181	HIS
1	A	207	ASN
1	A	217	ASN
1	A	307	HIS
1	A	309	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	130	HIS
1	B	207	ASN
1	B	217	ASN
1	B	369	ASN
1	C	217	ASN
1	C	225	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	C	8002	-	4,4,4	0.93	0	6,6,6	0.48	0
2	TRS	B	7362	-	7,7,7	0.31	0	9,9,9	0.46	0
4	NHF	B	438	-	11,11,11	5.47	2 (18%)	11,15,15	2.39	1 (9%)
3	PO4	A	8000	-	4,4,4	0.93	0	6,6,6	0.49	0
3	PO4	B	8001	-	4,4,4	0.94	0	6,6,6	0.46	0
2	TRS	A	7360	-	7,7,7	0.32	0	9,9,9	0.31	0
2	TRS	A	7361	-	7,7,7	0.30	0	9,9,9	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	B	7362	-	-	6/9/9/9	-
4	NHF	B	438	-	-	1/2/19/19	0/1/1/1
2	TRS	A	7360	-	-	3/9/9/9	-
2	TRS	A	7361	-	-	6/9/9/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	438	NHF	C1-C2	-16.48	1.34	1.51
4	B	438	NHF	O2-C2	7.33	1.33	1.21

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	438	NHF	C1-O5-C5	7.40	122.06	111.67

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	7361	TRS	N-C-C1-O1
2	B	7362	TRS	C3-C-C1-O1
2	B	7362	TRS	N-C-C1-O1
2	B	7362	TRS	N-C-C2-O2
2	A	7360	TRS	C3-C-C2-O2
2	A	7360	TRS	N-C-C2-O2
2	A	7361	TRS	C1-C-C3-O3
2	A	7361	TRS	N-C-C3-O3
2	B	7362	TRS	C2-C-C1-O1
2	A	7360	TRS	C1-C-C2-O2
2	A	7361	TRS	C3-C-C1-O1
2	A	7361	TRS	C2-C-C3-O3
2	A	7361	TRS	C2-C-C1-O1
2	B	7362	TRS	C3-C-C2-O2
4	B	438	NHF	C4-C5-C6-O6
2	B	7362	TRS	C1-C-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/439 (100%)	0.27	19 (4%) 40 35	20, 42, 52, 58	2 (0%)
1	B	439/439 (100%)	0.49	30 (6%) 23 20	22, 43, 50, 56	1 (0%)
1	C	439/439 (100%)	0.22	8 (1%) 67 64	22, 42, 49, 56	1 (0%)
All	All	1317/1317 (100%)	0.33	57 (4%) 40 35	20, 42, 50, 58	4 (0%)

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	221	CYS	5.4
1	B	-1	ARG	5.0
1	A	394	SER	4.8
1	A	395	ARG	4.6
1	C	-1	ARG	4.6
1	A	262	GLN	4.1
1	B	218	GLY	4.0
1	B	413	SER	3.9
1	A	221	CYS	3.9
1	B	395	ARG	3.6
1	A	-1	ARG	3.5
1	B	341	PHE	3.4
1	C	164[A]	HIS	3.2
1	A	248	GLU	3.1
1	B	342	GLY	3.0
1	B	394	SER	3.0
1	A	396	SER	2.9
1	B	221	CYS	2.8
1	A	410	MET	2.8
1	B	361	GLY	2.7
1	B	364	ARG	2.7
1	B	362	GLY	2.7
1	A	255	ILE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	219	ILE	2.7
1	B	316	MET	2.7
1	A	258	PHE	2.7
1	B	0	HIS	2.7
1	A	0	HIS	2.6
1	A	282	GLN	2.5
1	C	394	SER	2.5
1	C	395	ARG	2.5
1	B	300	ALA	2.5
1	B	302	SER	2.5
1	A	350	CYS	2.4
1	B	412	PHE	2.4
1	B	346	LEU	2.4
1	C	93	TRP	2.3
1	B	305	GLU	2.3
1	B	396	SER	2.3
1	B	393	LEU	2.3
1	A	316[A]	MET	2.2
1	B	303	LEU	2.2
1	A	397	ASP	2.2
1	B	344	VAL	2.2
1	B	350	CYS	2.2
1	B	282	GLN	2.2
1	C	47	GLN	2.2
1	A	256	GLY	2.2
1	A	260	ARG	2.2
1	C	0	HIS	2.1
1	B	343	LEU	2.1
1	B	409	ALA	2.1
1	A	253	MET	2.1
1	B	222	SER	2.0
1	A	164[A]	HIS	2.0
1	B	373	ILE	2.0
1	B	242	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	TRS	B	7362	8/8	0.67	0.20	84,84,84,84	0
4	NHF	B	438	11/11	0.75	0.17	65,66,66,67	0
3	PO4	C	8002	5/5	0.80	0.29	85,85,85,86	0
2	TRS	A	7361	8/8	0.80	0.19	56,57,57,57	0
2	TRS	A	7360	8/8	0.84	0.19	60,61,61,61	0
3	PO4	B	8001	5/5	0.85	0.17	93,93,93,93	0
3	PO4	A	8000	5/5	0.86	0.27	81,81,81,82	0

6.5 Other polymers [i](#)

There are no such residues in this entry.