



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:22 PM UTC

PDB ID : 7FRR / pdb_00007frr
Title : PanDDA analysis group deposition – Crystal structure of PTP1B in complex with Z2856434906
Authors : Mehlman, T.; Biel, J.; Azeem, S.M.; Nelson, E.R.; Hossain, S.; Dunnett, L.E.; Paterson, N.G.; Douangamath, A.; Talon, R.; Axford, D.; Orins, H.; von Delft, F.; Keedy, D.A.
Deposited on : 2022-10-25
Resolution : 1.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtrriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	54% 79% 12% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	O1J	A	702	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5261 atoms, of which 2533 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	297	5054	1607	2512	439	480	16	0	14	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	SER	CYS	engineered mutation	UNP P18031
A	92	VAL	CYS	engineered mutation	UNP P18031


- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	20	4	12	1	3	0	0

- Molecule 3 is (benzyloxy)acetic acid (CCD ID: O1J) (formula: $C_9H_{10}O_3$) (labeled as "Ligand

of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			21	9	9	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	162	Total	O	0	5
			166	166		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.53Å 89.53Å 106.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	62.63 – 1.83 62.63 – 1.83	Depositor EDS
% Data completeness (in resolution range)	99.9 (62.63-1.83) 99.9 (62.63-1.83)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.31 (at 1.83Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.175 , 0.214 0.278 , 0.289	Depositor DCC
R_{free} test set	1727 reflections (3.93%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 60.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	5261	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: O1J, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.04	1/2598 (0.0%)	1.29	3/3501 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	206	PRO	C-O	-5.23	1.17	1.24

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	187	SER	O-C-N	-7.63	116.83	121.71
1	A	187	SER	CA-C-N	-7.10	112.86	119.82
1	A	187	SER	C-N-CA	-7.10	112.86	119.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2542	2512	2488	40	1
2	A	8	12	12	3	0
3	A	12	9	0	0	0
4	A	166	0	0	14	2

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2728	2533	2500	40	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272[A]:LEU:HD11	4:A:916:HOH:O	1.55	1.05
1:A:272[A]:LEU:HD21	4:A:916:HOH:O	1.75	0.86
1:A:233[A]:LEU:HA	4:A:899:HOH:O	1.77	0.85
1:A:233[B]:LEU:HA	4:A:899:HOH:O	1.79	0.81
1:A:272[A]:LEU:CD1	4:A:916:HOH:O	2.19	0.80
1:A:157[B]:GLN:NE2	1:A:170:GLU:OE1	2.18	0.76
1:A:152:TYR:CZ	1:A:190:SER:HB2	2.26	0.71
1:A:272[A]:LEU:CD2	4:A:916:HOH:O	2.34	0.70
1:A:152:TYR:C	1:A:177:THR:HG1	2.00	0.70
1:A:3:MET:CE	1:A:246[B]:ILE:HD11	2.22	0.70
1:A:71:LEU:O	4:A:803:HOH:O	2.12	0.67
1:A:152:TYR:CE2	1:A:190:SER:HB2	2.32	0.65
1:A:152:TYR:C	1:A:177:THR:OG1	2.42	0.63
1:A:80[A]:SER:O	4:A:802:HOH:O	2.07	0.60
1:A:83:LEU:HD11	1:A:226[B]:CYS:SG	2.42	0.59
1:A:54:HIS:HB2	2:A:701:TRS:H32	1.86	0.56
1:A:45:ARG:NH2	1:A:87:PRO:O	2.39	0.55
1:A:272[B]:LEU:HD22	4:A:916:HOH:O	2.08	0.53
1:A:79:ARG:CZ	1:A:233[B]:LEU:HD11	2.39	0.51
1:A:148:ASP:OD1	1:A:150:LYS:HE3	2.12	0.49
1:A:54:HIS:HB2	2:A:701:TRS:C3	2.43	0.48
1:A:9:GLN:HA	1:A:9:GLN:OE1	2.14	0.48
1:A:272[B]:LEU:HD13	4:A:916:HOH:O	2.13	0.48
1:A:3:MET:CE	1:A:246[B]:ILE:CD1	2.92	0.47
1:A:74:MET:HE2	1:A:230:THR:HG21	1.97	0.47
1:A:152:TYR:O	1:A:177:THR:OG1	2.29	0.47
1:A:54:HIS:ND1	2:A:701:TRS:H21	2.30	0.46
1:A:272[B]:LEU:CD2	4:A:916:HOH:O	2.63	0.45
1:A:32[A]:SER:N	4:A:805:HOH:O	2.48	0.45
1:A:32[B]:SER:N	4:A:805:HOH:O	2.48	0.45
1:A:77:ALA:HB1	1:A:233[A]:LEU:HD23	2.00	0.44
1:A:75:GLU:O	1:A:78[B]:GLN:NE2	2.36	0.43
1:A:3:MET:HE1	1:A:246[B]:ILE:HD11	1.95	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:SER:O	1:A:188:PRO:C	2.61	0.41
1:A:219:ILE:O	1:A:260:LEU:HA	2.19	0.41
1:A:28:SER:HB3	1:A:30:PHE:CE2	2.56	0.41
1:A:188:PRO:HA	1:A:269:PHE:CZ	2.56	0.41
1:A:272[B]:LEU:CD1	4:A:916:HOH:O	2.67	0.41
1:A:246[A]:ILE:CG2	1:A:271:TYR:CZ	3.04	0.41
1:A:76:GLU:OE2	1:A:238:ARG:NH1	2.54	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:MET:SD	4:A:920:HOH:O[2_655]	1.63	0.57
4:A:845:HOH:O	4:A:856:HOH:O[4_655]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	309/321 (96%)	299 (97%)	9 (3%)	1 (0%)	36 25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/294 (97%)	276 (97%)	8 (3%)	38	20

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	ARG
1	A	139	ASN
1	A	150	LYS
1	A	216	SER
1	A	238	ARG
1	A	289	ASP
1	A	292	LYS
1	A	293	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	HIS
1	A	102	GLN
1	A	208	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	O1J	A	702	-	12,12,12	2.75	5 (41%)	14,14,14	1.48	2 (14%)
2	TRS	A	701	-	7,7,7	0.34	0	9,9,9	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	O1J	A	702	-	-	2/6/6/6	0/1/1/1
2	TRS	A	701	-	-	6/9/9/9	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	O1J	C7-C8	6.01	1.49	1.38
3	A	702	O1J	C5-C4	5.60	1.48	1.38
3	A	702	O1J	C1-C	2.61	1.58	1.50
3	A	702	O1J	C8-C3	2.26	1.43	1.38
3	A	702	O1J	C6-C5	2.25	1.43	1.38

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	O1J	C2-O2-C1	4.22	117.19	112.36
3	A	702	O1J	O-C-C1	2.17	123.88	113.66

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	TRS	C1-C-C3-O3
2	A	701	TRS	C2-C-C3-O3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	701	TRS	N-C-C3-O3
3	A	702	O1J	C3-C2-O2-C1
3	A	702	O1J	C-C1-O2-C2
2	A	701	TRS	N-C-C2-O2
2	A	701	TRS	C1-C-C2-O2
2	A	701	TRS	C3-C-C2-O2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	TRS	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/321 (92%)	5.13	172 (57%) 0 0	5, 32, 74, 116	77 (25%)

All (172) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	179	TRP	21.9
1	A	272[A]	LEU	21.0
1	A	294	LEU	18.8
1	A	187	SER	18.7
1	A	196	PHE	18.6
1	A	226[A]	CYS	18.5
1	A	243	SER	17.6
1	A	269	PHE	17.6
1	A	191	PHE	17.5
1	A	277	GLY	17.4
1	A	296	HIS	17.4
1	A	184	VAL	17.3
1	A	278	ALA	17.1
1	A	183	GLY	17.1
1	A	149	ILE	16.7
1	A	78[A]	GLN	16.7
1	A	241	PRO	16.6
1	A	280	PHE	16.5
1	A	180	PRO	16.5
1	A	185	PRO	16.5
1	A	178	THR	16.2
1	A	240	ASP	16.2
1	A	46	TYR	16.1
1	A	153	TYR	16.0
1	A	292	LYS	15.8
1	A	246[A]	ILE	15.7
1	A	244	VAL	15.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	152	TYR	15.5
1	A	193	ASN	15.5
1	A	151	SER	15.4
1	A	224	THR	15.4
1	A	291	TRP	15.4
1	A	233[A]	LEU	15.3
1	A	287	VAL	15.2
1	A	265	ASP	15.2
1	A	242	SER	15.1
1	A	298	ASP	15.0
1	A	236	ASP	14.4
1	A	186	GLU	14.4
1	A	155	VAL	14.3
1	A	279	LYS	14.1
1	A	237	LYS	13.9
1	A	281	ILE	13.9
1	A	72[A]	ILE	13.8
1	A	293	GLU	13.8
1	A	154	THR	13.7
1	A	283	GLY	13.3
1	A	80[A]	SER	13.3
1	A	150	LYS	13.2
1	A	238	ARG	13.2
1	A	235	MET	12.9
1	A	217	ALA	12.9
1	A	252[A]	GLU	12.8
1	A	297	GLU	12.8
1	A	181	ASP	12.7
1	A	216	SER	12.7
1	A	157[A]	GLN	12.6
1	A	182	PHE	12.4
1	A	290	GLN	12.1
1	A	276	GLU	11.8
1	A	288	GLN	11.8
1	A	285	SER	11.5
1	A	262	GLN	11.5
1	A	221	ARG	11.4
1	A	207[A]	GLU	10.9
1	A	115	GLU	10.9
1	A	25[A]	HIS	10.8
1	A	282	MET	10.7
1	A	239	LYS	10.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	253[A]	MET	10.4
1	A	289	ASP	10.3
1	A	295	SER	10.2
1	A	45	ARG	10.1
1	A	284	ASP	9.6
1	A	286	SER	9.5
1	A	24	ARG	8.5
1	A	32[A]	SER	8.0
1	A	12	LYS	7.2
1	A	23	ILE	6.8
1	A	105[A]	ARG	6.4
1	A	3	MET	5.8
1	A	166	GLN	5.8
1	A	7	PHE	5.6
1	A	8	GLU	5.4
1	A	5	LYS	5.4
1	A	16	TRP	5.1
1	A	139	ASN	5.0
1	A	41	LYS	4.9
1	A	131	LYS	4.9
1	A	161	GLU	4.9
1	A	29	ASP	4.7
1	A	75	GLU	4.6
1	A	264	ALA	4.6
1	A	140	LEU	4.3
1	A	164	THR	4.3
1	A	47	ARG	4.3
1	A	36	LYS	4.1
1	A	61	GLN	4.0
1	A	135	PHE	3.9
1	A	127	GLN	3.9
1	A	39	LYS	3.9
1	A	138	THR	3.8
1	A	63	ASP	3.8
1	A	2	GLU	3.8
1	A	251	LEU	3.7
1	A	33	ARG	3.7
1	A	15	SER	3.6
1	A	35	ALA	3.6
1	A	136	GLU	3.6
1	A	38	PRO	3.5
1	A	165	THR	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	42	ASN	3.4
1	A	21	GLN	3.3
1	A	234	LEU	3.3
1	A	48	ASP	3.2
1	A	116	LYS	3.2
1	A	248	LYS	3.2
1	A	10	ILE	3.1
1	A	77	ALA	3.1
1	A	4	GLU	3.1
1	A	40	ASN	3.1
1	A	67	ILE	3.1
1	A	208	HIS	3.0
1	A	158	LEU	3.0
1	A	163	LEU	3.0
1	A	62	GLU	3.0
1	A	27	ALA	3.0
1	A	202	GLY	2.9
1	A	197	LYS	2.9
1	A	134	ILE	2.8
1	A	128	LYS	2.8
1	A	37	LEU	2.8
1	A	250	LEU	2.7
1	A	188	PRO	2.7
1	A	195	LEU	2.7
1	A	271	TYR	2.6
1	A	160	LEU	2.6
1	A	227	LEU	2.6
1	A	26	GLU	2.6
1	A	261	ILE	2.6
1	A	86	GLY	2.6
1	A	270	SER	2.5
1	A	49	VAL	2.5
1	A	120	LYS	2.5
1	A	20	TYR	2.5
1	A	204	LEU	2.5
1	A	260	LEU	2.5
1	A	267	LEU	2.4
1	A	14	GLY	2.4
1	A	66	TYR	2.4
1	A	73	LYS	2.4
1	A	103	LYS	2.4
1	A	19	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	18	ALA	2.4
1	A	211	VAL	2.4
1	A	9	GLN	2.3
1	A	58	LYS	2.3
1	A	232	LEU	2.3
1	A	206	PRO	2.2
1	A	59	LEU	2.2
1	A	171	ILE	2.2
1	A	92	VAL	2.2
1	A	268	ARG	2.2
1	A	13	SER	2.2
1	A	203	SER	2.2
1	A	88	LEU	2.2
1	A	192	LEU	2.2
1	A	162	ASN	2.1
1	A	11	ASP	2.1
1	A	254	ARG	2.1
1	A	51	PRO	2.1
1	A	274	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	O1J	A	702	12/12	0.64	0.51	34,44,45,46	21
2	TRS	A	701	8/8	0.76	0.21	63,77,90,90	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.