



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 04:51 AM UTC

PDB ID : 8FRB / pdb_00008frb
Title : Mouse acidic mammalian chitinase, catalytic domain in complex with N,N'-diacetylchitobiose at pH 5.25
Authors : Diaz, R.E.; Fraser, J.S.
Deposited on : 2023-01-06
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	
1	B	397	
1	C	397	
1	D	397	
2	E	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 100%
2	G	2	 50% 50%
2	H	2	 100%
2	I	2	 50% 50%
2	J	2	 100%
2	K	2	 100%
3	L	2	 50% 50%
4	M	2	 50% 50%
4	N	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	G	2[A]	-	-	-	X
2	NAG	G	2[C]	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 25661 atoms, of which 11675 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acidic mammalian chitinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	369	Total	C	H	N	O	S	0	6	0
			5819	1924	2830	487	565	13			
1	B	370	Total	C	H	N	O	S	0	5	0
			5811	1918	2828	486	566	13			
1	C	370	Total	C	H	N	O	S	0	6	0
			5831	1927	2837	487	567	13			
1	D	369	Total	C	H	N	O	S	0	4	0
			5788	1912	2816	485	562	13			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	392	HIS	-	expression tag	UNP Q91XA9
A	393	HIS	-	expression tag	UNP Q91XA9
A	394	HIS	-	expression tag	UNP Q91XA9
A	395	HIS	-	expression tag	UNP Q91XA9
A	396	HIS	-	expression tag	UNP Q91XA9
A	397	HIS	-	expression tag	UNP Q91XA9
B	392	HIS	-	expression tag	UNP Q91XA9
B	393	HIS	-	expression tag	UNP Q91XA9
B	394	HIS	-	expression tag	UNP Q91XA9
B	395	HIS	-	expression tag	UNP Q91XA9
B	396	HIS	-	expression tag	UNP Q91XA9
B	397	HIS	-	expression tag	UNP Q91XA9
C	392	HIS	-	expression tag	UNP Q91XA9
C	393	HIS	-	expression tag	UNP Q91XA9
C	394	HIS	-	expression tag	UNP Q91XA9
C	395	HIS	-	expression tag	UNP Q91XA9
C	396	HIS	-	expression tag	UNP Q91XA9
C	397	HIS	-	expression tag	UNP Q91XA9
D	392	HIS	-	expression tag	UNP Q91XA9
D	393	HIS	-	expression tag	UNP Q91XA9
D	394	HIS	-	expression tag	UNP Q91XA9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	395	HIS	-	expression tag	UNP Q91XA9
D	396	HIS	-	expression tag	UNP Q91XA9
D	397	HIS	-	expression tag	UNP Q91XA9

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	2	Total 57	C 16	H 28	N 2	O 11	0	2	0
2	F	2	Total 114	C 32	H 56	N 4	O 22	0	2	0
2	G	2	Total 114	C 32	H 56	N 4	O 22	0	2	0
2	H	2	Total 57	C 16	H 28	N 2	O 11	0	2	0
2	I	2	Total 57	C 16	H 28	N 2	O 11	0	2	0
2	J	2	Total 114	C 32	H 56	N 4	O 22	0	2	0
2	K	2	Total 57	C 16	H 28	N 2	O 11	0	2	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetylaminio-2-deoxy-alpha-L-idopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	2	Total	C	H	N	O	0	2	0
			57	16	28	2	11			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	M	2	Total	C	H	N	O	0	2	0
			57	16	28	2	11			
4	N	2	Total	C	H	N	O	0	2	0
			57	16	28	2	11			

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	B	1	Total	Mg	0	0
			1	1		
5	C	1	Total	Mg	0	0
			1	1		
5	D	2	Total	Mg	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	402	Total	O	0	0
			402	402		
6	B	449	Total	O	0	0
			449	449		
6	C	415	Total	O	0	0
			415	415		
6	D	400	Total	O	0	0
			400	400		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acidic mammalian chitinase




- Molecule 1: Acidic mammalian chitinase



- Molecule 1: Acidic mammalian chitinase



- Molecule 1: Acidic mammalian chitinase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  50%  50%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  50%  50%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  100%

MAG1
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetylamino-2-deoxy-alpha-L-idopyranose

Chain L:  50% 50%


HSQ1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose

Chain M:  50% 50%

NDG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose

Chain N:  100%

NDG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	91.93Å 106.96Å 146.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.29 – 1.70 57.29 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (57.29-1.70) 100.0 (57.29-1.70)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.99 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.142 , 0.178 0.143 , 0.178	Depositor DCC
R_{free} test set	7993 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	12.5	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	25661	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 52.12 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.0888e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDG, NAG, HSQ, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.20	0/3079	0.52	4/4191 (0.1%)
1	B	0.19	0/3072	0.52	2/4182 (0.0%)
1	C	0.21	0/3084	0.52	4/4198 (0.1%)
1	D	0.18	0/3061	0.51	4/4167 (0.1%)
All	All	0.19	0/12296	0.52	14/16738 (0.1%)

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	360	TRP	CA-C-N	5.93	132.37	121.70
1	B	360	TRP	C-N-CA	5.93	132.37	121.70
1	D	57	ALA	CA-C-N	5.89	132.31	121.70
1	D	57	ALA	C-N-CA	5.89	132.31	121.70
1	A	360	TRP	CA-C-N	5.75	132.06	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2989	2830	2824	11	0
1	B	2983	2828	2822	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2994	2837	2830	9	0
1	D	2972	2816	2812	9	0
2	E	29	28	15	0	0
2	F	58	56	53	2	0
2	G	58	56	53	1	0
2	H	29	28	26	0	0
2	I	29	28	23	1	0
2	J	58	56	53	2	0
2	K	29	28	26	0	0
3	L	29	28	25	1	0
4	M	29	28	14	2	0
4	N	29	28	22	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	2	0	0	0	0
6	A	402	0	0	2	2
6	B	449	0	0	0	1
6	C	415	0	0	2	1
6	D	400	0	0	1	0
All	All	13986	11675	11598	39	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 39 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:782:HOH:O	2:F:2[A]:NAG:H81	1.84	0.76
1:A:138[B]:ASP:OD2	4:M:1[B]:NDG:H8C2	1.99	0.63
1:D:359[B]:ILE:HD11	1:D:383:LEU:HD12	1.91	0.52
1:D:54:LEU:HG	1:D:90:LEU:HD11	1.92	0.51
1:A:86:ARG:NH1	6:A:508:HOH:O	2.43	0.51

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:645:HOH:O	6:B:728:HOH:O[3_545]	2.05	0.15
6:A:852:HOH:O	6:C:869:HOH:O[4_556]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	373/397 (94%)	368 (99%)	5 (1%)	0	100	100
1	B	373/397 (94%)	367 (98%)	6 (2%)	0	100	100
1	C	374/397 (94%)	368 (98%)	6 (2%)	0	100	100
1	D	371/397 (94%)	367 (99%)	4 (1%)	0	100	100
All	All	1491/1588 (94%)	1470 (99%)	21 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/331 (95%)	310 (98%)	5 (2%)	55	41
1	B	315/331 (95%)	313 (99%)	2 (1%)	78	72
1	C	316/331 (96%)	314 (99%)	2 (1%)	78	72
1	D	313/331 (95%)	310 (99%)	3 (1%)	68	58
All	All	1259/1324 (95%)	1247 (99%)	12 (1%)	70	58

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	358	MET
1	C	391	THR
1	D	358	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	275	ASN
1	A	349	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	374	GLN
1	C	384	ASN
1	D	374	GLN
1	D	192	GLN
1	D	297	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

26 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	E	1[C]	2	15,15,15	2.66	6 (40%)	21,21,21	2.32	4 (19%)
2	NAG	E	2[C]	2	14,14,15	2.32	5 (35%)	17,19,21	1.47	2 (11%)
2	NAG	F	1[A]	2	15,15,15	2.37	7 (46%)	21,21,21	1.08	1 (4%)
2	NAG	F	1[B]	2	15,15,15	2.32	6 (40%)	21,21,21	1.05	1 (4%)
2	NAG	F	2[A]	2	14,14,15	2.24	6 (42%)	17,19,21	1.30	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	F	2[B]	2	14,14,15	2.41	5 (35%)	17,19,21	1.17	1 (5%)
2	NAG	G	1[A]	2	15,15,15	2.28	8 (53%)	21,21,21	1.20	1 (4%)
2	NAG	G	1[C]	2	15,15,15	2.35	6 (40%)	21,21,21	2.60	5 (23%)
2	NAG	G	2[A]	2	14,14,15	2.28	5 (35%)	17,19,21	1.09	0
2	NAG	G	2[C]	2	14,14,15	2.05	4 (28%)	17,19,21	1.22	2 (11%)
2	NAG	H	1[B]	2	15,15,15	1.96	3 (20%)	21,21,21	1.02	2 (9%)
2	NAG	H	2[B]	2	14,14,15	2.01	4 (28%)	17,19,21	0.87	0
2	NAG	I	1[C]	2	15,15,15	2.61	8 (53%)	21,21,21	2.35	4 (19%)
2	NAG	I	2[C]	2	14,14,15	2.55	6 (42%)	17,19,21	1.20	2 (11%)
2	NAG	J	1[A]	2	15,15,15	2.55	7 (46%)	21,21,21	0.97	1 (4%)
2	NAG	J	1[B]	2	15,15,15	2.32	5 (33%)	21,21,21	1.27	3 (14%)
2	NAG	J	2[A]	2	14,14,15	2.52	8 (57%)	17,19,21	1.09	1 (5%)
2	NAG	J	2[B]	2	14,14,15	2.20	8 (57%)	17,19,21	1.12	1 (5%)
2	NAG	K	1[C]	2	15,15,15	2.43	6 (40%)	21,21,21	2.50	4 (19%)
2	NAG	K	2[C]	2	14,14,15	1.84	6 (42%)	17,19,21	1.21	3 (17%)
3	HSQ	L	1[B]	3	15,15,15	1.76	3 (20%)	21,21,21	2.34	7 (33%)
3	NAG	L	2[B]	3	14,14,15	1.87	5 (35%)	17,19,21	1.06	1 (5%)
4	NDG	M	1[B]	4	15,15,15	2.38	4 (26%)	21,21,21	1.48	3 (14%)
4	NAG	M	2[B]	4	14,14,15	2.41	6 (42%)	17,19,21	1.65	3 (17%)
4	NDG	N	1[A]	4	15,15,15	2.77	7 (46%)	21,21,21	1.66	4 (19%)
4	NAG	N	2[A]	4	14,14,15	2.32	7 (50%)	17,19,21	1.00	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1[C]	2	-	2/6/26/26	0/1/1/1
2	NAG	E	2[C]	2	-	0/6/23/26	0/1/1/1
2	NAG	F	1[A]	2	-	0/6/26/26	0/1/1/1
2	NAG	F	1[B]	2	-	0/6/26/26	0/1/1/1
2	NAG	F	2[A]	2	-	2/6/23/26	0/1/1/1
2	NAG	F	2[B]	2	-	4/6/23/26	0/1/1/1
2	NAG	G	1[A]	2	-	0/6/26/26	0/1/1/1
2	NAG	G	1[C]	2	-	1/6/26/26	0/1/1/1
2	NAG	G	2[A]	2	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	G	2[C]	2	-	0/6/23/26	0/1/1/1
2	NAG	H	1[B]	2	-	0/6/26/26	0/1/1/1
2	NAG	H	2[B]	2	-	2/6/23/26	0/1/1/1
2	NAG	I	1[C]	2	-	1/6/26/26	0/1/1/1
2	NAG	I	2[C]	2	-	0/6/23/26	0/1/1/1
2	NAG	J	1[A]	2	-	0/6/26/26	0/1/1/1
2	NAG	J	1[B]	2	-	0/6/26/26	0/1/1/1
2	NAG	J	2[A]	2	-	2/6/23/26	0/1/1/1
2	NAG	J	2[B]	2	-	4/6/23/26	0/1/1/1
2	NAG	K	1[C]	2	-	1/6/26/26	0/1/1/1
2	NAG	K	2[C]	2	-	0/6/23/26	0/1/1/1
3	HSQ	L	1[B]	3	-	0/6/26/26	0/1/1/1
3	NAG	L	2[B]	3	-	2/6/23/26	0/1/1/1
4	NDG	M	1[B]	4	-	0/6/26/26	0/1/1/1
4	NAG	M	2[B]	4	-	0/6/23/26	0/1/1/1
4	NDG	N	1[A]	4	-	0/6/26/26	0/1/1/1
4	NAG	N	2[A]	4	-	0/6/23/26	0/1/1/1

The worst 5 of 151 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	2[C]	NAG	C1-C2	6.66	1.61	1.52
2	F	2[B]	NAG	C1-C2	6.30	1.60	1.52
4	M	2[B]	NAG	C1-C2	6.25	1.60	1.52
2	E	2[C]	NAG	C1-C2	6.21	1.60	1.52
4	N	1[A]	NDG	C1-C2	6.16	1.60	1.52

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	1[C]	NAG	O5-C1-C2	10.25	119.81	109.52
2	K	1[C]	NAG	O5-C1-C2	9.80	119.36	109.52
2	E	1[C]	NAG	O5-C1-C2	8.56	118.12	109.52
2	I	1[C]	NAG	O5-C1-C2	8.07	117.62	109.52
3	L	1[B]	HSQ	O5-C5-C4	5.46	119.53	109.70

There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	1[C]	NAG	C1-C2-N2-C7

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	G	1[C]	NAG	C1-C2-N2-C7
2	I	1[C]	NAG	C1-C2-N2-C7
2	K	1[C]	NAG	C1-C2-N2-C7
2	F	2[B]	NAG	O5-C5-C6-O6


There are no ring outliers.


8 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	1[A]	NAG	1	0
2	G	1[A]	NAG	1	0
2	J	2[B]	NAG	1	0
2	J	1[B]	NAG	1	0
2	F	2[A]	NAG	1	0
3	L	2[B]	NAG	1	0
2	I	1[C]	NAG	1	0
4	M	1[B]	NDG	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.


























5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	369/397 (92%)	-0.58	1 (0%) 90 91	5, 14, 31, 55	6 (1%)
1	B	370/397 (93%)	-0.67	1 (0%) 90 91	4, 13, 28, 54	5 (1%)
1	C	370/397 (93%)	-0.68	3 (0%) 82 85	5, 13, 27, 69	6 (1%)
1	D	369/397 (92%)	-0.57	2 (0%) 87 89	5, 14, 33, 59	4 (1%)
All	All	1478/1588 (93%)	-0.62	7 (0%) 87 89	4, 13, 31, 69	21 (1%)

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	391	THR	4.9
1	D	390	SER	2.7
1	C	171	ILE	2.7
1	D	86	ARG	2.6
1	A	278	ASP	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	G	2[A]	14/15	0.49	0.52	17,22,27,32	28
2	NAG	G	2[C]	14/15	0.49	0.52	8,13,18,18	28

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	N	2[A]	14/15	0.77	0.27	19,25,38,57	28
2	NAG	I	1[C]	15/15	0.78	0.17	8,18,25,27	29
2	NAG	E	1[C]	15/15	0.79	0.12	9,16,20,25	29
4	NDG	N	1[A]	15/15	0.84	0.17	15,25,33,39	29
2	NAG	E	2[C]	14/15	0.85	0.14	4,18,29,29	28
2	NAG	I	2[C]	14/15	0.85	0.15	15,25,36,37	28
4	NAG	M	2[B]	14/15	0.87	0.13	6,15,23,34	28
2	NAG	J	2[B]	14/15	0.90	0.11	13,19,26,35	28
2	NAG	J	2[A]	14/15	0.90	0.11	11,20,31,37	28
2	NAG	J	1[B]	15/15	0.91	0.09	15,24,32,37	29
2	NAG	F	1[A]	15/15	0.91	0.09	14,22,29,29	29
2	NAG	F	1[B]	15/15	0.91	0.09	16,22,28,28	29
2	NAG	F	2[A]	14/15	0.91	0.10	12,18,23,29	28
2	NAG	F	2[B]	14/15	0.91	0.10	12,20,25,29	28
2	NAG	J	1[A]	15/15	0.91	0.09	16,23,28,29	29
4	NDG	M	1[B]	15/15	0.93	0.09	12,17,21,26	29
2	NAG	G	1[A]	15/15	0.93	0.07	8,13,16,18	29
2	NAG	G	1[C]	15/15	0.93	0.07	4,7,11,20	29
3	NAG	L	2[B]	14/15	0.93	0.07	10,12,21,29	0
3	HSQ	L	1[B]	15/15	0.94	0.07	9,16,26,29	0
2	NAG	H	2[B]	14/15	0.94	0.07	8,13,18,29	0
2	NAG	K	1[C]	15/15	0.96	0.06	8,12,16,34	0
2	NAG	H	1[B]	15/15	0.97	0.05	9,17,22,24	0
2	NAG	K	2[C]	14/15	0.97	0.05	8,12,17,18	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain L:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain M:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	MG	B	401	1/1	0.96	0.06	17,17,17,17	1
5	MG	C	401	1/1	0.96	0.05	20,20,20,20	1
5	MG	A	401	1/1	0.98	0.03	18,18,18,18	1
5	MG	D	401	1/1	0.99	0.02	21,21,21,21	1
5	MG	D	402	1/1	0.99	0.03	19,19,19,19	1

6.5 Other polymers [i](#)

There are no such residues in this entry.