



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:14 AM UTC

PDB ID : 3FSN / pdb_00003fsn
Title : Crystal structure of RPE65 at 2.14 angstrom resolution
Authors : Kiser, P.D.; Lodowski, D.T.; Palczewski, K.
Deposited on : 2009-01-11
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	533	
1	B	533	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8635 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retinal pigment epithelium-specific 65 kDa protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	513	Total	C	N	O	S	0	3	0
			4161	2696	683	767	15			
1	B	514	Total	C	N	O	S	0	3	0
			4172	2704	686	768	14			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	341	LEU	SER	variant	UNP Q28175
B	341	LEU	SER	variant	UNP Q28175

- Molecule 2 is FE (II) ION (CCD ID: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		
2	B	1	Total	Fe	0	0
			1	1		

- Molecule 3 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	1
			13	8	5		
3	B	1	Total	C	O	0	1
			13	8	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	131	Total	O	0	0
			131	131		
4	B	143	Total	O	0	0
			143	143		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Retinal pigment epithelium-specific 65 kDa protein



- Molecule 1: Retinal pigment epithelium-specific 65 kDa protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	176.52Å 176.52Å 86.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.22 – 2.14 48.22 – 2.14	Depositor EDS
% Data completeness (in resolution range)	99.4 (48.22-2.14) 99.5 (48.22-2.14)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.14Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.180 , 0.216 (Not available) , 0.214	Depositor DCC
R_{free} test set	4299 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	32.2	Xtriage
Anisotropy	0.403	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 26.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.065 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8635	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/4292	0.83	7/5843 (0.1%)
1	B	0.80	0/4305	0.82	3/5862 (0.1%)
All	All	0.78	0/8597	0.82	10/11705 (0.1%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	271	TRP	N-CA-C	7.32	119.40	107.32
1	A	201	SER	N-CA-C	7.30	120.29	108.76
1	A	321	HIS	CA-C-N	-7.09	108.00	121.54
1	A	321	HIS	C-N-CA	-7.09	108.00	121.54
1	B	321	HIS	CA-C-N	-6.53	112.01	122.66
1	B	321	HIS	C-N-CA	-6.53	112.01	122.66
1	A	226	VAL	CB-CA-C	-6.36	103.70	112.04
1	A	134	VAL	N-CA-C	6.14	117.16	111.45
1	A	137	TYR	N-CA-C	5.97	115.17	108.25
1	B	201	SER	N-CA-C	5.38	118.20	108.69

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4161	0	4061	33	0
1	B	4172	0	4070	27	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	13	0	18	0	0
3	B	13	0	18	3	0
4	A	131	0	0	1	0
4	B	143	0	0	0	0
All	All	8635	0	8167	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195[B]:CYS:SG	1:A:196:PHE:N	2.54	0.80
1:B:291:ILE:HD13	1:B:324:LEU:HD12	1.67	0.77
1:B:291:ILE:CD1	1:B:324:LEU:HD12	2.23	0.68
1:A:333:GLY:HA2	1:B:394:ILE:HD11	1.74	0.68
1:A:103:PHE:CZ	1:A:276:MET:HE2	2.28	0.67
1:B:248:ASN:HB3	3:B:535[B]:PG4:H82	1.79	0.64
1:A:267:SER:CB	1:A:273:ALA:HB2	2.28	0.63
1:B:270:LEU:HD12	1:B:271:TRP:N	2.16	0.59
1:A:460:TRP:CZ2	1:A:462:GLU:HG3	2.38	0.59
1:A:264:PHE:HB2	1:A:278:CYS:SG	2.42	0.58
1:A:205:ASN:HB3	1:A:225:ILE:HD12	1.86	0.56
1:A:491:VAL:HG12	1:A:493:PRO:HD3	1.87	0.56
1:B:270:LEU:HD12	1:B:271:TRP:H	1.70	0.56
1:A:26:LEU:C	1:A:26:LEU:HD23	2.31	0.56
1:B:159:THR:OG1	1:B:161:GLU:HG2	2.07	0.54
1:A:276:MET:HG2	1:A:342:TYR:OH	2.08	0.53
1:B:205:ASN:HB3	1:B:225:ILE:HD12	1.90	0.53
1:B:460:TRP:CZ2	1:B:462:GLU:HG3	2.44	0.53
1:A:103:PHE:CE2	1:A:276:MET:HE2	2.44	0.53
1:B:491:VAL:HG12	1:B:493:PRO:HD3	1.91	0.53
1:A:333:GLY:HA2	1:B:394:ILE:CD1	2.38	0.52
1:A:172:VAL:HG23	1:A:174:VAL:HG22	1.93	0.51
1:A:267:SER:HB2	1:A:273:ALA:HB2	1.91	0.51
1:A:267:SER:HB3	1:A:273:ALA:HB2	1.91	0.51
1:A:455:LYS:NZ	4:A:723:HOH:O	2.38	0.50
1:B:310:ASN:HB2	1:B:331:TRP:HB2	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:THR:HB	1:A:256:PRO:HD2	1.94	0.48
1:B:215:ASP:O	1:B:216:LYS:HB2	2.15	0.47
1:B:483:ASP:OD1	1:B:508:LYS:HE2	2.15	0.47
1:A:34:ILE:HD13	1:A:77:VAL:CG2	2.45	0.46
1:A:247:PRO:O	1:A:294:LYS:HE3	2.16	0.45
1:A:103:PHE:CE2	1:A:276:MET:CE	2.99	0.45
1:A:47:PRO:HD3	1:A:526:PHE:CE2	2.51	0.45
1:B:307:SER:HB2	1:B:308:PRO:HD2	1.98	0.44
1:B:248:ASN:OD1	3:B:535[B]:PG4:H81	2.17	0.44
1:B:173:SER:O	1:B:173:SER:OG	2.32	0.43
1:B:264:PHE:HB2	1:B:278:CYS:SG	2.58	0.43
1:B:218:ASP:OD1	1:B:218:ASP:C	2.62	0.43
1:A:103:PHE:CZ	1:A:276:MET:CE	2.99	0.43
1:B:322:GLU:OE2	1:B:322:GLU:HA	2.19	0.42
1:A:15:LEU:HD23	1:A:491:VAL:HG21	2.01	0.42
1:A:71:ASP:O	1:A:77:VAL:HA	2.19	0.42
1:A:101:THR:HG23	1:A:106:CYS:HA	2.02	0.42
1:A:394:ILE:HD11	1:B:333:GLY:HA2	2.02	0.42
1:A:218:ASP:O	1:A:221:SER:OG	2.34	0.42
1:A:34:ILE:HG12	1:A:504:ILE:HD12	2.02	0.41
1:B:248:ASN:CB	3:B:535[B]:PG4:H82	2.48	0.41
1:A:307:SER:HB2	1:A:308:PRO:HD2	2.01	0.41
1:B:100:ILE:HD13	1:B:127:GLU:HG2	2.02	0.41
1:B:415:ALA:O	1:B:438:GLY:HA2	2.21	0.41
1:A:529:LEU:HD12	1:A:529:LEU:C	2.46	0.41
1:B:420:GLN:HG3	1:B:470:PRO:O	2.21	0.40
1:A:141:GLU:HG3	1:A:212:LEU:HD22	2.02	0.40
1:A:226:VAL:HG13	1:A:297:LYS:HG2	2.04	0.40
1:A:264:PHE:O	1:A:267:SER:N	2.54	0.40
1:B:412:PRO:O	1:B:414[A]:GLN:HG2	2.21	0.40
1:B:503:LEU:HD21	1:B:505:LEU:HD21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	512/533 (96%)	499 (98%)	13 (2%)	0	100	100
1	B	513/533 (96%)	498 (97%)	15 (3%)	0	100	100
All	All	1025/1066 (96%)	997 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	459/475 (97%)	451 (98%)	8 (2%)	53	58
1	B	460/475 (97%)	456 (99%)	4 (1%)	70	77
All	All	919/950 (97%)	907 (99%)	12 (1%)	61	67

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	SER
1	A	26	LEU
1	A	174	VAL
1	A	183	ILE
1	A	215	ASP
1	A	310	ASN
1	A	358	ARG
1	A	519	GLU
1	B	3	SER
1	B	183	ILE
1	B	201	SER
1	B	310	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	HIS
1	A	205	ASN
1	A	228	GLN
1	A	461	GLN
1	A	497	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PG4	A	535[A]	-	12,12,12	0.64	0	11,11,11	0.46	0
3	PG4	B	535[B]	-	12,12,12	0.71	0	11,11,11	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PG4	A	535[A]	-	-	6/10/10/10	-
3	PG4	B	535[B]	-	-	7/10/10/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	535[B]	PG4	O3-C5-C6-O4
3	B	535[B]	PG4	O4-C7-C8-O5
3	A	535[A]	PG4	O4-C7-C8-O5
3	A	535[A]	PG4	O1-C1-C2-O2
3	A	535[A]	PG4	C1-C2-O2-C3
3	A	535[A]	PG4	C4-C3-O2-C2
3	B	535[B]	PG4	C1-C2-O2-C3
3	B	535[B]	PG4	O1-C1-C2-O2
3	A	535[A]	PG4	C3-C4-O3-C5
3	B	535[B]	PG4	C6-C5-O3-C4
3	A	535[A]	PG4	C5-C6-O4-C7
3	B	535[B]	PG4	C8-C7-O4-C6
3	B	535[B]	PG4	C4-C3-O2-C2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	535[B]	PG4	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	513/533 (96%)	-1.27	0 100 100	20, 34, 45, 70	3 (0%)
1	B	514/533 (96%)	-1.27	0 100 100	20, 34, 47, 72	3 (0%)
All	All	1027/1066 (96%)	-1.27	0 100 100	20, 34, 46, 72	6 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PG4	B	535[B]	13/13	0.97	0.07	65,66,67,67	13
3	PG4	A	535[A]	13/13	0.98	0.06	45,52,53,53	13
2	FE2	A	534	1/1	1.00	0.02	28,28,28,28	0
2	FE2	B	534	1/1	1.00	0.02	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.