



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 18, 2026 – 08:15 AM UTC

PDB ID : 7FS2 / pdb\_00007fs2  
Title : Structure of liver pyruvate kinase in complex with allosteric modulator 13  
Authors : Lulla, A.; Nilsson, O.; Brear, P.; Nain-Perez, A.; Grotli, M.; Hyvonen, M.  
Deposited on : 2022-12-18  
Resolution : 2.37 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7164 (2.40-2.36)
Clashscore	190562	7722 (2.40-2.36)
Ramachandran outliers	187476	7626 (2.40-2.36)
Sidechain outliers	187428	7627 (2.40-2.36)
RSRZ outliers	180081	7170 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	447	 12% 83% 11% • 6%
1	B	447	 13% 87% 9% ••
1	C	447	 6% 84% 10% • 5%
1	D	447	 2% 87% 7% • 5%
1	E	447	 10% 84% 8% • 6%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	447	
1	G	447	
1	H	447	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	OXL	A	602	-	X	-	-
3	OXL	B	602	-	X	-	-
3	OXL	C	602	-	X	-	-
3	OXL	D	602	-	X	-	-
3	OXL	E	602	-	X	-	-
3	OXL	F	602	-	X	-	-
3	OXL	G	602	-	X	-	-
3	OXL	H	602	-	X	-	-

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 27517 atoms, of which 84 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate kinase PKLR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	422	Total 3236	C 2034	N 585	O 597	S 20	0	6	0
1	B	436	Total 3329	C 2090	N 604	O 615	S 20	0	4	0
1	C	425	Total 3247	C 2040	N 585	O 603	S 19	0	4	0
1	D	425	Total 3252	C 2042	N 590	O 601	S 19	0	6	0
1	E	419	Total 3210	C 2018	N 579	O 593	S 20	0	5	0
1	F	432	Total 3321	C 2090	N 597	O 614	S 20	0	7	0
1	G	421	Total 3231	C 2031	N 581	O 600	S 19	0	6	0
1	H	425	Total 3251	C 2040	N 594	O 598	S 19	0	4	0

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P30613
A	0	SER	-	expression tag	UNP P30613
A	1	MET	-	expression tag	UNP P30613
A	2	GLU	-	expression tag	UNP P30613
A	12	ASP	SER	conflict	UNP P30613
A	130	GLY	-	linker	UNP P30613
A	131	SER	-	linker	UNP P30613
A	132	GLY	-	linker	UNP P30613
B	-1	GLY	-	expression tag	UNP P30613
B	0	SER	-	expression tag	UNP P30613
B	1	MET	-	expression tag	UNP P30613
B	2	GLU	-	expression tag	UNP P30613
B	12	ASP	SER	conflict	UNP P30613

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	130	GLY	-	linker	UNP P30613
B	131	SER	-	linker	UNP P30613
B	132	GLY	-	linker	UNP P30613
C	-1	GLY	-	expression tag	UNP P30613
C	0	SER	-	expression tag	UNP P30613
C	1	MET	-	expression tag	UNP P30613
C	2	GLU	-	expression tag	UNP P30613
C	12	ASP	SER	conflict	UNP P30613
C	130	GLY	-	linker	UNP P30613
C	131	SER	-	linker	UNP P30613
C	132	GLY	-	linker	UNP P30613
D	-1	GLY	-	expression tag	UNP P30613
D	0	SER	-	expression tag	UNP P30613
D	1	MET	-	expression tag	UNP P30613
D	2	GLU	-	expression tag	UNP P30613
D	12	ASP	SER	conflict	UNP P30613
D	130	GLY	-	linker	UNP P30613
D	131	SER	-	linker	UNP P30613
D	132	GLY	-	linker	UNP P30613
E	-1	GLY	-	expression tag	UNP P30613
E	0	SER	-	expression tag	UNP P30613
E	1	MET	-	expression tag	UNP P30613
E	2	GLU	-	expression tag	UNP P30613
E	12	ASP	SER	conflict	UNP P30613
E	228	GLY	-	linker	UNP P30613
E	229	SER	-	linker	UNP P30613
E	230	GLY	-	linker	UNP P30613
F	-1	GLY	-	expression tag	UNP P30613
F	0	SER	-	expression tag	UNP P30613
F	1	MET	-	expression tag	UNP P30613
F	2	GLU	-	expression tag	UNP P30613
F	12	ASP	SER	conflict	UNP P30613
F	228	GLY	-	linker	UNP P30613
F	229	SER	-	linker	UNP P30613
F	230	GLY	-	linker	UNP P30613
G	-1	GLY	-	expression tag	UNP P30613
G	0	SER	-	expression tag	UNP P30613
G	1	MET	-	expression tag	UNP P30613
G	2	GLU	-	expression tag	UNP P30613
G	12	ASP	SER	conflict	UNP P30613
G	130	GLY	-	linker	UNP P30613
G	131	SER	-	linker	UNP P30613

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	132	GLY	-	linker	UNP P30613
H	-1	GLY	-	expression tag	UNP P30613
H	0	SER	-	expression tag	UNP P30613
H	1	MET	-	expression tag	UNP P30613
H	2	GLU	-	expression tag	UNP P30613
H	12	ASP	SER	conflict	UNP P30613
H	130	GLY	-	linker	UNP P30613
H	131	SER	-	linker	UNP P30613
H	132	GLY	-	linker	UNP P30613

- Molecule 2 is 1,6-di-O-phosphono-beta-D-fructofuranose (CCD ID: FBP) (formula:  $C_6H_{14}O_{12}P_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	C	O	P	0	0
			20	6	12	2		
2	B	1	Total	C	O	P	0	0
			20	6	12	2		
2	C	1	Total	C	O	P	0	0
			20	6	12	2		
2	D	1	Total	C	O	P	0	0
			20	6	12	2		
2	E	1	Total	C	O	P	0	0
			20	6	12	2		
2	F	1	Total	C	O	P	0	0
			20	6	12	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	G	1	Total	C	O	P	0	0
			20	6	12	2		
2	H	1	Total	C	O	P	0	0
			20	6	12	2		

- Molecule 3 is OXALATE ION (CCD ID: OXL) (formula: C<sub>2</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	2	4		
3	B	1	Total	C	O	0	0
			6	2	4		
3	C	1	Total	C	O	0	0
			6	2	4		
3	D	1	Total	C	O	0	0
			6	2	4		
3	E	1	Total	C	O	0	0
			6	2	4		
3	F	1	Total	C	O	0	0
			6	2	4		
3	G	1	Total	C	O	0	0
			6	2	4		
3	H	1	Total	C	O	0	0
			6	2	4		


- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	B	1	Total Mg 1 1	0	0
4	C	1	Total Mg 1 1	0	0
4	D	1	Total Mg 1 1	0	0
4	E	1	Total Mg 1 1	0	0
4	F	1	Total Mg 1 1	0	0
4	G	1	Total Mg 1 1	0	0
4	H	1	Total Mg 1 1	0	0

- Molecule 5 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total K 1 1	0	0
5	B	1	Total K 1 1	0	0
5	C	1	Total K 1 1	0	0
5	D	1	Total K 1 1	0	0
5	E	1	Total K 1 1	0	0
5	F	1	Total K 1 1	0	0
5	G	1	Total K 1 1	0	0
5	H	1	Total K 1 1	0	0

- Molecule 6 is (1P)-3',4,4',5-tetrahydroxy-N-{{4-(3-hydroxybenzene-1-sulfonyl)phenyl}methyl}[1,1'-biphenyl]-2-sulfonamide (CCD ID: O7X) (formula: C<sub>25</sub>H<sub>21</sub>NO<sub>9</sub>S<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			S
6	A	1	58	25	21	1	9	2	21	0
6	B	1	58	25	21	1	9	2	21	0
6	F	1	58	25	21	1	9	2	21	0
6	G	1	58	25	21	1	9	2	21	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	65	Total	O	0	0
			65	65		
7	B	84	Total	O	0	0
			84	84		
7	C	121	Total	O	0	0
			121	121		
7	D	156	Total	O	0	0
			156	156		
7	E	90	Total	O	0	0
			90	90		
7	F	120	Total	O	0	0
			120	120		
7	G	164	Total	O	0	0
			164	164		
7	H	184	Total	O	0	0
			184	184		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pyruvate kinase PKLR




- Molecule 1: Pyruvate kinase PKLR





- Molecule 1: Pyruvate kinase PKLR









- Molecule 1: Pyruvate kinase PKLR




- Molecule 1: Pyruvate kinase PKLR




- Molecule 1: Pyruvate kinase PKLR



- Molecule 1: Pyruvate kinase PKLR



- Molecule 1: Pyruvate kinase PKLR



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	207.67Å 113.01Å 187.77Å 90.00° 92.41° 90.00°	Depositor
Resolution (Å)	187.60 – 2.37 187.60 – 2.37	Depositor EDS
% Data completeness (in resolution range)	67.7 (187.60-2.37) 67.7 (187.60-2.37)	Depositor EDS
$R_{merge}$	0.22	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.53 (at 2.37Å)	Xtrriage
Refinement program	BUSTER 2.10.4 (16-JUL-2021)	Depositor
R, $R_{free}$	0.203 , 0.239 0.193 , 0.224	Depositor DCC
$R_{free}$ test set	6074 reflections (3.47%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	43.2	Xtrriage
Anisotropy	0.025	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 52.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.002 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	27517	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 54.12 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.8203e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: O7X, FBP, K, OXL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	1/3307 (0.0%)	1.04	5/4469 (0.1%)
1	B	0.71	1/3396 (0.0%)	1.04	2/4592 (0.0%)
1	C	0.74	1/3313 (0.0%)	1.04	3/4479 (0.1%)
1	D	0.75	0/3326	1.04	4/4497 (0.1%)
1	E	0.68	0/3278	1.04	2/4430 (0.0%)
1	F	0.75	1/3396 (0.0%)	1.05	6/4591 (0.1%)
1	G	0.76	2/3303 (0.1%)	1.03	3/4465 (0.1%)
1	H	0.79	0/3316	1.07	3/4483 (0.1%)
All	All	0.73	6/26635 (0.0%)	1.05	28/36006 (0.1%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	374	SER	CA-C	6.00	1.55	1.52
1	G	525	THR	CA-CB	5.38	1.61	1.53
1	B	374	SER	CA-C	5.21	1.55	1.52
1	A	374	SER	CA-C	5.20	1.55	1.52
1	G	537	MET	SD-CE	-5.15	1.66	1.79

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	422	GLU	CB-CG-CD	7.78	125.82	112.60
1	A	422	GLU	CB-CG-CD	7.27	124.96	112.60
1	D	453	LEU	CA-C-N	6.23	128.62	120.28
1	D	453	LEU	C-N-CA	6.23	128.62	120.28
1	B	453	LEU	CA-C-N	6.03	128.36	120.28

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3236	0	3299	25	0
1	B	3329	0	3394	23	1
1	C	3247	0	3299	26	0
1	D	3252	0	3310	16	0
1	E	3210	0	3270	23	0
1	F	3321	0	3393	28	0
1	G	3231	0	3284	28	0
1	H	3251	0	3306	27	0
2	A	20	0	10	2	0
2	B	20	0	10	0	0
2	C	20	0	10	0	0
2	D	20	0	10	0	0
2	E	20	0	10	0	0
2	F	20	0	10	0	0
2	G	20	0	10	0	0
2	H	20	0	10	0	0
3	A	6	0	0	0	0
3	B	6	0	0	0	0
3	C	6	0	0	0	0
3	D	6	0	0	0	0
3	E	6	0	0	0	0
3	F	6	0	0	0	0
3	G	6	0	0	0	0
3	H	6	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	F	1	0	0	0	0
5	G	1	0	0	0	0
5	H	1	0	0	0	0
6	A	37	21	0	1	0
6	B	37	21	0	3	0
6	F	37	21	0	3	0
6	G	37	21	0	0	0
7	A	65	0	0	0	0
7	B	84	0	0	1	0
7	C	121	0	0	0	0
7	D	156	0	0	0	0
7	E	90	0	0	0	0
7	F	120	0	0	2	0
7	G	164	0	0	0	0
7	H	184	0	0	2	0
All	All	27433	84	26635	174	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 174 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:411:ARG:HG3	1:C:426:ILE:HD11	1.49	0.94
1:A:408:GLU:OE2	1:B:411:ARG:NH2	2.01	0.93
1:B:411:ARG:HG3	1:B:426:ILE:HD11	1.57	0.85
1:G:537:MET:HE3	1:H:537:MET:HG2	1.64	0.79
1:G:422[A]:GLU:HG3	1:G:452:LEU:HD13	1.69	0.75

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:511:LEU:O	1:B:511:LEU:O[2_555]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	424/447 (95%)	419 (99%)	4 (1%)	1 (0%)	43	56
1	B	438/447 (98%)	433 (99%)	4 (1%)	1 (0%)	43	56
1	C	425/447 (95%)	418 (98%)	5 (1%)	2 (0%)	24	34
1	D	429/447 (96%)	424 (99%)	4 (1%)	1 (0%)	43	56
1	E	420/447 (94%)	416 (99%)	3 (1%)	1 (0%)	43	56
1	F	435/447 (97%)	431 (99%)	3 (1%)	1 (0%)	43	56
1	G	423/447 (95%)	415 (98%)	6 (1%)	2 (0%)	24	34
1	H	427/447 (96%)	421 (99%)	5 (1%)	1 (0%)	43	56
All	All	3421/3576 (96%)	3377 (99%)	34 (1%)	10 (0%)	36	48

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	271	GLY
1	G	271	GLY
1	A	340	THR
1	B	340	THR
1	C	340	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	340/352 (97%)	325 (96%)	15 (4%)	25	40

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	349/352 (99%)	333 (95%)	16 (5%)	24	38
1	C	341/352 (97%)	334 (98%)	7 (2%)	47	67
1	D	342/352 (97%)	331 (97%)	11 (3%)	34	53
1	E	338/352 (96%)	325 (96%)	13 (4%)	29	46
1	F	350/352 (99%)	333 (95%)	17 (5%)	22	36
1	G	340/352 (97%)	332 (98%)	8 (2%)	43	63
1	H	340/352 (97%)	334 (98%)	6 (2%)	51	71
All	All	2740/2816 (97%)	2647 (97%)	93 (3%)	35	51

5 of 93 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	412	ARG
1	F	284	GLU
1	E	537[A]	MET
1	F	76[A]	MET
1	F	517	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	90	HIS
1	G	470	GLN
1	E	390	GLN
1	H	390	GLN
1	G	275	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 36 ligands modelled in this entry, 16 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FBP	C	601	-	18,20,20	0.87	1 (5%)	21,32,32	0.70	0
2	FBP	D	601	-	18,20,20	0.89	1 (5%)	21,32,32	1.05	1 (4%)
3	OXL	A	602	4	5,5,5	2.05	2 (40%)	6,6,6	2.14	3 (50%)
2	FBP	F	601	-	18,20,20	0.50	0	21,32,32	0.85	0
2	FBP	H	601	-	18,20,20	1.03	1 (5%)	21,32,32	1.28	1 (4%)
3	OXL	D	602	4	5,5,5	2.02	2 (40%)	6,6,6	3.27	4 (66%)
3	OXL	B	602	4	5,5,5	2.26	2 (40%)	6,6,6	1.73	2 (33%)
3	OXL	F	602	4	5,5,5	3.00	4 (80%)	6,6,6	2.39	3 (50%)
6	O7X	G	605	-	40,40,40	0.33	0	59,60,60	0.78	1 (1%)
3	OXL	H	602	4	5,5,5	2.11	2 (40%)	6,6,6	1.65	1 (16%)
2	FBP	G	601	-	18,20,20	0.50	0	21,32,32	1.04	2 (9%)
3	OXL	E	602	4	5,5,5	2.41	2 (40%)	6,6,6	1.11	1 (16%)
3	OXL	C	602	4	5,5,5	2.02	2 (40%)	6,6,6	1.07	1 (16%)
2	FBP	A	601	-	18,20,20	0.57	0	21,32,32	0.70	0
6	O7X	B	605	-	40,40,40	0.32	0	59,60,60	0.88	3 (5%)
2	FBP	E	601	-	18,20,20	0.64	0	21,32,32	0.91	1 (4%)
6	O7X	F	605	-	40,40,40	0.32	0	59,60,60	0.95	3 (5%)
6	O7X	A	605	-	40,40,40	0.19	0	59,60,60	0.80	3 (5%)
3	OXL	G	602	4	5,5,5	2.33	3 (60%)	6,6,6	2.59	3 (50%)
2	FBP	B	601	-	18,20,20	0.55	0	21,32,32	0.84	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.  
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FBP	C	601	-	-	6/13/32/32	0/1/1/1
2	FBP	D	601	-	-	2/13/32/32	0/1/1/1
3	OXL	A	602	4	-	4/4/4/4	-
2	FBP	F	601	-	-	2/13/32/32	0/1/1/1
2	FBP	H	601	-	-	5/13/32/32	0/1/1/1
3	OXL	D	602	4	-	4/4/4/4	-
3	OXL	B	602	4	-	4/4/4/4	-
3	OXL	F	602	4	-	4/4/4/4	-
6	O7X	G	605	-	-	2/28/28/28	0/4/4/4
3	OXL	H	602	4	-	4/4/4/4	-
2	FBP	G	601	-	-	2/13/32/32	0/1/1/1
3	OXL	E	602	4	-	4/4/4/4	-
3	OXL	C	602	4	-	4/4/4/4	-
2	FBP	A	601	-	-	7/13/32/32	0/1/1/1
6	O7X	B	605	-	-	1/28/28/28	0/4/4/4
2	FBP	E	601	-	-	3/13/32/32	0/1/1/1
6	O7X	F	605	-	-	2/28/28/28	0/4/4/4
6	O7X	A	605	-	-	1/28/28/28	0/4/4/4
3	OXL	G	602	4	-	4/4/4/4	-
2	FBP	B	601	-	-	3/13/32/32	0/1/1/1

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	602	OXL	O1-C1	4.51	1.33	1.22
3	B	602	OXL	O2-C2	4.41	1.33	1.22
3	A	602	OXL	O2-C2	3.77	1.31	1.22
3	F	602	OXL	O2-C2	3.71	1.31	1.22
3	F	602	OXL	O1-C1	3.68	1.31	1.22

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	602	OXL	O4-C2-C1	5.40	123.31	112.83
2	H	601	FBP	O5P-P2-O6	4.56	118.56	106.67
3	G	602	OXL	O3-C1-C2	4.43	121.43	112.83

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	F	605	O7X	C11-C10-C15	4.17	124.92	120.25
3	D	602	OXL	O3-C1-C2	4.01	120.62	112.83

There are no chirality outliers.

5 of 68 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	FBP	O1-C1-C2-O2
2	A	601	FBP	O1-C1-C2-C3
2	A	601	FBP	O1-C1-C2-O5
2	A	601	FBP	C4-C5-C6-O6
2	B	601	FBP	C1-O1-P1-O2P


There are no ring outliers.


4 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	FBP	2	0
6	B	605	O7X	3	0
6	F	605	O7X	3	0
6	A	605	O7X	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






























## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	422/447 (94%)	1.02	52 (12%) 8 7	32, 62, 97, 119	6 (1%)
1	B	436/447 (97%)	0.90	57 (13%) 7 6	30, 54, 92, 99	4 (0%)
1	C	425/447 (95%)	0.49	27 (6%) 25 24	24, 49, 76, 111	4 (0%)
1	D	425/447 (95%)	0.07	9 (2%) 63 62	17, 37, 67, 118	6 (1%)
1	E	419/447 (93%)	0.84	45 (10%) 11 10	28, 56, 90, 109	5 (1%)
1	F	432/447 (96%)	0.43	19 (4%) 39 38	26, 46, 73, 93	7 (1%)
1	G	421/447 (94%)	0.16	18 (4%) 40 39	23, 41, 65, 92	7 (1%)
1	H	425/447 (95%)	-0.08	11 (2%) 57 56	17, 35, 60, 109	4 (0%)
All	All	3405/3576 (95%)	0.48	238 (6%) 22 21	17, 48, 85, 119	43 (1%)

The worst 5 of 238 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	511	LEU	7.0
1	F	231	PRO	6.7
1	A	232	GLY	6.0
1	B	517	VAL	5.6
1	B	115	LEU	5.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	K	E	604	1/1	0.86	0.12	125,125,125,125	0
3	OXL	E	602	6/6	0.87	0.13	71,71,72,72	0
3	OXL	C	602	6/6	0.89	0.14	74,74,75,75	0
6	O7X	A	605	37/37	0.89	0.15	57,63,70,70	21
6	O7X	B	605	37/37	0.89	0.17	53,62,72,72	21
6	O7X	G	605	37/37	0.89	0.17	55,64,71,72	21
3	OXL	A	602	6/6	0.90	0.09	79,80,80,80	0
6	O7X	F	605	37/37	0.90	0.15	49,56,65,67	21
5	K	A	604	1/1	0.90	0.10	90,90,90,90	0
3	OXL	B	602	6/6	0.91	0.11	61,62,63,63	0
3	OXL	F	602	6/6	0.91	0.12	67,68,69,70	0
5	K	C	604	1/1	0.92	0.10	74,74,74,74	0
4	MG	B	603	1/1	0.93	0.14	40,40,40,40	0
3	OXL	D	602	6/6	0.93	0.12	62,62,63,64	0
5	K	B	604	1/1	0.93	0.08	78,78,78,78	0
5	K	F	604	1/1	0.94	0.06	82,82,82,82	0
5	K	G	604	1/1	0.94	0.09	78,78,78,78	0
2	FBP	E	601	20/20	0.94	0.09	54,57,59,60	0
3	OXL	G	602	6/6	0.94	0.08	50,50,51,52	0
4	MG	A	603	1/1	0.94	0.11	41,41,41,41	0
2	FBP	B	601	20/20	0.94	0.08	54,57,58,58	0
2	FBP	A	601	20/20	0.95	0.08	56,59,61,61	0
2	FBP	F	601	20/20	0.96	0.06	44,51,54,55	0
3	OXL	H	602	6/6	0.96	0.10	56,57,58,58	0
4	MG	C	603	1/1	0.96	0.15	31,31,31,31	0
5	K	H	604	1/1	0.96	0.09	50,50,50,50	0
4	MG	D	603	1/1	0.97	0.15	26,26,26,26	0
5	K	D	604	1/1	0.97	0.08	51,51,51,51	0
2	FBP	C	601	20/20	0.98	0.05	32,34,34,35	0
2	FBP	H	601	20/20	0.98	0.06	24,29,32,32	0
4	MG	E	603	1/1	0.98	0.08	33,33,33,33	0
4	MG	F	603	1/1	0.98	0.13	27,27,27,27	0
2	FBP	G	601	20/20	0.99	0.04	31,35,37,38	0
2	FBP	D	601	20/20	0.99	0.05	29,32,34,34	0
4	MG	G	603	1/1	1.00	0.07	15,15,15,15	0
4	MG	H	603	1/1	1.00	0.06	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around O7X B 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around O7X G 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around O7X F 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FBP E 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FBP B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FBP A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FBP F 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FBP C 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around FBP H 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FBP G 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.