



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:04 PM UTC

PDB ID : 8FS1 / pdb_00008fs1
Title : CamA Adenine Methyltransferase Complexed to Cognate Substrate DNA and Inhibitor 11a (YD905)
Authors : Zhou, J.; Horton, J.R.; Cheng, X.
Deposited on : 2023-01-09
Resolution : 2.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1819 (2.76-2.72)
Clashscore	190562	1866 (2.76-2.72)
Ramachandran outliers	187476	1830 (2.76-2.72)
Sidechain outliers	187428	1831 (2.76-2.72)
RSRZ outliers	180081	1819 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	577	3% 85% 8% 6%
1	B	577	3% 86% 9% 5%
1	C	577	4% 84% 9% 7%
2	E	14	7% 50% 50%
2	G	14	79% 21%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	I	14	 86%14%
3	D	14	 71%29%
3	F	14	 86%14%
3	H	14	 93%7%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 15653 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Site-specific DNA-methyltransferase (adenine-specific).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	544	Total	C	N	O	S	0	0	0
			4478	2914	714	833	17			
1	B	551	Total	C	N	O	S	0	0	0
			4536	2954	729	836	17			
1	C	537	Total	C	N	O	S	0	0	0
			4295	2789	687	802	17			

- Molecule 2 is a DNA chain called DNA (5'-D(*AP*TP*GP*GP*GP*AP*CP*TP*TP*TP*TP*TP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	14	Total	C	N	O	P	0	0	0
			287	139	50	85	13			
2	G	14	Total	C	N	O	P	0	0	0
			287	139	50	85	13			
2	I	14	Total	C	N	O	P	0	0	0
			287	139	50	85	13			

- Molecule 3 is a DNA chain called DNA (5'-D(*AP*TP*GP*GP*GP*AP*CP*TP*TP*TP*TP*TP*GP*A)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	14	Total	C	N	O	P	0	0	0
			281	136	53	79	13			
3	F	14	Total	C	N	O	P	0	0	0
			281	136	53	79	13			
3	H	14	Total	C	N	O	P	0	0	0
			281	136	53	79	13			

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 5'-S-{2-[N'-(cyclohexylmethyl)carbamimidamido]ethyl}-N-(3-phenylpropyl)-5'-thioadenosine (CCD ID: YB0) (formula: C₂₉H₄₂N₈O₃S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			41	29	8	3	1		
5	B	1	Total	C	N	O	S	0	0
			41	29	8	3	1		
5	C	1	Total	C	N	O	S	0	0
			41	29	8	3	1		

- Molecule 6 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	K	0	0
			2	2		
6	B	3	Total	K	0	0
			3	3		
6	C	2	Total	K	0	0
			2	2		
6	D	1	Total	K	0	0
			1	1		
6	F	1	Total	K	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	147	Total	O	0	0
			147	147		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	162	Total 162	O 162	0	0
7	C	61	Total 61	O 61	0	0
7	E	15	Total 15	O 15	0	0
7	D	24	Total 24	O 24	0	0
7	F	25	Total 25	O 25	0	0
7	G	20	Total 20	O 20	0	0
7	H	16	Total 16	O 16	0	0
7	I	10	Total 10	O 10	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Site-specific DNA-methyltransferase (adenine-specific)



- Molecule 1: Site-specific DNA-methyltransferase (adenine-specific)




- Molecule 1: Site-specific DNA-methyltransferase (adenine-specific)








- Molecule 2: DNA (5'-D(*AP*TP*GP*GP*GP*AP*CP*TP*TP*TP*TP*TP*GP*A)-3')



- Molecule 2: DNA (5'-D(*AP*TP*GP*GP*GP*AP*CP*TP*TP*TP*TP*TP*GP*A)-3')




- Molecule 2: DNA (5'-D(*AP*TP*GP*GP*GP*AP*CP*TP*TP*TP*TP*TP*GP*A)-3')



- Molecule 3: DNA (5'-D(*AP*TP*GP*GP*GP*AP*CP*TP*TP*TP*TP*TP*GP*A)-3')



- Molecule 3: DNA (5'-D(*AP*TP*GP*GP*GP*AP*CP*TP*TP*TP*TP*TP*GP*A)-3')



- Molecule 3: DNA (5'-D(*AP*TP*GP*GP*GP*AP*CP*TP*TP*TP*TP*TP*GP*A)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	80.91Å 160.72Å 229.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.55 – 2.74 48.55 – 2.74	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.55-2.74) 99.6 (48.55-2.74)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.73Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.171 , 0.218 (Not available) , 0.195	Depositor DCC
R_{free} test set	2000 reflections (2.52%)	wwPDB-VP
Wilson B-factor (Å ²)	48.5	Xtriage
Anisotropy	0.373	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15653	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, YB0, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/4571	0.46	0/6159
1	B	0.26	0/4630	0.46	0/6234
1	C	0.23	0/4381	0.44	0/5917
2	E	0.37	0/321	0.56	0/495
2	G	0.39	0/321	0.63	0/495
2	I	0.33	0/321	0.55	0/495
3	D	0.34	0/315	0.54	0/483
3	F	0.36	0/315	0.54	0/483
3	H	0.31	0/315	0.55	0/483
All	All	0.26	0/15490	0.47	0/21244

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4478	0	4406	29	0
1	B	4536	0	4497	26	0
1	C	4295	0	4135	27	0
2	E	287	0	162	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	G	287	0	162	2	0
2	I	287	0	162	2	0
3	D	281	0	159	2	0
3	F	281	0	159	1	0
3	H	281	0	159	1	0
4	A	4	0	6	1	0
4	B	20	0	30	1	0
4	F	4	0	6	0	0
5	A	41	0	0	0	0
5	B	41	0	0	0	0
5	C	41	0	0	0	0
6	A	2	0	0	0	0
6	B	3	0	0	0	0
6	C	2	0	0	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
7	A	147	0	0	1	0
7	B	162	0	0	0	0
7	C	61	0	0	1	0
7	D	24	0	0	0	0
7	E	15	0	0	0	0
7	F	25	0	0	0	0
7	G	20	0	0	0	0
7	H	16	0	0	0	0
7	I	10	0	0	0	0
All	All	15653	0	14043	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:482:GLU:HG3	1:B:482:GLU:HG2	1.59	0.84
1:C:57:ARG:NH1	1:C:158:LYS:O	2.18	0.76
1:C:334:CYS:O	1:C:529:ARG:NH1	2.28	0.66
1:C:280:ILE:HD11	1:C:285:ILE:HG12	1.77	0.65
2:G:1:DA:H1'	2:G:2:DT:H5''	1.80	0.63
1:C:496:ILE:HD11	1:C:530:ILE:HD12	1.82	0.60
1:C:252:ILE:HD12	1:C:287:ILE:HD12	1.84	0.59
1:C:39:TYR:OH	1:C:243:GLU:OE2	2.16	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:13:DC:H2''	3:D:14:DA:C8	2.38	0.58
1:A:202:LYS:HD2	1:A:234:TYR:CZ	2.38	0.58
1:C:43:LYS:HD2	1:C:264:THR:HG21	1.86	0.58
1:B:252:ILE:HD12	1:B:287:ILE:HD12	1.86	0.57
1:A:319:VAL:HG11	1:A:508:ILE:HG23	1.87	0.56
1:A:168:TYR:CE1	1:A:219:PRO:HD3	2.41	0.56
1:A:252:ILE:HD12	1:A:287:ILE:HD12	1.88	0.56
2:E:1:DA:H2''	2:E:2:DT:H5''	1.86	0.56
2:E:2:DT:H2''	2:E:3:DG:C8	2.41	0.55
1:A:66:GLY:HA3	1:A:123:LEU:HD13	1.91	0.53
1:B:58:ILE:HD13	1:B:76:LEU:HD11	1.92	0.52
1:A:367:LEU:HD22	1:A:408:ARG:HG2	1.90	0.52
1:B:66:GLY:HA3	1:B:123:LEU:HD13	1.91	0.52
1:A:502:TYR:HA	1:A:505:TYR:HB3	1.92	0.52
1:A:281:LYS:HE3	1:A:303:GLU:HB3	1.91	0.52
1:B:168:TYR:CE1	1:B:219:PRO:HD3	2.45	0.51
1:C:317:ILE:HG22	1:C:319:VAL:HG22	1.93	0.51
1:C:516:ASN:ND2	7:C:701:HOH:O	2.42	0.51
1:C:61:ILE:HG23	1:C:150:SER:HB3	1.93	0.51
2:I:2:DT:H2''	2:I:3:DG:C8	2.45	0.51
1:C:168:TYR:CE1	1:C:219:PRO:HD3	2.46	0.51
1:C:58:ILE:HG13	1:C:161:TYR:HB2	1.93	0.51
1:A:143:ILE:HG22	1:A:145:LEU:HG	1.93	0.50
1:C:364:LYS:HE2	1:C:413:ILE:HG12	1.94	0.50
2:G:2:DT:H2''	2:G:3:DG:C8	2.47	0.50
1:B:280:ILE:HD11	1:B:285:ILE:CG1	2.42	0.49
1:B:88:LYS:HE3	4:B:601:EDO:O1	2.12	0.49
1:B:280:ILE:HD11	1:B:285:ILE:HG12	1.94	0.49
1:C:213:ILE:HD11	1:C:268:LYS:HE3	1.95	0.48
1:A:433:LYS:HB2	1:A:436:GLU:HG3	1.96	0.48
1:A:376:TRP:CZ2	1:A:472:SER:HB2	2.49	0.48
1:A:464:ILE:HG13	1:A:494:VAL:HG11	1.95	0.47
1:A:348:ILE:O	1:A:472:SER:HA	2.15	0.47
2:E:10:DT:H2''	2:E:11:DT:H72	1.97	0.47
1:B:512:LYS:HD3	1:B:518:TYR:CE2	2.50	0.46
1:A:176:LYS:HZ1	2:E:12:DT:P	2.39	0.46
1:C:284:ASP:OD1	1:C:515:LYS:HD3	2.16	0.46
1:B:348:ILE:O	1:B:472:SER:HA	2.16	0.46
3:D:3:DC:H2'	3:D:4:DA:C8	2.51	0.46
1:A:279:LYS:NZ	4:A:601:EDO:O1	2.49	0.45
1:C:350:THR:HB	1:C:356:PHE:CD2	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:350:THR:HB	1:B:356:PHE:CD2	2.51	0.45
1:B:143:ILE:HG22	1:B:145:LEU:HG	1.98	0.45
1:C:378:LYS:HB3	1:C:378:LYS:HE3	1.83	0.45
1:C:444:LEU:HD12	1:C:444:LEU:HA	1.81	0.45
1:B:317:ILE:HG22	1:B:319:VAL:HG22	1.97	0.45
3:F:13:DC:H2''	3:F:14:DA:C8	2.52	0.45
1:C:348:ILE:O	1:C:472:SER:HA	2.18	0.44
1:B:374:LYS:HB3	1:B:470:PHE:CD1	2.52	0.44
1:A:374:LYS:HB3	1:A:470:PHE:CD1	2.53	0.44
1:B:194:ALA:HB1	1:B:198:PHE:CD2	2.52	0.43
1:B:433:LYS:HB2	1:B:436:GLU:HG3	2.00	0.43
1:B:284:ASP:OD1	1:B:515:LYS:HD3	2.18	0.43
1:A:455:TYR:CZ	1:A:456:LYS:HD2	2.53	0.43
1:C:203:LYS:O	1:C:207:ILE:HG12	2.19	0.43
1:B:188:GLU:O	1:B:230:ASP:HB3	2.18	0.43
1:C:418:LYS:HE3	1:C:434:TRP:CE2	2.54	0.43
1:B:566:LYS:HE3	1:B:566:LYS:HB2	1.77	0.43
1:B:162:ILE:HG21	1:B:204:ILE:HG12	2.01	0.42
1:B:502:TYR:HA	1:B:505:TYR:HB3	2.02	0.42
1:A:60:ASP:HB3	1:A:63:CYS:HB3	2.01	0.42
1:B:532:ARG:HE	1:B:532:ARG:HB2	1.58	0.42
1:C:319:VAL:HB	1:C:323:ASP:HB2	2.01	0.42
1:B:464:ILE:HG13	1:B:494:VAL:HG11	2.00	0.42
1:A:203:LYS:O	1:A:207:ILE:HG12	2.19	0.42
1:C:338:LEU:HD23	1:C:338:LEU:HA	1.89	0.42
3:H:14:DA:H61	2:I:2:DT:H3	1.67	0.42
1:B:51:ILE:HD11	1:B:83:ASN:HB3	2.02	0.42
1:A:384:LYS:NZ	7:A:702:HOH:O	2.36	0.41
1:A:43:LYS:HD2	1:A:264:THR:HG21	2.02	0.41
1:C:114:ASP:O	1:C:147:CYS:HA	2.20	0.41
1:C:539:ILE:HG23	1:C:567:MET:HG3	2.01	0.41
1:A:326:PHE:HE1	1:A:577:ILE:HD12	1.86	0.41
1:A:357:ILE:C	1:A:358:LEU:HD12	2.46	0.41
1:C:244:ILE:HG12	1:C:263:LEU:CD1	2.51	0.41
1:C:376:TRP:CZ2	1:C:472:SER:HB2	2.55	0.41
1:A:317:ILE:HG22	1:A:319:VAL:HG22	2.03	0.41
1:B:418:LYS:HG3	1:B:434:TRP:CE3	2.55	0.41
1:A:572:MET:HG2	1:A:577:ILE:HD11	2.02	0.40
1:A:67:ASN:OD1	1:A:67:ASN:N	2.54	0.40
1:A:192:ASP:HA	2:E:13:DG:O3'	2.22	0.40
1:A:452:MET:HE2	1:A:475:VAL:HG11	2.02	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:330:ILE:HG12	1:B:575:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	540/577 (94%)	522 (97%)	18 (3%)	0	100	100
1	B	549/577 (95%)	529 (96%)	20 (4%)	0	100	100
1	C	533/577 (92%)	515 (97%)	18 (3%)	0	100	100
All	All	1622/1731 (94%)	1566 (96%)	56 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	457/547 (84%)	449 (98%)	8 (2%)	51	69
1	B	492/547 (90%)	482 (98%)	10 (2%)	48	67
1	C	452/547 (83%)	445 (98%)	7 (2%)	57	73
All	All	1401/1641 (85%)	1376 (98%)	25 (2%)	51	69

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	ILE
1	A	67	ASN
1	A	88	LYS
1	A	115	ILE
1	A	168	TYR
1	A	432	ARG
1	A	455	TYR
1	A	482	GLU
1	B	31	TYR
1	B	115	ILE
1	B	121	SER
1	B	143	ILE
1	B	168	TYR
1	B	263	LEU
1	B	292	THR
1	B	455	TYR
1	B	508	ILE
1	B	577	ILE
1	C	97	THR
1	C	115	ILE
1	C	168	TYR
1	C	296	LEU
1	C	363	VAL
1	C	444	LEU
1	C	455	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	282	ASN
1	B	83	ASN
1	B	107	ASN
1	B	282	ASN
1	B	438	GLN
1	B	541	ASN
1	B	545	GLN
1	C	144	ASN
1	C	282	ASN
1	C	406	ASN
1	C	438	GLN
1	C	469	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 19 ligands modelled in this entry, 9 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	YB0	A	602	-	42,45,45	0.18	0	59,60,60	0.64	0
4	EDO	B	606	-	3,3,3	0.52	0	2,2,2	0.16	0
4	EDO	B	601	-	3,3,3	0.43	0	2,2,2	0.24	0
4	EDO	B	602	-	3,3,3	0.51	0	2,2,2	0.28	0
4	EDO	F	101	-	3,3,3	0.48	0	2,2,2	0.23	0
5	YB0	C	601	-	42,45,45	0.16	0	59,60,60	0.86	2 (3%)
4	EDO	A	601	-	3,3,3	0.53	0	2,2,2	0.10	0
4	EDO	B	604	6	3,3,3	0.43	0	2,2,2	0.44	0
4	EDO	B	603	-	3,3,3	0.49	0	2,2,2	0.19	0
5	YB0	B	605	-	42,45,45	0.16	0	59,60,60	0.89	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	YB0	A	602	-	-	3/24/48/48	0/5/5/5
4	EDO	B	606	-	-	0/1/1/1	-
4	EDO	B	601	-	-	0/1/1/1	-
4	EDO	B	602	-	-	0/1/1/1	-
4	EDO	F	101	-	-	0/1/1/1	-
5	YB0	C	601	-	-	6/24/48/48	0/5/5/5
4	EDO	A	601	-	-	0/1/1/1	-
4	EDO	B	604	6	-	1/1/1/1	-
4	EDO	B	603	-	-	0/1/1/1	-
5	YB0	B	605	-	-	6/24/48/48	0/5/5/5

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	605	YB0	N10-C11-N12	3.90	124.75	119.28
5	C	601	YB0	C9-N10-C11	3.88	130.58	123.46
5	B	605	YB0	C9-N10-C11	3.84	130.50	123.46
5	C	601	YB0	N10-C11-N12	3.84	124.66	119.28
5	B	605	YB0	N12-C11-N20	-2.08	116.42	120.26

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	602	YB0	O21-C5-C6-S7
5	B	605	YB0	C2-C5-C6-S7
5	B	605	YB0	N12-C13-C14-C15
5	B	605	YB0	N12-C13-C14-C19
5	C	601	YB0	S7-C8-C9-N10
4	B	604	EDO	O1-C1-C2-O2
5	A	602	YB0	N30-C31-N32-C33
5	B	605	YB0	N30-C31-N32-C33
5	C	601	YB0	N30-C31-N32-C33
5	C	601	YB0	N12-C13-C14-C19
5	A	602	YB0	C26-C31-N32-C33
5	B	605	YB0	C26-C31-N32-C33
5	C	601	YB0	C26-C31-N32-C33
5	B	605	YB0	O21-C5-C6-S7
5	C	601	YB0	N12-C13-C14-C15
5	C	601	YB0	C3-C22-N23-C24


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	601	EDO	1	0
4	A	601	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	544/577 (94%)	-0.35	16 (2%) 53 51	26, 47, 84, 135	0
1	B	551/577 (95%)	-0.45	18 (3%) 49 47	25, 43, 81, 168	0
1	C	537/577 (93%)	-0.03	24 (4%) 38 37	33, 62, 115, 155	0
2	E	14/14 (100%)	-0.46	1 (7%) 22 21	35, 41, 159, 190	0
2	G	14/14 (100%)	-1.08	0 100 100	32, 38, 59, 60	0
2	I	14/14 (100%)	-0.46	0 100 100	41, 50, 146, 170	0
3	D	14/14 (100%)	-0.42	0 100 100	32, 40, 129, 153	0
3	F	14/14 (100%)	-0.88	0 100 100	29, 34, 73, 76	0
3	H	14/14 (100%)	-0.34	0 100 100	39, 50, 130, 139	0
All	All	1716/1815 (94%)	-0.29	59 (3%) 48 46	25, 49, 105, 190	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	156	ARG	5.0
1	A	139	SER	4.9
1	B	272	GLU	4.1
1	B	136	LEU	3.9
1	B	31	TYR	3.8
1	B	132	VAL	3.8
1	A	138	GLU	3.8
1	B	135	ASP	3.8
1	C	31	TYR	3.7
1	A	140	ASP	3.6
1	B	402	ASN	3.6
1	B	133	VAL	3.6
1	C	273	THR	3.6
1	C	287	ILE	3.5
1	A	272	GLU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	130	LYS	3.5
1	C	157	TYR	3.5
1	B	27	SER	3.5
1	A	401	ASP	3.5
1	A	30	TYR	3.4
1	C	94	ASN	3.3
1	B	138	GLU	3.3
1	B	139	SER	3.3
1	A	273	THR	3.2
1	A	31	TYR	3.2
1	C	142	LYS	3.2
1	C	272	GLU	3.0
1	C	143	ILE	2.9
1	C	96	TRP	2.9
1	A	285	ILE	2.9
1	A	555	ILE	2.9
1	B	140	ASP	2.9
1	B	134	ASN	2.8
1	A	132	VAL	2.8
1	C	95	TYR	2.8
1	C	554	SER	2.8
1	C	141	ILE	2.7
1	C	556	ASP	2.7
1	C	404	ASN	2.7
1	B	137	ASP	2.7
1	C	155	TRP	2.7
1	A	270	THR	2.6
1	A	554	SER	2.5
1	B	270	THR	2.5
1	B	405	THR	2.5
1	A	286	CYS	2.5
1	C	405	THR	2.5
1	C	298	LYS	2.5
1	C	85	TYR	2.4
1	B	273	THR	2.4
1	B	28	GLY	2.4
1	B	29	ILE	2.3
1	C	555	ILE	2.3
1	C	93	GLU	2.3
1	C	51	ILE	2.2
2	E	1	DA	2.2
1	A	404	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	73	TYR	2.0
1	C	98	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EDO	B	606	4/4	0.77	0.23	54,69,80,85	0
6	K	F	102	1/1	0.83	0.21	94,94,94,94	0
4	EDO	A	601	4/4	0.87	0.19	59,63,63,63	0
4	EDO	B	602	4/4	0.89	0.21	55,57,58,62	0
6	K	D	101	1/1	0.90	0.15	81,81,81,81	0
4	EDO	B	603	4/4	0.91	0.22	58,58,77,81	0
5	YB0	C	601	41/41	0.92	0.12	61,75,101,104	0
6	K	C	603	1/1	0.92	0.10	85,85,85,85	0
4	EDO	B	604	4/4	0.93	0.14	40,46,49,60	0
5	YB0	A	602	41/41	0.93	0.12	43,52,101,111	0
4	EDO	F	101	4/4	0.94	0.11	26,46,50,61	0
5	YB0	B	605	41/41	0.95	0.12	36,44,99,112	0
4	EDO	B	601	4/4	0.95	0.10	39,44,48,53	0
6	K	B	609	1/1	0.95	0.07	86,86,86,86	0
6	K	A	603	1/1	0.97	0.05	56,56,56,56	0
6	K	B	607	1/1	0.97	0.06	61,61,61,61	0
6	K	B	608	1/1	0.98	0.04	54,54,54,54	0
6	K	A	604	1/1	0.98	0.08	74,74,74,74	0
6	K	C	602	1/1	0.98	0.04	69,69,69,69	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around YB0 C 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around YB0 A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around YB0 B 605:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.