



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:10 AM UTC

PDB ID : 8FSS / pdb_00008fss
Title : Complex Structure of YejA-S481A with Microcin C7
Authors : Nair, S.K.; Dong, S.-H.
Deposited on : 2023-01-11
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	586	 3% 87% 9% ..
2	B	7	 14% 71% 14%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5275 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YejA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	571	Total	C	N	O	S	Se	0	2	0
			4677	3011	797	855	2	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	initiating methionine	UNP P33913
A	481	ALA	SER	engineered mutation	UNP P33913

- Molecule 2 is a protein called Microcin C7.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	7	Total	C	N	O	P	S	0	0	0
			80	42	18	18	1	1			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	7	7MD	ASN	modified residue	UNP Q47505


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	510	Total	O	0	0
			510	510		
3	B	8	Total	O	0	0
			8	8		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Yeja



• Molecule 2: Microcin C7



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	90.09Å 102.83Å 144.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	76.55 – 2.00 76.55 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.8 (76.55-2.00) 98.8 (76.55-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8	Depositor
R, R_{free}	0.219 , (Not available) 0.219 , 0.247	Depositor DCC
R_{free} test set	2243 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	24.0	Xtriage
Anisotropy	1.090	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5275	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 7MD, FME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	2/4811 (0.0%)	0.57	0/6530
2	B	5.49	14/34 (41.2%)	2.29	1/44 (2.3%)
All	All	0.66	16/4845 (0.3%)	0.60	1/6574 (0.0%)

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	5	ASN	C-N	11.79	1.49	1.33
2	B	4	GLY	C-N	10.38	1.47	1.33
2	B	4	GLY	CA-C	10.17	1.61	1.52
2	B	3	THR	N-CA	-9.30	1.33	1.46
2	B	2	ARG	NE-CZ	8.81	1.42	1.33
2	B	5	ASN	CG-ND2	8.52	1.51	1.33
2	B	3	THR	C-N	7.02	1.42	1.33
2	B	5	ASN	C-O	-6.84	1.15	1.23
2	B	3	THR	C-O	-6.67	1.15	1.24
2	B	2	ARG	CZ-NH2	6.53	1.42	1.33
2	B	3	THR	CA-C	6.52	1.61	1.52
2	B	2	ARG	CZ-NH1	-6.47	1.23	1.32
1	A	57	ASN	C-O	-6.14	1.18	1.24
2	B	4	GLY	C-O	-5.45	1.16	1.23
2	B	2	ARG	C-N	5.28	1.41	1.33
1	A	453	THR	C-O	-5.04	1.18	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	ARG	CG-CD-NE	-5.35	100.22	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4677	0	4523	31	0
2	B	80	0	68	12	0
3	A	510	0	0	13	2
3	B	8	0	0	2	0
All	All	5275	0	4591	40	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:7:7MD:O4'	2:B:7:7MD:C4'	1.66	1.20
1:A:17:ARG:NH1	3:A:601:HOH:O	2.01	0.91
2:B:7:7MD:O4'	3:B:201:HOH:O	1.94	0.84
2:B:1:FME:HCN	3:B:207:HOH:O	1.77	0.83
1:A:271:PHE:O	3:A:602:HOH:O	2.02	0.77
1:A:270:ASN:N	3:A:606:HOH:O	2.19	0.73
1:A:358:LYS:O	3:A:603:HOH:O	2.06	0.73
1:A:579:PRO:O	3:A:604:HOH:O	2.07	0.72
2:B:7:7MD:H8	2:B:7:7MD:O5'	1.94	0.67
1:A:413:LEU:HD13	1:A:440:ILE:HG12	1.78	0.66
2:B:7:7MD:C33	2:B:7:7MD:H5'A	2.26	0.66
1:A:281:GLN:O	1:A:282:LYS:HB2	1.97	0.64
1:A:275:TYR:N	3:A:611:HOH:O	2.30	0.63
1:A:447:VAL:HG23	1:A:452:ILE:HG12	1.80	0.63
1:A:116:ARG:HG3	3:A:667:HOH:O	1.99	0.62
1:A:467:VAL:H	2:B:7:7MD:H34	1.64	0.62
2:B:7:7MD:C8	2:B:7:7MD:C5'	2.83	0.56
1:A:419:LEU:HD11	1:A:428:VAL:HG21	1.91	0.53
1:A:359:LYS:HG2	3:A:790:HOH:O	2.08	0.52
2:B:7:7MD:H5'A	2:B:7:7MD:C32	2.41	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:VAL:HG22	1:A:13:LEU:HD13	1.95	0.48
1:A:72:THR:HA	1:A:79:SER:O	2.13	0.47
1:A:284:GLU:H	1:A:284:GLU:HG2	1.52	0.47
1:A:339:GLU:HG3	3:A:981:HOH:O	2.13	0.47
1:A:107:PHE:CE2	1:A:113:ILE:HG12	2.52	0.45
1:A:379:GLY:HA2	3:A:614:HOH:O	2.16	0.45
1:A:266:TYR:CG	1:A:278:LYS:HE3	2.52	0.45
1:A:292:LEU:HD13	1:A:538:MSE:SE	2.67	0.45
1:A:270:ASN:ND2	3:A:605:HOH:O	2.12	0.44
1:A:279:ASP:OD2	3:A:607:HOH:O	2.21	0.44
1:A:154:LEU:HG	1:A:162:MSE:HE2	2.00	0.44
2:B:7:7MD:H8	2:B:7:7MD:C5'	2.48	0.42
1:A:466:ARG:HA	2:B:7:7MD:C34	2.49	0.42
1:A:282:LYS:HD2	1:A:543:GLU:HG3	2.01	0.42
1:A:246:PHE:HB2	1:A:251:PHE:CZ	2.55	0.41
2:B:7:7MD:C4'	2:B:7:7MD:N9	2.75	0.41
1:A:466:ARG:HA	2:B:7:7MD:H34A	2.02	0.41
1:A:245:ALA:HB1	1:A:250:ALA:HB3	2.02	0.41
1:A:283:ASN:OD1	1:A:283:ASN:N	2.54	0.40
1:A:275:TYR:HB2	3:A:611:HOH:O	2.20	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:927:HOH:O	3:A:1026:HOH:O[4_556]	1.99	0.21
3:A:831:HOH:O	3:A:1050:HOH:O[6_654]	2.08	0.12

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	567/586 (97%)	550 (97%)	16 (3%)	1 (0%)	43	42
2	B	5/7 (71%)	5 (100%)	0	0	100	100
All	All	572/593 (96%)	555 (97%)	16 (3%)	1 (0%)	43	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	282	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/491 (100%)	467 (95%)	26 (5%)	20	18
2	B	3/3 (100%)	3 (100%)	0	100	100
All	All	496/494 (100%)	470 (95%)	26 (5%)	20	18

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	VAL
1	A	13	LEU
1	A	15	GLU
1	A	92	ASP
1	A	135	LEU
1	A	163	LEU
1	A	164	SER
1	A	187	THR
1	A	201	LYS
1	A	218	ASN
1	A	221	VAL
1	A	230	THR
1	A	240	ASN
1	A	282	LYS
1	A	283	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	284	GLU
1	A	292	LEU
1	A	339	GLU
1	A	386	LEU
1	A	391	LEU
1	A	400	LYS
1	A	413	LEU
1	A	423	SER
1	A	429	LEU
1	A	469	ARG
1	A	519	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	424	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FME	B	1	2	8,9,10	2.96	3 (37%)	8,9,11	1.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FME	B	1	2	-	2/7/9/11	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	FME	CA-N	-6.05	1.37	1.46
2	B	1	FME	CB-CA	3.93	1.61	1.53
2	B	1	FME	CN-N	3.15	1.43	1.33

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	FME	C-CA-CB-CG
2	B	1	FME	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	FME	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	559/586 (95%)	0.46	20 (3%)	46	45	16, 26, 42, 62	2 (0%)
2	B	5/7 (71%)	0.40	0	100	100	25, 27, 28, 33	0
All	All	564/593 (95%)	0.46	20 (3%)	47	46	16, 26, 41, 62	2 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	275	TYR	3.8
1	A	283	ASN	3.1
1	A	268	GLY	3.0
1	A	376	LYS	3.0
1	A	5	LYS	2.9
1	A	407	ALA	2.9
1	A	409	THR	2.8
1	A	277	ILE	2.6
1	A	270	ASN	2.5
1	A	271	PHE	2.5
1	A	281	GLN	2.3
1	A	408	THR	2.3
1	A	326	ASN	2.3
1	A	55	ARG	2.3
1	A	579	PRO	2.2
1	A	400	LYS	2.1
1	A	187	THR	2.1
1	A	469	ARG	2.1
1	A	285	SER	2.1
1	A	359	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FME	B	1	10/11	0.89	0.11	23,32,37,45	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.