



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 08:50 PM UTC

PDB ID : 5FTD / pdb_00005ftd
Title : Crystal structure of Pif1 helicase from Bacteroides apo form
Authors : Chen, W.-F.; Dai, Y.-X.; Duan, X.-L.; Liu, N.-N.; Shi, W.; Li, M.; Dou, S.-X.;
Li, N.; Dong, Y.-H.; Rety, S.; Xi, X.-G.
Deposited on : 2016-01-12
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7381 atoms, of which 3473 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TPR DOMAIN PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	429	6934	2224	3473	570	651	16	0	0	0

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	442	442	442	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TPR DOMAIN PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.78Å 70.75Å 103.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.56 – 1.70 48.56 – 1.70	Depositor EDS
% Data completeness (in resolution range)	87.1 (48.56-1.70) 87.1 (48.56-1.70)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 1.70Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.168 , 0.194 0.173 , 0.199	Depositor DCC
R_{free} test set	2357 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	21.7	Xtrriage
Anisotropy	0.609	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7381	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/3529	0.83	4/4767 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	191	GLU	CA-C-N	7.43	126.70	118.97
1	A	191	GLU	C-N-CA	7.43	126.70	118.97
1	A	146	LEU	CA-C-N	6.02	126.58	120.38
1	A	146	LEU	C-N-CA	6.02	126.58	120.38

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	189	GLN	Peptide
1	A	3	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3461	3473	3473	43	0
2	A	5	0	0	0	0
3	A	442	0	0	17	5
All	All	3908	3473	3473	43	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:318:ARG:NH2	3:A:2323:HOH:O	1.91	1.00
1:A:145:GLN:NE2	3:A:2184:HOH:O	2.04	0.89
1:A:18:GLN:NE2	3:A:2014:HOH:O	2.06	0.86
1:A:29:LYS:NZ	3:A:2038:HOH:O	2.02	0.78
1:A:9:GLU:OE2	3:A:2008:HOH:O	2.04	0.75
1:A:389:SER:OG	1:A:390:ARG:NH2	2.20	0.74
1:A:228:ASN:O	3:A:2246:HOH:O	2.05	0.73
1:A:146:LEU:O	3:A:2186:HOH:O	2.07	0.73
1:A:318:ARG:NH1	1:A:319:THR:O	2.22	0.72
1:A:224:SER:O	3:A:2243:HOH:O	2.11	0.68
1:A:390:ARG:NH2	3:A:2408:HOH:O	2.31	0.62
1:A:188:ARG:NH2	3:A:2210:HOH:O	2.13	0.59
1:A:185:LYS:O	1:A:187:PHE:N	2.35	0.59
1:A:7:THR:HG21	1:A:185:LYS:HE2	1.84	0.57
1:A:185:LYS:HG2	1:A:186:ILE:H	1.70	0.56
1:A:155:ARG:O	1:A:159:SER:OG	2.07	0.55
1:A:329:ASN:O	1:A:344:ILE:HD13	2.07	0.54
1:A:3:ASP:OD1	1:A:3:ASP:N	2.41	0.53
1:A:101:LEU:CD2	1:A:103:ILE:HG13	2.39	0.51
1:A:163:ASP:OD1	3:A:2194:HOH:O	2.20	0.47
1:A:411:GLU:OE2	3:A:2417:HOH:O	2.21	0.47
1:A:52:ALA:HA	1:A:65:VAL:O	2.16	0.45
1:A:418:GLU:HA	1:A:421:TYR:CE2	2.51	0.45
1:A:335:LYS:NZ	3:A:2369:HOH:O	2.50	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:185:LYS:HG2	1:A:186:ILE:N	2.30	0.44
1:A:108:SER:C	1:A:109:MET:HE2	2.43	0.44
1:A:344:ILE:HD13	1:A:344:ILE:H	1.82	0.44
1:A:77:PRO:HG3	1:A:154:GLU:HA	2.00	0.43
1:A:31:GLY:O	1:A:186:ILE:HA	2.18	0.43
1:A:270:PRO:HB2	3:A:2067:HOH:O	2.18	0.43
1:A:29:LYS:HE2	1:A:385:TYR:OH	2.19	0.42
1:A:151:LYS:N	1:A:151:LYS:HD2	2.34	0.42
1:A:430:LYS:HE3	1:A:431:ARG:HG3	2.01	0.42
1:A:49:CYS:SG	1:A:101:LEU:HD13	2.60	0.42
1:A:155:ARG:HG3	1:A:165:PHE:CZ	2.55	0.42
1:A:38:LEU:HD11	1:A:103:ILE:HG21	2.02	0.42
1:A:101:LEU:C	1:A:101:LEU:HD23	2.44	0.42
1:A:109:MET:HE3	1:A:145:GLN:HG2	2.01	0.41
1:A:224:SER:HB2	3:A:2243:HOH:O	2.20	0.41
1:A:180:ILE:HG21	1:A:421:TYR:CZ	2.55	0.41
1:A:229:GLU:HB3	3:A:2250:HOH:O	2.20	0.40
1:A:29:LYS:HE2	3:A:2406:HOH:O	2.20	0.40
1:A:180:ILE:HD12	1:A:180:ILE:N	2.37	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:2103:HOH:O	3:A:2224:HOH:O[4_545]	1.97	0.23
3:A:2351:HOH:O	3:A:2421:HOH:O[2_554]	2.01	0.19
3:A:2054:HOH:O	3:A:2235:HOH:O[1_655]	2.14	0.06
3:A:2053:HOH:O	3:A:2235:HOH:O[1_655]	2.15	0.05
3:A:2348:HOH:O	3:A:2430:HOH:O[2_554]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	427/433 (99%)	414 (97%)	10 (2%)	3 (1%)	18 7

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	ILE
1	A	188	ARG
1	A	190	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	385/390 (99%)	367 (95%)	18 (5%)	23 9

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	4	MET
1	A	17	ILE
1	A	84	LEU
1	A	185	LYS
1	A	189	GLN
1	A	190	THR
1	A	202	ARG
1	A	218	LYS
1	A	269	ILE
1	A	282	ARG
1	A	318	ARG
1	A	341	LYS
1	A	344	ILE
1	A	394	LEU
1	A	395	GLU
1	A	398	VAL
1	A	422	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	HIS
1	A	145	GLN
1	A	236	HIS
1	A	296	ASN
1	A	361	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	1432	-	4,4,4	1.09	0	6,6,6	0.70	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	429/433 (99%)	0.11	31 (7%) 21 23	16, 28, 59, 100	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	187	PHE	7.3
1	A	190	THR	6.0
1	A	148	PRO	5.8
1	A	149	VAL	5.8
1	A	186	ILE	5.5
1	A	192	PRO	3.4
1	A	147	PRO	3.4
1	A	318	ARG	3.3
1	A	150	THR	3.2
1	A	185	LYS	3.2
1	A	189	GLN	3.1
1	A	75	PHE	3.1
1	A	188	ARG	3.0
1	A	191	GLU	2.9
1	A	155	ARG	2.9
1	A	3	ASP	2.9
1	A	151	LYS	2.9
1	A	124	TRP	2.8
1	A	215	SER	2.7
1	A	165	PHE	2.7
1	A	152	LYS	2.6
1	A	86	ASN	2.4
1	A	146	LEU	2.4
1	A	158	LEU	2.4
1	A	154	GLU	2.2
1	A	290	SER	2.1
1	A	375	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	377	HIS	2.1
1	A	395	GLU	2.0
1	A	208	SER	2.0
1	A	87	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	A	1432	5/5	0.97	0.07	26,27,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.