



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 12:26 AM UTC

PDB ID : 6FTA / pdb_00006fta
Title : Crystal structure of human phosphodiesterase 4D2 catalytic domain with inhibitor NPD-3098
Authors : Singh, A.K.; Brown, D.G.
Deposited on : 2018-02-20
Resolution : 2.34 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	
1	B	364	
1	C	364	
1	D	364	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 11415 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	329	2667	1686	456	511	14	0	1	0
1	B	324	2622	1659	448	501	14	0	0	0
1	C	323	2613	1654	446	499	14	0	0	0
1	D	324	2628	1663	449	502	14	0	1	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	75	GLY	-	expression tag	UNP Q08499
A	76	SER	-	expression tag	UNP Q08499
A	77	HIS	-	expression tag	UNP Q08499
A	78	MET	-	expression tag	UNP Q08499
B	75	GLY	-	expression tag	UNP Q08499
B	76	SER	-	expression tag	UNP Q08499
B	77	HIS	-	expression tag	UNP Q08499
B	78	MET	-	expression tag	UNP Q08499
C	75	GLY	-	expression tag	UNP Q08499
C	76	SER	-	expression tag	UNP Q08499
C	77	HIS	-	expression tag	UNP Q08499
C	78	MET	-	expression tag	UNP Q08499
D	75	GLY	-	expression tag	UNP Q08499
D	76	SER	-	expression tag	UNP Q08499
D	77	HIS	-	expression tag	UNP Q08499
D	78	MET	-	expression tag	UNP Q08499


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		
2	D	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is (4 {a} {S},8 {a} {S})-4-[4-methoxy-3-[(2-methoxyphenyl)methoxy]phenyl]-2-[1-(3-nitroimidazo[1,2-b]pyridazin-6-yl)piperidin-4-yl]-4 {a},5,6,7,8,8 {a}-hexahydrophthalazin-1-one (CCD ID: E6N) (formula: C₃₄H₃₇N₇O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			47	34	7	6		
5	B	1	Total	C	N	O	0	0
			47	34	7	6		
5	C	1	Total	C	N	O	0	0
			47	34	7	6		
5	D	1	Total	C	N	O	0	0
			47	34	7	6		

- Molecule 6 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
6	A	1	Total 15	8	2	4	1	0	0
6	B	1	Total 15	8	2	4	1	0	0
6	B	1	Total 15	8	2	4	1	0	0
6	C	1	Total 15	8	2	4	1	0	0
6	D	1	Total 15	8	2	4	1	0	0

- Molecule 7 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	D	1	Total C O 7 4 3	0	0
7	D	1	Total C O 7 4 3	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	127	Total O 127 127	0	0
8	B	109	Total O 109 109	0	0
8	C	94	Total O 94 94	0	0
8	D	154	Total O 154 154	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4D




- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4D




- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4D



- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4D





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	98.99Å 110.76Å 161.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	80.67 – 2.34 80.67 – 2.34	Depositor EDS
% Data completeness (in resolution range)	100.0 (80.67-2.34) 100.0 (80.67-2.34)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.181 , 0.240 0.191 , 0.245	Depositor DCC
R_{free} test set	3788 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	46.7	Xtrriage
Anisotropy	0.265	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11415	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, ZN, EPE, EDO, MG, E6N

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.12	2/2721 (0.1%)	1.14	13/3696 (0.4%)
1	B	1.19	5/2676 (0.2%)	1.08	10/3636 (0.3%)
1	C	1.15	2/2667 (0.1%)	1.08	6/3624 (0.2%)
1	D	1.22	6/2685 (0.2%)	1.05	3/3648 (0.1%)
All	All	1.17	15/10749 (0.1%)	1.09	32/14604 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	351	GLY	C-N	13.40	1.52	1.33
1	C	337	MET	C-O	-7.06	1.15	1.24
1	D	388	VAL	C-O	-6.98	1.15	1.24
1	A	351	GLY	CA-C	-6.65	1.42	1.51
1	D	120	VAL	C-O	-6.19	1.17	1.24

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	351	GLY	CA-C-N	13.26	145.56	121.70
1	A	351	GLY	C-N-CA	13.26	145.56	121.70
1	A	348	ARG	N-CA-C	-9.70	98.85	112.13
1	D	376	ILE	CB-CA-C	-9.20	103.89	111.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	290	LYS	N-CA-C	7.72	120.98	108.63

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	351	GLY	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2667	0	2622	30	0
1	B	2622	0	2578	17	0
1	C	2613	0	2570	25	0
1	D	2628	0	2586	15	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	28	0	42	0	0
4	B	28	0	42	0	0
4	C	12	0	18	0	0
4	D	48	0	72	3	0
5	A	47	0	0	1	0
5	B	47	0	0	0	0
5	C	47	0	0	1	0
5	D	47	0	0	0	0
6	A	15	0	17	3	0
6	B	30	0	35	0	0
6	C	15	0	17	0	0
6	D	15	0	17	1	0
7	D	14	0	20	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	127	0	0	2	0
8	B	109	0	0	1	0
8	C	94	0	0	0	0
8	D	154	0	0	0	0
All	All	11415	0	10636	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 84 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:306:ARG:HH11	1:C:306:ARG:HG2	1.49	0.75
1:C:306:ARG:HG2	1:C:306:ARG:NH1	2.02	0.73
1:A:357:MET:HG3	1:A:368:SER:OG	1.91	0.69
1:A:331:GLN:HE22	6:A:511:EPE:H102	1.57	0.68
1:A:350:ARG:O	1:A:351:GLY:O	2.13	0.67

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	328/364 (90%)	307 (94%)	16 (5%)	5 (2%)	8 6
1	B	322/364 (88%)	316 (98%)	6 (2%)	0	100 100
1	C	321/364 (88%)	305 (95%)	9 (3%)	7 (2%)	5 3
1	D	323/364 (89%)	320 (99%)	2 (1%)	1 (0%)	36 41
All	All	1294/1456 (89%)	1248 (96%)	33 (3%)	13 (1%)	12 11

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	352	MET
1	A	354	ILE
1	C	360	LYS
1	C	361	HIS
1	C	363	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/331 (91%)	283 (94%)	18 (6%)	17	20
1	B	296/331 (89%)	284 (96%)	12 (4%)	27	35
1	C	295/331 (89%)	281 (95%)	14 (5%)	23	30
1	D	297/331 (90%)	288 (97%)	9 (3%)	36	47
All	All	1189/1324 (90%)	1136 (96%)	53 (4%)	24	32

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	360	LYS
1	C	286	MET
1	D	327	GLN
1	B	390	PRO
1	C	178	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	210	GLN
1	C	250	GLN
1	C	242	GLN
1	C	258	GLN
1	A	407	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 48 ligands modelled in this entry, 8 are monoatomic - leaving 40 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	EDO	D	504	-	3,3,3	0.42	0	2,2,2	0.51	0
5	E6N	D	514	-	49,53,53	1.19	4 (8%)	60,76,76	2.08	12 (20%)
6	EPE	B	512	-	15,15,15	2.75	1 (6%)	19,20,20	1.51	2 (10%)
4	EDO	C	507	-	3,3,3	0.63	0	2,2,2	0.16	0
6	EPE	B	511	-	15,15,15	1.91	1 (6%)	19,20,20	1.34	2 (10%)
5	E6N	B	508	-	49,53,53	1.15	4 (8%)	60,76,76	2.05	13 (21%)
5	E6N	C	506	-	49,53,53	1.27	4 (8%)	60,76,76	1.89	8 (13%)
4	EDO	D	507	-	3,3,3	0.78	0	2,2,2	0.16	0
4	EDO	C	503	-	3,3,3	0.58	0	2,2,2	0.20	0
4	EDO	C	504	-	3,3,3	0.61	0	2,2,2	0.47	0
4	EDO	B	510	-	3,3,3	0.48	0	2,2,2	0.34	0
4	EDO	A	505	-	3,3,3	0.34	0	2,2,2	0.46	0
4	EDO	A	504	-	3,3,3	0.52	0	2,2,2	0.30	0
4	EDO	A	503	-	3,3,3	0.42	0	2,2,2	0.36	0
4	EDO	A	506	-	3,3,3	0.35	0	2,2,2	0.46	0
4	EDO	D	515	-	3,3,3	0.71	0	2,2,2	0.17	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	507	-	3,3,3	0.50	0	2,2,2	0.35	0
4	EDO	D	506	-	3,3,3	0.26	0	2,2,2	0.57	0
4	EDO	D	505	-	3,3,3	0.39	0	2,2,2	0.37	0
4	EDO	B	509	-	3,3,3	0.38	0	2,2,2	0.44	0
4	EDO	D	501	-	3,3,3	0.40	0	2,2,2	0.69	0
5	E6N	A	510	-	49,53,53	1.18	4 (8%)	60,76,76	2.06	13 (21%)
7	PEG	D	513	-	6,6,6	0.27	0	5,5,5	0.82	0
4	EDO	A	509	-	3,3,3	0.55	0	2,2,2	0.09	0
4	EDO	B	507	-	3,3,3	0.45	0	2,2,2	0.31	0
4	EDO	D	517	-	3,3,3	0.55	0	2,2,2	0.21	0
4	EDO	D	511	-	3,3,3	0.39	0	2,2,2	0.11	0
4	EDO	A	508	-	3,3,3	0.46	0	2,2,2	0.41	0
4	EDO	D	509	-	3,3,3	0.30	0	2,2,2	0.56	0
4	EDO	D	516	-	3,3,3	0.38	0	2,2,2	0.51	0
4	EDO	D	508	-	3,3,3	0.65	0	2,2,2	0.22	0
6	EPE	C	505	-	15,15,15	2.91	1 (6%)	19,20,20	5.70	8 (42%)
7	PEG	D	512	-	6,6,6	0.33	0	5,5,5	0.58	0
4	EDO	B	505	-	3,3,3	0.33	0	2,2,2	0.60	0
4	EDO	B	503	-	3,3,3	0.47	0	2,2,2	0.11	0
4	EDO	B	504	-	3,3,3	0.38	0	2,2,2	0.47	0
4	EDO	B	506	-	3,3,3	0.72	0	2,2,2	0.59	0
6	EPE	A	511	-	15,15,15	2.86	1 (6%)	19,20,20	1.62	5 (26%)
6	EPE	D	518	-	15,15,15	2.85	1 (6%)	19,20,20	1.57	5 (26%)
4	EDO	D	510	-	3,3,3	0.42	0	2,2,2	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	D	504	-	-	0/1/1/1	-
5	E6N	D	514	-	-	6/23/62/62	0/7/7/7
6	EPE	B	512	-	-	5/9/19/19	0/1/1/1
4	EDO	C	507	-	-	1/1/1/1	-
6	EPE	B	511	-	-	4/9/19/19	0/1/1/1
5	E6N	B	508	-	-	7/23/62/62	0/7/7/7
5	E6N	C	506	-	-	7/23/62/62	0/7/7/7
4	EDO	D	507	-	-	1/1/1/1	-
4	EDO	C	503	-	-	1/1/1/1	-
4	EDO	C	504	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	510	-	-	0/1/1/1	-
4	EDO	A	505	-	-	1/1/1/1	-
4	EDO	A	504	-	-	1/1/1/1	-
4	EDO	A	503	-	-	1/1/1/1	-
4	EDO	A	506	-	-	1/1/1/1	-
4	EDO	D	515	-	-	1/1/1/1	-
4	EDO	A	507	-	-	1/1/1/1	-
4	EDO	D	506	-	-	1/1/1/1	-
4	EDO	D	505	-	-	1/1/1/1	-
4	EDO	B	509	-	-	0/1/1/1	-
4	EDO	D	501	-	-	0/1/1/1	-
5	E6N	A	510	-	-	7/23/62/62	0/7/7/7
7	PEG	D	513	-	-	0/4/4/4	-
4	EDO	A	509	-	-	1/1/1/1	-
4	EDO	B	507	-	-	1/1/1/1	-
4	EDO	D	517	-	-	1/1/1/1	-
4	EDO	D	511	-	-	0/1/1/1	-
4	EDO	A	508	-	-	1/1/1/1	-
4	EDO	D	509	-	-	1/1/1/1	-
4	EDO	D	516	-	-	1/1/1/1	-
4	EDO	D	508	-	-	1/1/1/1	-
6	EPE	C	505	-	-	2/9/19/19	0/1/1/1
7	PEG	D	512	-	-	3/4/4/4	-
4	EDO	B	505	-	-	1/1/1/1	-
4	EDO	B	503	-	-	1/1/1/1	-
4	EDO	B	504	-	-	1/1/1/1	-
4	EDO	B	506	-	-	1/1/1/1	-
6	EPE	A	511	-	-	7/9/19/19	0/1/1/1
6	EPE	D	518	-	-	5/9/19/19	0/1/1/1
4	EDO	D	510	-	-	1/1/1/1	-

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	505	EPE	C10-S	-10.98	1.62	1.77
6	A	511	EPE	C10-S	-10.94	1.62	1.77
6	D	518	EPE	C10-S	-10.63	1.62	1.77
6	B	512	EPE	C10-S	-10.43	1.63	1.77
6	B	511	EPE	C10-S	-6.78	1.68	1.77

The worst 5 of 68 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	505	EPE	O1S-S-C10	-19.25	77.64	106.73
6	C	505	EPE	O2S-S-C10	-12.82	87.36	106.73
5	C	506	E6N	C29-C30-C25	-7.46	96.19	110.83
5	D	514	E6N	C29-C30-C25	-7.31	96.49	110.83
6	C	505	EPE	O3S-S-C10	-6.45	93.39	106.00

There are no chirality outliers.

5 of 77 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	A	510	E6N	C20-C21-N5-O3
5	A	510	E6N	N6-C21-N5-O3
5	C	506	E6N	C20-C21-N5-O3
5	D	514	E6N	N7-C16-N3-C15
6	A	511	EPE	C9-C10-S-O1S



There are no ring outliers.

7 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	506	E6N	1	0
4	D	506	EDO	3	0
5	A	510	E6N	1	0
7	D	513	PEG	1	0
7	D	512	PEG	2	0
6	A	511	EPE	3	0
6	D	518	EPE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	329/364 (90%)	0.44	21 (6%) 25 30	22, 48, 87, 137	14 (4%)
1	B	324/364 (89%)	0.13	8 (2%) 58 64	30, 50, 82, 130	0
1	C	323/364 (88%)	0.55	16 (4%) 34 40	19, 51, 89, 133	11 (3%)
1	D	324/364 (89%)	-0.09	7 (2%) 62 67	25, 42, 79, 144	1 (0%)
All	All	1300/1456 (89%)	0.26	52 (4%) 42 48	19, 48, 86, 144	26 (2%)

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	360	LYS	25.7
1	A	354	ILE	15.7
1	C	354	ILE	15.6
1	C	358	CYS	15.3
1	C	356	PRO	12.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EDO	D	515	4/4	0.76	0.19	62,68,70,74	0
4	EDO	D	507	4/4	0.77	0.28	71,81,81,85	0
4	EDO	B	510	4/4	0.81	0.17	75,76,76,76	0
4	EDO	C	504	4/4	0.81	0.24	78,78,88,89	0
4	EDO	C	507	4/4	0.83	0.22	68,80,83,84	0
4	EDO	D	506	4/4	0.84	0.18	53,60,64,76	0
4	EDO	B	509	4/4	0.85	0.14	66,67,68,69	0
7	PEG	D	513	7/7	0.85	0.18	54,58,66,71	0
4	EDO	B	504	4/4	0.86	0.21	77,79,79,81	0
4	EDO	D	508	4/4	0.87	0.17	51,59,69,75	0
4	EDO	D	501	4/4	0.88	0.17	57,59,61,69	0
4	EDO	B	505	4/4	0.88	0.20	66,68,72,76	0
5	E6N	A	510	47/47	0.88	0.16	58,87,139,150	0
5	E6N	C	506	47/47	0.88	0.15	48,67,111,125	0
6	EPE	A	511	15/15	0.88	0.16	71,102,120,122	0
6	EPE	B	511	15/15	0.88	0.18	76,99,110,114	0
4	EDO	B	506	4/4	0.88	0.17	62,70,79,80	0
4	EDO	A	507	4/4	0.89	0.20	77,81,84,87	0
5	E6N	D	514	47/47	0.89	0.14	46,69,126,132	0
4	EDO	A	509	4/4	0.89	0.18	54,60,66,69	0
4	EDO	D	516	4/4	0.89	0.19	69,73,75,76	0
6	EPE	B	512	15/15	0.89	0.17	63,100,115,117	0
4	EDO	A	506	4/4	0.89	0.15	51,52,55,64	0
4	EDO	B	507	4/4	0.90	0.22	74,80,93,100	0
4	EDO	A	504	4/4	0.90	0.16	52,66,69,70	0
4	EDO	A	508	4/4	0.90	0.16	76,76,78,80	0
5	E6N	B	508	47/47	0.90	0.13	46,62,112,117	0
4	EDO	D	510	4/4	0.90	0.16	66,74,84,92	0
7	PEG	D	512	7/7	0.91	0.12	53,54,61,62	0
6	EPE	D	518	15/15	0.91	0.18	46,109,128,132	0
4	EDO	D	509	4/4	0.92	0.14	67,68,73,77	0
4	EDO	C	503	4/4	0.92	0.14	63,64,71,75	0
4	EDO	D	511	4/4	0.92	0.12	53,54,57,57	0
4	EDO	A	503	4/4	0.92	0.14	77,78,82,85	0
4	EDO	D	504	4/4	0.92	0.12	76,77,80,82	0
4	EDO	B	503	4/4	0.93	0.15	67,68,69,74	0
4	EDO	D	517	4/4	0.95	0.10	51,54,56,57	0
4	EDO	A	505	4/4	0.95	0.08	51,52,53,54	0
6	EPE	C	505	15/15	0.95	0.11	53,75,90,95	0
4	EDO	D	505	4/4	0.96	0.08	49,52,52,54	0
2	ZN	A	501	1/1	0.99	0.02	43,43,43,43	0
3	MG	C	502	1/1	1.00	0.01	23,23,23,23	0
3	MG	D	503	1/1	1.00	0.01	22,22,22,22	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	B	501	1/1	1.00	0.01	44,44,44,44	0
2	ZN	C	501	1/1	1.00	0.01	42,42,42,42	0
2	ZN	D	502	1/1	1.00	0.01	38,38,38,38	0
3	MG	A	502	1/1	1.00	0.01	29,29,29,29	0
3	MG	B	502	1/1	1.00	0.03	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around E6N C 506:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around E6N D 514:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.