



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 09:35 PM EDT

PDB ID : 7FT6 / pdb_00007ft6
Title : SDCBP PanDDA analysis group deposition – The PDZ domains of SDCBP in complex with POB0008
Authors : Bradshaw, W.J.; Katis, V.L.; Bountra, C.; von Delft, F.; Brennan, P.E.
Deposited on : 2023-01-24
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	195	
1	B	195	
1	C	195	
1	D	195	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	SO4	D	305	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6411 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Syntenin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	S	0	1	0
			1481	935	262	275	9			
1	B	193	Total	C	N	O	S	0	1	0
			1495	943	264	279	9			
1	C	193	Total	C	N	O	S	0	3	0
			1514	953	270	282	9			
1	D	191	Total	C	N	O	S	0	1	0
			1481	935	262	275	9			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	104	SER	-	expression tag	UNP O00560
A	105	MET	-	expression tag	UNP O00560
B	104	SER	-	expression tag	UNP O00560
B	105	MET	-	expression tag	UNP O00560
C	104	SER	-	expression tag	UNP O00560
C	105	MET	-	expression tag	UNP O00560
D	104	SER	-	expression tag	UNP O00560
D	105	MET	-	expression tag	UNP O00560

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is D-GLUTAMIC ACID (CCD ID: DGL) (formula: $C_5H_9NO_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			10	5	1	4		
3	D	1	Total	C	N	O	0	0
			10	5	1	4		

- Molecule 4 is GLYCINE (CCD ID: GLY) (formula: $C_2H_5NO_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	N	O	0	0
			5	2	1	2		
4	D	1	Total	C	N	O	0	0
			5	2	1	2		

- Molecule 5 is methyl 1-(methanesulfonyl)-2-methyl-D-prolinate (CCD ID: YEK) (formula: $C_8H_{15}NO_4S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	D	1	Total	C	N	O	S	0	0
			14	8	1	4	1		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O_4S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is ALANINE (CCD ID: ALA) (formula: $C_3H_7NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	D	1	Total	C	N	O	0	0
			6	3	1	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	49	Total	O	0	0
			49	49		
8	B	105	Total	O	0	1
			105	105		
8	C	79	Total	O	0	0
			79	79		
8	D	110	Total	O	0	1
			110	110		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Syntenin-1




• Molecule 1: Syntenin-1




• Molecule 1: Syntenin-1



• Molecule 1: Syntenin-1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.69Å 49.42Å 114.67Å 90.00° 95.15° 90.00°	Depositor
Resolution (Å)	80.32 – 1.85 80.32 – 1.85	Depositor EDS
% Data completeness (in resolution range)	93.9 (80.32-1.85) 93.9 (80.32-1.85)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.20 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.215 , 0.262 0.225 , 0.262	Depositor DCC
R_{free} test set	3795 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	35.2	Xtriage
Anisotropy	0.229	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6411	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.02 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.8462e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DGL, EDO, YEK, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	0/1502	1.42	2/2019 (0.1%)
1	B	1.16	2/1516 (0.1%)	1.43	4/2038 (0.2%)
1	C	1.18	0/1535	1.45	2/2063 (0.1%)
1	D	1.24	3/1502 (0.2%)	1.49	7/2019 (0.3%)
All	All	1.18	5/6055 (0.1%)	1.45	15/8139 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	292	ASP	C-O	-6.75	1.15	1.23
1	D	238	ILE	C-O	-6.65	1.17	1.24
1	D	251	ASP	C-O	5.74	1.30	1.24
1	D	190	ILE	C-O	5.51	1.30	1.24
1	B	198	THR	C-O	-5.23	1.17	1.23

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	182	PHE	CB-CA-C	7.89	121.52	110.16
1	B	294	THR	CB-CA-C	-6.85	97.52	109.64
1	A	266	THR	CA-CB-OG1	-6.81	99.38	109.60
1	D	244	GLN	CA-C-N	6.25	130.27	121.02
1	D	244	GLN	C-N-CA	6.25	130.27	121.02
1	A	244	GLN	CB-CG-CD	-5.73	102.86	112.60
1	B	194	PRO	CB-CA-C	-5.64	103.84	111.85
1	D	153	ARG	CG-CD-NE	-5.51	99.88	112.00
1	D	254	ILE	N-CA-C	-5.50	105.14	110.42
1	B	295	ILE	CB-CA-C	5.48	116.04	110.94
1	D	182	PHE	CA-CB-CG	-5.33	108.47	113.80
1	C	278	ILE	CA-C-O	-5.32	114.88	120.57

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	198	THR	CA-CB-OG1	-5.28	101.69	109.60
1	C	200	THR	CA-CB-OG1	-5.15	101.88	109.60
1	D	170	SER	CA-C-O	-5.06	116.00	121.87

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1481	0	1530	16	0
1	B	1495	0	1541	9	0
1	C	1514	0	1558	14	0
1	D	1481	0	1530	23	0
2	A	4	0	6	0	0
2	B	8	0	12	0	0
2	C	12	0	18	0	0
2	D	8	0	12	0	0
3	B	10	0	7	0	0
3	D	10	0	7	0	0
4	C	5	0	2	0	0
4	D	5	0	2	1	0
5	D	14	0	0	0	0
6	D	15	0	0	0	0
7	D	6	0	4	0	0
8	A	49	0	0	1	0
8	B	105	0	0	1	0
8	C	79	0	0	1	0
8	D	110	0	0	4	0
All	All	6411	0	6229	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:108:ILE:HD12	1:C:198:THR:HG23	1.65	0.77
1:A:201:MET:HE1	1:A:267:ILE:HD12	1.67	0.76
1:A:292:ASP:OD1	1:A:294:THR:OG1	2.10	0.70
1:D:118:CYS:SG	1:D:184:GLU:O	2.45	0.67
1:A:286:ILE:HG23	1:A:290:LEU:HD12	1.80	0.64
1:D:117:LEU:HD12	1:D:117:LEU:C	2.23	0.64
1:D:197:ARG:NH1	8:D:403:HOH:O	2.25	0.60
4:D:309:GLY:OXT	8:D:402:HOH:O	2.16	0.60
1:B:177:VAL:O	8:B:401:HOH:O	2.15	0.60
1:C:108:ILE:HG22	1:C:108:ILE:O	2.03	0.59
1:D:206:THR:HG1	1:D:208:HIS:HD1	1.49	0.58
1:D:154:PHE:CD1	1:D:247:ILE:CD1	2.87	0.57
1:C:261:SER:OG	1:C:262:GLY:N	2.38	0.57
1:B:286:ILE:HG23	1:B:290:LEU:HD12	1.86	0.56
1:D:117:LEU:HD12	1:D:117:LEU:O	2.07	0.55
1:A:242:ASN:ND2	1:A:261:SER:OG	2.40	0.54
1:D:197:ARG:NH2	8:D:401:HOH:O	2.11	0.54
1:B:201:MET:HE1	1:B:267:ILE:HD12	1.88	0.54
1:C:140:LEU:HD12	1:C:291:MET:HE2	1.90	0.53
1:B:109:LYS:N	1:B:192:ASP:OD2	2.41	0.53
1:D:117:LEU:C	1:D:117:LEU:CD1	2.82	0.52
1:B:112:ILE:CG2	1:B:189:THR:HG22	2.39	0.52
1:A:290:LEU:HD13	1:B:182:PHE:HB2	1.93	0.49
1:A:114:GLU:OE1	1:A:187:THR:HB	2.13	0.49
1:D:248:GLY:HA2	1:D:287:MET:HE3	1.94	0.49
1:D:210:GLY:HA3	1:D:225:SER:HB2	1.96	0.47
1:A:244:GLN:HE21	1:A:257:ILE:HD13	1.79	0.47
1:D:210:GLY:HA3	1:D:225:SER:CB	2.45	0.47
1:C:160:GLN:HA	1:C:164:GLU:O	2.15	0.47
1:D:142:GLN:HA	1:D:290:LEU:O	2.14	0.47
1:C:210:GLY:HA3	1:C:225:SER:HB2	1.98	0.46
1:A:183:GLY:O	1:A:184:GLU:C	2.59	0.46
1:C:196:GLU:O	1:C:197[B]:ARG:HD3	2.16	0.46
1:C:248:GLY:O	1:C:288:LYS:HG2	2.15	0.46
1:D:201:MET:HE1	1:D:267:ILE:HD12	1.98	0.46
1:A:197:ARG:HD3	1:C:194:PRO:O	2.16	0.45
1:D:153:ARG:NH1	8:D:405:HOH:O	2.37	0.45
1:A:282:MET:HE3	1:A:287:MET:HG3	1.98	0.44
1:D:193:ARG:N	1:D:194:PRO:HD3	2.32	0.44
1:D:154:PHE:CE1	1:D:247:ILE:HD13	2.53	0.44
1:C:182:PHE:HA	8:C:440:HOH:O	2.18	0.43
1:D:137:PHE:CE2	1:D:195:PHE:CE2	3.06	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:127:LEU:HD23	1:C:127:LEU:HA	1.83	0.43
1:A:128:ARG:HB2	1:A:140:LEU:HB2	2.01	0.42
1:C:108:ILE:CD1	1:C:198:THR:HG23	2.42	0.42
1:B:106:ALA:HB3	1:B:266:THR:HG21	2.01	0.42
1:D:280:LYS:O	1:D:281:ARG:HB2	2.20	0.41
1:D:286:ILE:HG23	1:D:290:LEU:HD12	2.01	0.41
1:D:137:PHE:CZ	1:D:195:PHE:CE2	3.07	0.41
1:A:298:VAL:OXT	1:A:298:VAL:CG2	2.68	0.41
1:C:280:LYS:O	1:C:281:ARG:HB2	2.21	0.41
1:D:204:ASP:OD1	1:D:204:ASP:C	2.64	0.41
1:A:283:ALA:HB3	1:A:286:ILE:HD12	2.03	0.41
1:B:160:GLN:HA	1:B:164:GLU:O	2.20	0.41
1:A:267:ILE:HG22	1:A:269:ILE:HG23	2.03	0.41
1:C:128:ARG:HD3	1:C:140:LEU:HD22	2.03	0.41
1:D:198:THR:HA	1:D:267:ILE:O	2.21	0.40
1:B:198:THR:HG22	1:B:268:THR:OG1	2.22	0.40
1:A:118:CYS:N	8:A:410:HOH:O	2.53	0.40
1:A:158:VAL:CG1	1:A:188:MET:HE2	2.52	0.40
1:D:285:SER:O	1:D:289:SER:HB3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	190/195 (97%)	181 (95%)	6 (3%)	3 (2%)	7	2
1	B	192/195 (98%)	187 (97%)	5 (3%)	0	100	100
1	C	194/195 (100%)	187 (96%)	7 (4%)	0	100	100
1	D	190/195 (97%)	183 (96%)	7 (4%)	0	100	100
All	All	766/780 (98%)	738 (96%)	25 (3%)	3 (0%)	30	17

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	184	GLU
1	A	109	LYS
1	A	183	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/168 (99%)	158 (95%)	8 (5%)	23	8
1	B	167/168 (99%)	162 (97%)	5 (3%)	36	21
1	C	169/168 (101%)	163 (96%)	6 (4%)	31	16
1	D	166/168 (99%)	161 (97%)	5 (3%)	36	21
All	All	668/672 (99%)	644 (96%)	24 (4%)	31	16

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	114	GLU
1	A	140	LEU
1	A	171	SER
1	A	199	ILE
1	A	205	SER
1	A	287	MET
1	A	295	ILE
1	A	298	VAL
1	B	114	GLU
1	B	144	ASN
1	B	179	LYS
1	B	220	SER
1	B	266	THR
1	C	107	GLU
1	C	108	ILE
1	C	110	GLN
1	C	128	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	150	VAL
1	C	288	LYS
1	D	108	ILE
1	D	117	LEU
1	D	125	ILE
1	D	223	LYS
1	D	289	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	GLN
1	A	230	ASN
1	A	237	ASN
1	A	244	GLN
1	B	142	GLN
1	B	160	GLN
1	B	165	ASN
1	B	202	HIS
1	B	208	HIS
1	B	215	ASN
1	C	230	ASN
1	D	110	GLN
1	D	215	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	C	303	-	3,3,3	0.36	0	2,2,2	0.49	0
2	EDO	D	303	-	3,3,3	0.16	0	2,2,2	0.25	0
2	EDO	C	301	-	3,3,3	0.14	0	2,2,2	0.14	0
2	EDO	D	302	-	3,3,3	0.23	0	2,2,2	0.19	0
6	SO4	D	305	-	4,4,4	0.35	0	6,6,6	0.07	0
2	EDO	C	302	-	3,3,3	0.19	0	2,2,2	0.37	0
4	GLY	D	309	-	4,4,4	1.22	1 (25%)	3,4,4	1.25	0
6	SO4	D	304	-	4,4,4	0.35	0	6,6,6	0.09	0
2	EDO	A	301	-	3,3,3	0.19	0	2,2,2	0.12	0
6	SO4	D	306	-	4,4,4	0.34	0	6,6,6	0.07	0
2	EDO	B	301	-	3,3,3	0.33	0	2,2,2	0.20	0
5	YEK	D	301	-	13,14,14	0.36	0	16,22,22	0.63	0
7	ALA	D	307	-	5,5,5	1.11	0	6,6,6	0.77	0
2	EDO	B	302	-	3,3,3	0.41	0	2,2,2	0.27	0
3	DGL	B	303	-	8,9,9	1.02	0	8,11,11	1.55	1 (12%)
3	DGL	D	308	-	8,9,9	1.19	0	8,11,11	0.92	0
4	GLY	C	304	-	4,4,4	1.01	0	3,4,4	1.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	C	303	-	-	1/1/1/1	-
2	EDO	D	303	-	-	1/1/1/1	-
2	EDO	C	301	-	-	1/1/1/1	-
2	EDO	D	302	-	-	1/1/1/1	-
2	EDO	C	302	-	-	1/1/1/1	-
4	GLY	D	309	-	-	0/2/2/2	-
2	EDO	A	301	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	301	-	-	1/1/1/1	-
5	YEK	D	301	-	-	5/14/27/27	0/1/1/1
7	ALA	D	307	-	-	0/4/4/4	-
2	EDO	B	302	-	-	0/1/1/1	-
3	DGL	B	303	-	-	4/9/9/9	-
3	DGL	D	308	-	-	2/9/9/9	-
4	GLY	C	304	-	-	0/2/2/2	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	309	GLY	OXT-C	-2.35	1.23	1.30

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	303	DGL	OXT-C-O	-2.86	117.58	124.08

There are no chirality outliers.

All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	D	301	YEK	C5-N-S-C4
5	D	301	YEK	C5-N-S-O2
5	D	301	YEK	C5-N-S-O3
5	D	301	YEK	C1-N-S-C4
5	D	301	YEK	C1-N-S-O3
2	C	303	EDO	O1-C1-C2-O2
3	B	303	DGL	C-CA-CB-CG
3	B	303	DGL	N-CA-CB-CG
3	D	308	DGL	O-C-CA-CB
3	D	308	DGL	OXT-C-CA-CB
2	B	301	EDO	O1-C1-C2-O2
2	C	301	EDO	O1-C1-C2-O2
2	C	302	EDO	O1-C1-C2-O2
2	D	302	EDO	O1-C1-C2-O2
2	D	303	EDO	O1-C1-C2-O2
3	B	303	DGL	OE2-CD-CG-CB
3	B	303	DGL	OE1-CD-CG-CB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	309	GLY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/195 (97%)	1.26	25 (13%) 7 7	12, 51, 78, 89	13 (6%)
1	B	193/195 (98%)	0.70	17 (8%) 15 15	11, 41, 58, 67	13 (6%)
1	C	193/195 (98%)	0.72	18 (9%) 14 14	11, 44, 66, 94	6 (3%)
1	D	191/195 (97%)	0.63	20 (10%) 11 11	10, 36, 52, 85	16 (8%)
All	All	768/780 (98%)	0.83	80 (10%) 11 11	10, 42, 69, 94	48 (6%)

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	111	GLY	14.1
1	A	112	ILE	13.2
1	A	108	ILE	12.7
1	B	117	LEU	11.8
1	B	118	CYS	11.7
1	D	187	THR	10.6
1	A	109	LYS	10.5
1	A	182	PHE	10.4
1	B	108	ILE	9.7
1	A	183	GLY	9.6
1	C	106	ALA	9.2
1	A	181	ALA	8.7
1	A	273[A]	PHE	8.6
1	B	106	ALA	8.5
1	B	119	LYS	8.2
1	D	108	ILE	8.1
1	D	112	ILE	8.0
1	D	273[A]	PHE	7.5
1	C	108	ILE	7.5
1	A	125	ILE	7.1
1	B	112	ILE	6.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	109	LYS	6.8
1	B	109	LYS	6.6
1	D	111	GLY	6.2
1	C	273[A]	PHE	6.1
1	B	273[A]	PHE	5.9
1	B	111	GLY	5.7
1	A	185	LYS	5.4
1	A	118	CYS	5.2
1	D	267	ILE	5.0
1	D	113	ARG	4.9
1	B	107	GLU	4.8
1	C	197[A]	ARG	4.8
1	D	117	LEU	4.7
1	D	182	PHE	4.2
1	A	184	GLU	4.1
1	C	144[A]	ASN	4.0
1	D	149	LEU	3.9
1	B	113	ARG	3.9
1	C	182	PHE	3.9
1	A	286	ILE	3.8
1	D	268	THR	3.7
1	D	121	GLN	3.7
1	D	256	ASP	3.6
1	A	282	MET	3.6
1	C	209	VAL	3.5
1	D	110	GLN	3.5
1	A	290	LEU	3.5
1	B	114	GLU	3.4
1	A	110	GLN	3.4
1	B	110	GLN	3.3
1	A	283	ALA	3.3
1	C	107	GLU	3.2
1	A	287	MET	3.1
1	D	118	CYS	3.1
1	D	148	SER	2.9
1	C	110	GLN	2.9
1	C	109	LYS	2.9
1	C	111	GLY	2.9
1	B	123	GLY	2.8
1	D	144	ASN	2.7
1	A	298	VAL	2.7
1	B	115	VAL	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	183	GLY	2.6
1	A	209	VAL	2.5
1	C	121	GLN	2.4
1	A	186	ILE	2.3
1	A	117	LEU	2.3
1	C	281	ARG	2.2
1	A	264	VAL	2.2
1	C	263	THR	2.2
1	C	151	GLY	2.2
1	B	182	PHE	2.2
1	D	184	GLU	2.1
1	D	125	ILE	2.1
1	B	177	VAL	2.1
1	A	179	LYS	2.1
1	A	263	THR	2.1
1	C	116	ILE	2.0
1	C	206	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	SO4	D	305	5/5	0.35	0.42	254,261,261,264	5
3	DGL	D	308	10/10	0.72	0.18	40,66,77,84	0
2	EDO	C	303	4/4	0.82	0.23	62,65,67,67	0
6	SO4	D	306	5/5	0.83	0.47	71,73,75,75	5
3	DGL	B	303	10/10	0.85	0.12	44,56,66,67	0
5	YEK	D	301	14/14	0.86	0.18	47,50,51,52	14

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	C	301	4/4	0.88	0.12	34,34,34,40	0
7	ALA	D	307	6/6	0.88	0.15	49,58,65,67	0
2	EDO	B	301	4/4	0.90	0.11	36,37,38,46	0
4	GLY	C	304	5/5	0.91	0.09	47,52,54,56	0
4	GLY	D	309	5/5	0.92	0.13	61,64,72,77	0
2	EDO	C	302	4/4	0.93	0.11	47,47,48,49	0
2	EDO	D	302	4/4	0.94	0.09	26,27,29,34	0
2	EDO	A	301	4/4	0.94	0.09	47,47,48,50	0
6	SO4	D	304	5/5	0.94	0.27	58,59,63,66	5
2	EDO	B	302	4/4	0.95	0.09	29,34,39,39	0
2	EDO	D	303	4/4	0.98	0.06	35,37,40,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around YEK D 301:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.