



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 12:20 PM UTC

PDB ID : 9FT0 / pdb_00009ft0
Title : Yeast 20S proteasome in complex with epoxyketone inhibitor 16
Authors : Maurits, E.; Huber, E.M.; Dekker, P.M.; Wang, X.; Heinemeyer, W.; Florea, B.I.; Groll, M.; Overkleeft, H.S.
Deposited on : 2024-06-23
Resolution : 2.75 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| R_{free} | 180053 | 1009 (2.76-2.76) |
| Clashscore | 190562 | 1044 (2.76-2.76) |
| Ramachandran outliers | 187476 | 1024 (2.76-2.76) |
| Sidechain outliers | 187428 | 1024 (2.76-2.76) |
| RSRZ outliers | 180081 | 1009 (2.76-2.76) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 250 | 94% 6% |
| 1 | O | 250 | 92% 7% |
| 2 | B | 258 | 85% 9% 5% |
| 2 | P | 258 | 86% 9% 5% |
| 3 | C | 254 | 86% 8% 5% |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 3 | Q | 254 | 87% 7% • 5% |
| 4 | D | 260 | 85% 8% • 7% |
| 4 | R | 260 | 85% 7% 7% |
| 5 | E | 234 | 87% 12% • |
| 5 | S | 234 | 88% 11% • |
| 6 | F | 288 | 79% 5% • 15% |
| 6 | T | 288 | 78% 6% • 15% |
| 7 | G | 252 | 90% 6% • • |
| 7 | U | 252 | 88% 7% • • |
| 8 | H | 231 | 89% 7% • |
| 8 | V | 231 | 89% 6% • |
| 9 | I | 205 | 92% 7% |
| 9 | W | 205 | 91% 8% |
| 10 | J | 198 | 89% 9% • • |
| 10 | X | 198 | 92% 6% • |
| 11 | K | 211 | 84% 13% • |
| 11 | Y | 211 | 85% 12% • |
| 12 | L | 222 | 91% 8% • |
| 12 | Z | 222 | 91% 8% • |
| 13 | M | 246 | 85% 10% 5% |
| 13 | a | 246 | 83% 11% 5% |
| 14 | N | 196 | 90% 10% |
| 14 | b | 196 | 93% 7% |

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|-----|------|-------|-----|-----------|----------|---------|------------------|
| 19 | SO4 | V | 302 | - | - | X | - |

2 Entry composition

There are 20 unique types of molecules in this entry. The entry contains 49929 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha type-2.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 1 | A | 250 | Total 1915 | C 1219 | N 315 | O 377 | S 4 | 0 | 0 | 0 |
| 1 | O | 250 | Total 1915 | C 1219 | N 315 | O 377 | S 4 | 0 | 0 | 0 |

- Molecule 2 is a protein called Proteasome subunit alpha type-3.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 2 | B | 244 | Total 1904 | C 1201 | N 321 | O 379 | S 3 | 0 | 0 | 0 |
| 2 | P | 244 | Total 1904 | C 1201 | N 321 | O 379 | S 3 | 0 | 0 | 0 |

- Molecule 3 is a protein called Proteasome subunit alpha type-4.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 3 | C | 241 | Total 1890 | C 1181 | N 331 | O 374 | S 4 | 0 | 0 | 0 |
| 3 | Q | 241 | Total 1890 | C 1181 | N 331 | O 374 | S 4 | 0 | 0 | 0 |

- Molecule 4 is a protein called Proteasome subunit alpha type-5.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 4 | D | 242 | Total 1861 | C 1162 | N 314 | O 378 | S 7 | 0 | 0 | 0 |
| 4 | R | 242 | Total 1861 | C 1162 | N 314 | O 378 | S 7 | 0 | 0 | 0 |

- Molecule 5 is a protein called Proteasome subunit alpha type-6.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 5 | E | 233 | Total 1795 | C 1129 | N 312 | O 350 | S 4 | 0 | 0 | 0 |
| 5 | S | 233 | Total 1795 | C 1129 | N 312 | O 350 | S 4 | 0 | 0 | 0 |

- Molecule 6 is a protein called Probable proteasome subunit alpha type-7.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 6 | F | 244 | Total 1896 | C 1205 | N 330 | O 357 | S 4 | 0 | 0 | 0 |
| 6 | T | 244 | Total 1896 | C 1205 | N 330 | O 357 | S 4 | 0 | 0 | 0 |

- Molecule 7 is a protein called Proteasome subunit alpha type-1.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 7 | G | 243 | Total 1921 | C 1221 | N 322 | O 370 | S 8 | 0 | 0 | 0 |
| 7 | U | 243 | Total 1921 | C 1221 | N 322 | O 370 | S 8 | 0 | 0 | 0 |

- Molecule 8 is a protein called Proteasome subunit beta type-2.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 8 | H | 221 | Total 1677 | C 1057 | N 292 | O 321 | S 7 | 0 | 0 | 0 |
| 8 | V | 221 | Total 1677 | C 1057 | N 292 | O 321 | S 7 | 0 | 0 | 0 |

- Molecule 9 is a protein called Proteasome subunit beta type-3.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 9 | I | 204 | Total 1581 | C 1010 | N 258 | O 305 | S 8 | 0 | 0 | 0 |
| 9 | W | 204 | Total 1581 | C 1010 | N 258 | O 305 | S 8 | 0 | 0 | 0 |

- Molecule 10 is a protein called Proteasome subunit beta type-4.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 10 | J | 198 | Total 1585 | C 1005 | N 269 | O 305 | S 6 | 0 | 0 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 10 | X | 198 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1585 | 1005 | 269 | 305 | 6 | | | |

- Molecule 11 is a protein called Proteasome subunit beta type-5.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 11 | K | 211 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1637 | 1041 | 279 | 310 | 7 | | | |
| 11 | Y | 211 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1637 | 1041 | 279 | 310 | 7 | | | |

- Molecule 12 is a protein called Proteasome subunit beta type-6.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 12 | L | 222 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1757 | 1115 | 303 | 335 | 4 | | | |
| 12 | Z | 222 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1757 | 1115 | 303 | 335 | 4 | | | |

- Molecule 13 is a protein called Proteasome subunit beta type-7.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 13 | M | 233 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1824 | 1154 | 312 | 351 | 7 | | | |
| 13 | a | 233 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1824 | 1154 | 312 | 351 | 7 | | | |

- Molecule 14 is a protein called Proteasome subunit beta type-1.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 14 | N | 196 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1512 | 955 | 250 | 300 | 7 | | | |
| 14 | b | 196 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1512 | 955 | 250 | 300 | 7 | | | |

- Molecule 15 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 15 | G | 1 | Total | Mg | 0 | 0 |
| | | | 1 | 1 | | |
| 15 | I | 1 | Total | Mg | 0 | 0 |
| | | | 1 | 1 | | |

Continued on next page...


Continued from previous page...

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 15 | K | 1 | Total Mg 1 1 | 0 | 0 |
| 15 | N | 1 | Total Mg 1 1 | 0 | 0 |
| 15 | V | 1 | Total Mg 1 1 | 0 | 0 |
| 15 | W | 1 | Total Mg 1 1 | 0 | 0 |
| 15 | X | 1 | Total Mg 1 1 | 0 | 0 |
| 15 | Y | 1 | Total Mg 1 1 | 0 | 0 |
| 15 | Z | 1 | Total Mg 1 1 | 0 | 0 |

- Molecule 16 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 16 | G | 1 | Total Cl 1 1 | 0 | 0 |
| 16 | N | 2 | Total Cl 2 2 | 0 | 0 |
| 16 | U | 1 | Total Cl 1 1 | 0 | 0 |
| 16 | b | 1 | Total Cl 1 1 | 0 | 0 |

- Molecule 17 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).




| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | |
|-----|-------|----------|-------|----|---|---|---|---------|---------|---|
| | | | Total | C | N | O | S | | | |
| 17 | H | 1 | Total | 12 | 6 | 1 | 4 | 1 | 0 | 0 |
| 17 | K | 1 | Total | 12 | 6 | 1 | 4 | 1 | 0 | 0 |
| 17 | X | 1 | Total | 12 | 6 | 1 | 4 | 1 | 0 | 0 |

- Molecule 18 is (2S)-N-[(2S)-1-[[[(1S)-2-cyclohexyl-1-[(2R,3S,6R,7S)-3-methanoyl-2,6-dimethyl-6,7-bis(oxidanyl)-1,4-oxazepan-7-yl]ethyl]amino]-3-(4-methoxyphenyl)-1-oxidanylidene-propan-2-yl]-2-(2-morpholin-4-ylethanoylamino)-4-oxidanyl-butanamide (CCD ID: A1IFL) (formula: C₃₆H₅₇N₅O₁₀) (labeled as "Ligand of Interest" by depositor).



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---|----|---------|---------|
| 18 | H | 1 | Total | C | N | O | 0 | 0 |
| | | | 51 | 36 | 5 | 10 | | |
| 18 | K | 1 | Total | C | N | O | 0 | 0 |
| | | | 51 | 36 | 5 | 10 | | |
| 18 | V | 1 | Total | C | N | O | 0 | 0 |
| | | | 51 | 36 | 5 | 10 | | |
| 18 | Y | 1 | Total | C | N | O | 0 | 0 |
| | | | 51 | 36 | 5 | 10 | | |

- Molecule 19 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|-----|---------|---------|
| 19 | K | 1 | Total | O S | 0 | 0 |
| | | | 5 | 4 1 | | |
| 19 | V | 1 | Total | O S | 0 | 0 |
| | | | 5 | 4 1 | | |
| 19 | Y | 1 | Total | O S | 0 | 0 |
| | | | 5 | 4 1 | | |

- Molecule 20 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---------|---------|
| 20 | A | 2 | Total | O | 0 | 0 |
| | | | 2 | 2 | | |
| 20 | B | 6 | Total | O | 0 | 0 |
| | | | 6 | 6 | | |
| 20 | C | 3 | Total | O | 0 | 0 |
| | | | 3 | 3 | | |

Continued on next page...


Continued from previous page...

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|------------------|---------|---------|
| 20 | D | 2 | Total O 2 2 | 0 | 0 |
| 20 | E | 4 | Total O 4 4 | 0 | 0 |
| 20 | F | 5 | Total O 5 5 | 0 | 0 |
| 20 | G | 4 | Total O 4 4 | 0 | 0 |
| 20 | H | 4 | Total O 4 4 | 0 | 0 |
| 20 | I | 3 | Total O 3 3 | 0 | 0 |
| 20 | J | 7 | Total O 7 7 | 0 | 0 |
| 20 | K | 6 | Total O 6 6 | 0 | 0 |
| 20 | L | 5 | Total O 5 5 | 0 | 0 |
| 20 | M | 5 | Total O 5 5 | 0 | 0 |
| 20 | N | 8 | Total O 8 8 | 0 | 0 |
| 20 | O | 2 | Total O 2 2 | 0 | 0 |
| 20 | P | 6 | Total O 6 6 | 0 | 0 |
| 20 | Q | 3 | Total O 3 3 | 0 | 0 |
| 20 | R | 2 | Total O 2 2 | 0 | 0 |
| 20 | S | 4 | Total O 4 4 | 0 | 0 |
| 20 | T | 7 | Total O 7 7 | 0 | 0 |
| 20 | U | 8 | Total O 8 8 | 0 | 0 |
| 20 | V | 4 | Total O 4 4 | 0 | 0 |
| 20 | W | 5 | Total O 5 5 | 0 | 0 |
| 20 | X | 11 | Total O 11 11 | 0 | 0 |



Continued on next page...

Continued from previous page...


| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|------------|--------------|-----------------|------------------|----------------|----------------|
| 20 | Y | 7 | Total O 7 7 | 0 | 0 |
| 20 | Z | 10 | Total O 10 10 | 0 | 0 |
| 20 | a | 10 | Total O 10 10 | 0 | 0 |
| 20 | b | 7 | Total O 7 7 | 0 | 0 |





- Molecule 3: Proteasome subunit alpha type-4




- Molecule 4: Proteasome subunit alpha type-5




- Molecule 4: Proteasome subunit alpha type-5




- Molecule 5: Proteasome subunit alpha type-6



- Molecule 5: Proteasome subunit alpha type-6




Chain I:  92% 7%




- Molecule 9: Proteasome subunit beta type-3

Chain W:  91% 8%




- Molecule 10: Proteasome subunit beta type-4

Chain J:  89% 9% 2%





- Molecule 10: Proteasome subunit beta type-4

Chain X:  92% 6% 2%




- Molecule 11: Proteasome subunit beta type-5

Chain K:  84% 13% 3%





- Molecule 11: Proteasome subunit beta type-5

Chain Y:  85% 12% 3%




- Molecule 12: Proteasome subunit beta type-6

Chain L:  91% 8%





- Molecule 12: Proteasome subunit beta type-6

Chain Z:  91% 8%





- Molecule 13: Proteasome subunit beta type-7

Chain M:  85% 10% 5%



- Molecule 13: Proteasome subunit beta type-7

Chain a:  83% 11% 5%



- Molecule 14: Proteasome subunit beta type-1

Chain N:  90% 10%



- Molecule 14: Proteasome subunit beta type-1

Chain b:  93% 7%



4 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 1 21 1 | Depositor |
| Cell constants a, b, c, α , β , γ | 135.52Å 299.48Å 145.00Å 90.00° 112.90° 90.00° | Depositor |
| Resolution (Å) | 30.00 – 2.75 30.00 – 2.75 | Depositor EDS |
| % Data completeness (in resolution range) | 96.3 (30.00-2.75) 96.3 (30.00-2.75) | Depositor EDS |
| R_{merge} | 0.10 | Depositor |
| R_{sym} | (Not available) | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 2.24 (at 2.76Å) | Xtrriage |
| Refinement program | REFMAC 5.8.0267 | Depositor |
| R, R_{free} | 0.190 , 0.232 (Not available) , 0.238 | Depositor DCC |
| R_{free} test set | 13235 reflections (5.00%) | wwPDB-VP |
| Wilson B-factor (Å ²) | 56.2 | Xtrriage |
| Anisotropy | 0.750 | Xtrriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.29 , 52.2 | EDS |
| L-test for twinning ² | $\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$ | Xtrriage |
| Estimated twinning fraction | No twinning to report. | Xtrriage |
| F_o, F_c correlation | 0.96 | EDS |
| Total number of atoms | 49929 | wwPDB-VP |
| Average B, all atoms (Å ²) | 73.0 | wwPDB-VP |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4, A1IFL, MES, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|-----------------|
| | | RMSZ | # $ Z > 5$ | RMSZ | # $ Z > 5$ |
| 1 | A | 1.03 | 0/1952 | 1.43 | 0/2642 |
| 1 | O | 1.03 | 0/1952 | 1.42 | 0/2642 |
| 2 | B | 1.03 | 0/1934 | 1.47 | 0/2618 |
| 2 | P | 1.03 | 0/1934 | 1.47 | 0/2618 |
| 3 | C | 1.04 | 0/1919 | 1.49 | 0/2598 |
| 3 | Q | 1.04 | 0/1919 | 1.49 | 0/2598 |
| 4 | D | 1.04 | 0/1886 | 1.48 | 0/2541 |
| 4 | R | 1.04 | 0/1886 | 1.49 | 0/2541 |
| 5 | E | 1.03 | 0/1823 | 1.45 | 0/2463 |
| 5 | S | 1.03 | 0/1823 | 1.45 | 2/2463 (0.1%) |
| 6 | F | 1.02 | 0/1936 | 1.48 | 2/2614 (0.1%) |
| 6 | T | 1.03 | 0/1936 | 1.49 | 2/2614 (0.1%) |
| 7 | G | 1.01 | 0/1959 | 1.43 | 0/2652 |
| 7 | U | 1.01 | 0/1959 | 1.43 | 0/2652 |
| 8 | H | 1.04 | 0/1708 | 1.43 | 2/2316 (0.1%) |
| 8 | V | 1.04 | 0/1708 | 1.45 | 2/2316 (0.1%) |
| 9 | I | 1.02 | 0/1611 | 1.43 | 1/2174 (0.0%) |
| 9 | W | 1.02 | 0/1611 | 1.43 | 1/2174 (0.0%) |
| 10 | J | 1.03 | 1/1613 (0.1%) | 1.45 | 2/2173 (0.1%) |
| 10 | X | 1.03 | 1/1613 (0.1%) | 1.43 | 1/2173 (0.0%) |
| 11 | K | 1.00 | 0/1674 | 1.47 | 1/2264 (0.0%) |
| 11 | Y | 1.01 | 0/1674 | 1.49 | 5/2264 (0.2%) |
| 12 | L | 1.02 | 0/1795 | 1.41 | 0/2420 |
| 12 | Z | 1.02 | 0/1795 | 1.41 | 0/2420 |
| 13 | M | 1.03 | 0/1855 | 1.39 | 2/2514 (0.1%) |
| 13 | a | 1.03 | 0/1855 | 1.39 | 0/2514 |
| 14 | N | 1.02 | 0/1541 | 1.45 | 0/2087 |
| 14 | b | 1.03 | 0/1541 | 1.45 | 0/2087 |
| All | All | 1.03 | 2/50412 (0.0%) | 1.45 | 23/68152 (0.0%) |

All (2) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|-------|-------------|----------|
| 10 | X | 133 | HIS | CE1-NE2 | -8.07 | 1.24 | 1.32 |
| 10 | J | 133 | HIS | CE1-NE2 | -7.46 | 1.25 | 1.32 |

The worst 5 of 23 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|-------|-------------|----------|
| 6 | T | 77 | LEU | CA-C-N | 7.34 | 124.87 | 120.24 |
| 6 | T | 77 | LEU | C-N-CA | 7.34 | 124.87 | 120.24 |
| 10 | J | 145 | ASP | CA-CB-CG | 6.77 | 119.37 | 112.60 |
| 11 | Y | 211 | ILE | N-CA-CB | -6.42 | 103.69 | 111.21 |
| 11 | K | 107 | LYS | CB-CA-C | -5.67 | 102.23 | 110.96 |

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 1915 | 0 | 1929 | 8 | 0 |
| 1 | O | 1915 | 0 | 1929 | 11 | 0 |
| 2 | B | 1904 | 0 | 1904 | 12 | 0 |
| 2 | P | 1904 | 0 | 1904 | 10 | 0 |
| 3 | C | 1890 | 0 | 1903 | 11 | 0 |
| 3 | Q | 1890 | 0 | 1903 | 8 | 0 |
| 4 | D | 1861 | 0 | 1839 | 11 | 0 |
| 4 | R | 1861 | 0 | 1839 | 8 | 0 |
| 5 | E | 1795 | 0 | 1800 | 19 | 0 |
| 5 | S | 1795 | 0 | 1800 | 15 | 0 |
| 6 | F | 1896 | 0 | 1889 | 5 | 0 |
| 6 | T | 1896 | 0 | 1889 | 9 | 0 |
| 7 | G | 1921 | 0 | 1913 | 11 | 0 |
| 7 | U | 1921 | 0 | 1913 | 12 | 0 |
| 8 | H | 1677 | 0 | 1678 | 6 | 0 |
| 8 | V | 1677 | 0 | 1678 | 4 | 0 |
| 9 | I | 1581 | 0 | 1574 | 8 | 0 |
| 9 | W | 1581 | 0 | 1574 | 10 | 0 |
| 10 | J | 1585 | 0 | 1590 | 15 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 10 | X | 1585 | 0 | 1590 | 14 | 0 |
| 11 | K | 1637 | 0 | 1585 | 20 | 0 |
| 11 | Y | 1637 | 0 | 1585 | 23 | 0 |
| 12 | L | 1757 | 0 | 1711 | 11 | 0 |
| 12 | Z | 1757 | 0 | 1711 | 14 | 0 |
| 13 | M | 1824 | 0 | 1832 | 8 | 0 |
| 13 | a | 1824 | 0 | 1832 | 11 | 0 |
| 14 | N | 1512 | 0 | 1481 | 13 | 0 |
| 14 | b | 1512 | 0 | 1481 | 8 | 0 |
| 15 | G | 1 | 0 | 0 | 0 | 0 |
| 15 | I | 1 | 0 | 0 | 0 | 0 |
| 15 | K | 1 | 0 | 0 | 0 | 0 |
| 15 | N | 1 | 0 | 0 | 0 | 0 |
| 15 | V | 1 | 0 | 0 | 0 | 0 |
| 15 | W | 1 | 0 | 0 | 0 | 0 |
| 15 | X | 1 | 0 | 0 | 0 | 0 |
| 15 | Y | 1 | 0 | 0 | 0 | 0 |
| 15 | Z | 1 | 0 | 0 | 0 | 0 |
| 16 | G | 1 | 0 | 0 | 0 | 0 |
| 16 | N | 2 | 0 | 0 | 1 | 0 |
| 16 | U | 1 | 0 | 0 | 0 | 0 |
| 16 | b | 1 | 0 | 0 | 0 | 0 |
| 17 | H | 12 | 0 | 13 | 2 | 0 |
| 17 | K | 12 | 0 | 13 | 0 | 0 |
| 17 | X | 12 | 0 | 13 | 0 | 0 |
| 18 | H | 51 | 0 | 0 | 1 | 0 |
| 18 | K | 51 | 0 | 0 | 2 | 0 |
| 18 | V | 51 | 0 | 0 | 4 | 0 |
| 18 | Y | 51 | 0 | 0 | 4 | 0 |
| 19 | K | 5 | 0 | 0 | 0 | 0 |
| 19 | V | 5 | 0 | 0 | 2 | 0 |
| 19 | Y | 5 | 0 | 0 | 0 | 0 |
| 20 | A | 2 | 0 | 0 | 0 | 0 |
| 20 | B | 6 | 0 | 0 | 0 | 0 |
| 20 | C | 3 | 0 | 0 | 0 | 0 |
| 20 | D | 2 | 0 | 0 | 0 | 0 |
| 20 | E | 4 | 0 | 0 | 0 | 0 |
| 20 | F | 5 | 0 | 0 | 0 | 0 |
| 20 | G | 4 | 0 | 0 | 0 | 0 |
| 20 | H | 4 | 0 | 0 | 0 | 0 |
| 20 | I | 3 | 0 | 0 | 0 | 0 |
| 20 | J | 7 | 0 | 0 | 1 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 20 | K | 6 | 0 | 0 | 0 | 0 |
| 20 | L | 5 | 0 | 0 | 0 | 0 |
| 20 | M | 5 | 0 | 0 | 0 | 0 |
| 20 | N | 8 | 0 | 0 | 0 | 0 |
| 20 | O | 2 | 0 | 0 | 0 | 0 |
| 20 | P | 6 | 0 | 0 | 0 | 0 |
| 20 | Q | 3 | 0 | 0 | 0 | 0 |
| 20 | R | 2 | 0 | 0 | 0 | 0 |
| 20 | S | 4 | 0 | 0 | 0 | 0 |
| 20 | T | 7 | 0 | 0 | 0 | 0 |
| 20 | U | 8 | 0 | 0 | 0 | 0 |
| 20 | V | 4 | 0 | 0 | 0 | 0 |
| 20 | W | 5 | 0 | 0 | 0 | 0 |
| 20 | X | 11 | 0 | 0 | 0 | 0 |
| 20 | Y | 7 | 0 | 0 | 0 | 0 |
| 20 | Z | 10 | 0 | 0 | 0 | 0 |
| 20 | a | 10 | 0 | 0 | 0 | 0 |
| 20 | b | 7 | 0 | 0 | 0 | 0 |
| All | All | 49929 | 0 | 49295 | 273 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 273 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|--------------------|--------------------------|-------------------|
| 11:Y:104:TYR:CZ | 11:Y:182:GLU:HG3 | 2.07 | 0.90 |
| 11:K:104:TYR:CZ | 11:K:182:GLU:HG3 | 2.08 | 0.88 |
| 19:V:302:SO4:O3 | 18:V:303:A1IFL:O58 | 1.92 | 0.87 |
| 6:T:91:GLU:HG2 | 6:T:111:ARG:HB3 | 1.67 | 0.76 |
| 6:F:91:GLU:HG2 | 6:F:111:ARG:HB3 | 1.68 | 0.75 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|----------|-------------|-----|
| 1 | A | 248/250 (99%) | 240 (97%) | 6 (2%) | 2 (1%) | 16 | 30 |
| 1 | O | 248/250 (99%) | 241 (97%) | 5 (2%) | 2 (1%) | 16 | 30 |
| 2 | B | 242/258 (94%) | 232 (96%) | 7 (3%) | 3 (1%) | 10 | 20 |
| 2 | P | 242/258 (94%) | 232 (96%) | 7 (3%) | 3 (1%) | 10 | 20 |
| 3 | C | 239/254 (94%) | 227 (95%) | 9 (4%) | 3 (1%) | 9 | 18 |
| 3 | Q | 239/254 (94%) | 228 (95%) | 8 (3%) | 3 (1%) | 9 | 18 |
| 4 | D | 240/260 (92%) | 230 (96%) | 8 (3%) | 2 (1%) | 16 | 30 |
| 4 | R | 240/260 (92%) | 230 (96%) | 8 (3%) | 2 (1%) | 16 | 30 |
| 5 | E | 231/234 (99%) | 218 (94%) | 11 (5%) | 2 (1%) | 14 | 28 |
| 5 | S | 231/234 (99%) | 218 (94%) | 11 (5%) | 2 (1%) | 14 | 28 |
| 6 | F | 242/288 (84%) | 234 (97%) | 8 (3%) | 0 | 100 | 100 |
| 6 | T | 242/288 (84%) | 234 (97%) | 8 (3%) | 0 | 100 | 100 |
| 7 | G | 241/252 (96%) | 234 (97%) | 6 (2%) | 1 (0%) | 30 | 50 |
| 7 | U | 241/252 (96%) | 234 (97%) | 6 (2%) | 1 (0%) | 30 | 50 |
| 8 | H | 219/231 (95%) | 214 (98%) | 5 (2%) | 0 | 100 | 100 |
| 8 | V | 219/231 (95%) | 214 (98%) | 5 (2%) | 0 | 100 | 100 |
| 9 | I | 202/205 (98%) | 194 (96%) | 8 (4%) | 0 | 100 | 100 |
| 9 | W | 202/205 (98%) | 194 (96%) | 8 (4%) | 0 | 100 | 100 |
| 10 | J | 196/198 (99%) | 191 (97%) | 5 (3%) | 0 | 100 | 100 |
| 10 | X | 196/198 (99%) | 191 (97%) | 5 (3%) | 0 | 100 | 100 |
| 11 | K | 209/211 (99%) | 204 (98%) | 5 (2%) | 0 | 100 | 100 |
| 11 | Y | 209/211 (99%) | 205 (98%) | 4 (2%) | 0 | 100 | 100 |
| 12 | L | 220/222 (99%) | 215 (98%) | 5 (2%) | 0 | 100 | 100 |
| 12 | Z | 220/222 (99%) | 215 (98%) | 5 (2%) | 0 | 100 | 100 |
| 13 | M | 231/246 (94%) | 220 (95%) | 11 (5%) | 0 | 100 | 100 |
| 13 | a | 231/246 (94%) | 221 (96%) | 10 (4%) | 0 | 100 | 100 |
| 14 | N | 194/196 (99%) | 186 (96%) | 8 (4%) | 0 | 100 | 100 |
| 14 | b | 194/196 (99%) | 185 (95%) | 9 (5%) | 0 | 100 | 100 |
| All | All | 6308/6610 (95%) | 6081 (96%) | 201 (3%) | 26 (0%) | 30 | 50 |

5 of 26 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 166 | LYS |
| 3 | C | 203 | THR |
| 5 | E | 201 | ARG |
| 1 | O | 166 | LYS |
| 3 | Q | 203 | THR |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|----------------|-----------|----------|-------------|----|
| 1 | A | 209/209 (100%) | 205 (98%) | 4 (2%) | 50 | 70 |
| 1 | O | 209/209 (100%) | 205 (98%) | 4 (2%) | 50 | 70 |
| 2 | B | 203/216 (94%) | 197 (97%) | 6 (3%) | 36 | 60 |
| 2 | P | 203/216 (94%) | 197 (97%) | 6 (3%) | 36 | 60 |
| 3 | C | 213/226 (94%) | 206 (97%) | 7 (3%) | 33 | 57 |
| 3 | Q | 213/226 (94%) | 206 (97%) | 7 (3%) | 33 | 57 |
| 4 | D | 198/215 (92%) | 191 (96%) | 7 (4%) | 32 | 56 |
| 4 | R | 198/215 (92%) | 191 (96%) | 7 (4%) | 32 | 56 |
| 5 | E | 192/193 (100%) | 186 (97%) | 6 (3%) | 35 | 59 |
| 5 | S | 192/193 (100%) | 186 (97%) | 6 (3%) | 35 | 59 |
| 6 | F | 201/239 (84%) | 192 (96%) | 9 (4%) | 24 | 46 |
| 6 | T | 201/239 (84%) | 192 (96%) | 9 (4%) | 24 | 46 |
| 7 | G | 207/210 (99%) | 200 (97%) | 7 (3%) | 32 | 57 |
| 7 | U | 207/210 (99%) | 200 (97%) | 7 (3%) | 32 | 57 |
| 8 | H | 180/189 (95%) | 174 (97%) | 6 (3%) | 33 | 57 |
| 8 | V | 180/189 (95%) | 174 (97%) | 6 (3%) | 33 | 57 |
| 9 | I | 172/173 (99%) | 169 (98%) | 3 (2%) | 53 | 71 |
| 9 | W | 172/173 (99%) | 169 (98%) | 3 (2%) | 53 | 71 |
| 10 | J | 175/175 (100%) | 166 (95%) | 9 (5%) | 21 | 40 |
| 10 | X | 175/175 (100%) | 169 (97%) | 6 (3%) | 32 | 57 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|-------------|----|
| 11 | K | 168/168 (100%) | 156 (93%) | 12 (7%) | 13 | 25 |
| 11 | Y | 168/168 (100%) | 156 (93%) | 12 (7%) | 13 | 25 |
| 12 | L | 185/185 (100%) | 176 (95%) | 9 (5%) | 22 | 43 |
| 12 | Z | 185/185 (100%) | 176 (95%) | 9 (5%) | 22 | 43 |
| 13 | M | 199/208 (96%) | 190 (96%) | 9 (4%) | 24 | 46 |
| 13 | a | 199/208 (96%) | 190 (96%) | 9 (4%) | 24 | 46 |
| 14 | N | 162/162 (100%) | 160 (99%) | 2 (1%) | 63 | 78 |
| 14 | b | 162/162 (100%) | 160 (99%) | 2 (1%) | 63 | 78 |
| All | All | 5328/5536 (96%) | 5139 (96%) | 189 (4%) | 32 | 56 |

5 of 189 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 4 | R | 124 | ARG |
| 8 | V | 22 | GLN |
| 4 | R | 235 | LEU |
| 6 | T | 181 | GLU |
| 9 | W | 182 | TRP |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 154 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 7 | U | 83 | ASN |
| 12 | Z | 80 | ASN |
| 7 | U | 167 | GLN |
| 10 | X | 63 | ASN |
| 13 | a | 108 | ASN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 14 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|-------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 18 | A1IFL | Y | 303 | 11 | 49,54,54 | 1.63 | 7 (14%) | 58,75,75 | 1.10 | 3 (5%) |
| 19 | SO4 | K | 303 | - | 4,4,4 | 0.58 | 0 | 6,6,6 | 0.09 | 0 |
| 19 | SO4 | Y | 302 | - | 4,4,4 | 0.33 | 0 | 6,6,6 | 0.08 | 0 |
| 17 | MES | X | 202 | - | 12,12,12 | 0.75 | 0 | 15,16,16 | 0.33 | 0 |
| 19 | SO4 | V | 302 | - | 4,4,4 | 0.49 | 0 | 6,6,6 | 0.12 | 0 |
| 18 | A1IFL | V | 303 | 8 | 49,54,54 | 1.18 | 3 (6%) | 58,75,75 | 1.14 | 5 (8%) |
| 18 | A1IFL | H | 302 | 8 | 49,54,54 | 1.15 | 2 (4%) | 58,75,75 | 1.25 | 7 (12%) |
| 17 | MES | H | 301 | - | 12,12,12 | 0.70 | 0 | 15,16,16 | 0.29 | 0 |
| 17 | MES | K | 302 | - | 12,12,12 | 0.73 | 0 | 15,16,16 | 0.45 | 0 |
| 18 | A1IFL | K | 304 | 11 | 49,54,54 | 1.86 | 9 (18%) | 58,75,75 | 1.17 | 6 (10%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|-------|-------|-----|------|---------|-------------|---------|
| 18 | A1IFL | Y | 303 | 11 | - | 9/37/85/85 | 0/3/4/4 |
| 17 | MES | X | 202 | - | - | 2/6/14/14 | 0/1/1/1 |
| 18 | A1IFL | V | 303 | 8 | - | 6/37/85/85 | 0/3/4/4 |
| 18 | A1IFL | H | 302 | 8 | - | 13/37/85/85 | 0/3/4/4 |
| 17 | MES | H | 301 | - | - | 2/6/14/14 | 0/1/1/1 |
| 17 | MES | K | 302 | - | - | 4/6/14/14 | 0/1/1/1 |
| 18 | A1IFL | K | 304 | 11 | - | 12/37/85/85 | 0/3/4/4 |

The worst 5 of 21 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|-------|---------|-------|-------------|----------|
| 18 | K | 304 | A1IFL | CB-CG | -6.89 | 1.35 | 1.51 |
| 18 | H | 302 | A1IFL | CB-CG | -6.21 | 1.36 | 1.51 |
| 18 | Y | 303 | A1IFL | CB-CG | -6.14 | 1.36 | 1.51 |
| 18 | V | 303 | A1IFL | CB-CG | -5.45 | 1.38 | 1.51 |
| 18 | K | 304 | A1IFL | O58-C11 | -5.12 | 1.37 | 1.44 |

The worst 5 of 21 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|-------|------------|-------|-------------|----------|
| 18 | H | 302 | A1IFL | CM-OH-CZ | -3.86 | 109.22 | 117.50 |
| 18 | V | 303 | A1IFL | CB1-CA3-N2 | -3.05 | 104.88 | 111.75 |
| 18 | V | 303 | A1IFL | CE2-CD2-NB | 3.04 | 114.74 | 110.12 |
| 18 | H | 302 | A1IFL | CA-NB-CD2 | -2.82 | 106.75 | 111.14 |
| 18 | H | 302 | A1IFL | C3-CA1-N | -2.73 | 105.51 | 110.91 |

There are no chirality outliers.

5 of 48 torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|-------|-----------------|
| 17 | K | 302 | MES | C7-C8-S-O1S |
| 17 | X | 202 | MES | C8-C7-N4-C3 |
| 18 | H | 302 | A1IFL | CA1-C3-C4-O3 |
| 18 | H | 302 | A1IFL | C21-C31-C48-C49 |
| 18 | H | 302 | A1IFL | C31-C48-C49-C54 |


There are no ring outliers.


6 monomers are involved in 13 short contacts:


| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|-------|---------|--------------|
| 18 | Y | 303 | A1IFL | 4 | 0 |
| 19 | V | 302 | SO4 | 2 | 0 |
| 18 | V | 303 | A1IFL | 4 | 0 |
| 18 | H | 302 | A1IFL | 1 | 0 |
| 17 | H | 301 | MES | 2 | 0 |
| 18 | K | 304 | A1IFL | 2 | 0 |


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|----------------|--------|--------------|-----------------------|-------|
| 1 | A | 250/250 (100%) | -0.21 | 1 (0%) 88 89 | 49, 65, 94, 137 | 0 |
| 1 | O | 250/250 (100%) | -0.13 | 3 (1%) 76 76 | 56, 73, 106, 145 | 0 |
| 2 | B | 244/258 (94%) | 0.12 | 5 (2%) 65 63 | 53, 72, 114, 147 | 0 |
| 2 | P | 244/258 (94%) | 0.03 | 7 (2%) 53 52 | 56, 73, 120, 153 | 0 |
| 3 | C | 241/254 (94%) | 0.03 | 8 (3%) 49 47 | 53, 75, 131, 143 | 0 |
| 3 | Q | 241/254 (94%) | 0.12 | 6 (2%) 58 56 | 57, 84, 146, 165 | 0 |
| 4 | D | 242/260 (93%) | 0.07 | 8 (3%) 49 47 | 55, 76, 114, 153 | 0 |
| 4 | R | 242/260 (93%) | 0.10 | 9 (3%) 45 42 | 58, 81, 121, 162 | 0 |
| 5 | E | 233/234 (99%) | 0.09 | 3 (1%) 75 75 | 57, 77, 112, 140 | 0 |
| 5 | S | 233/234 (99%) | 0.16 | 5 (2%) 63 61 | 57, 80, 114, 160 | 0 |
| 6 | F | 244/288 (84%) | -0.04 | 2 (0%) 82 82 | 52, 70, 106, 138 | 0 |
| 6 | T | 244/288 (84%) | -0.02 | 2 (0%) 82 82 | 55, 73, 112, 138 | 0 |
| 7 | G | 243/252 (96%) | -0.12 | 4 (1%) 70 69 | 50, 65, 96, 148 | 0 |
| 7 | U | 243/252 (96%) | -0.19 | 3 (1%) 76 76 | 53, 66, 93, 148 | 0 |
| 8 | H | 221/231 (95%) | -0.13 | 0 100 100 | 51, 62, 86, 141 | 0 |
| 8 | V | 221/231 (95%) | -0.12 | 1 (0%) 87 87 | 53, 66, 92, 136 | 0 |
| 9 | I | 204/205 (99%) | -0.26 | 0 100 100 | 48, 61, 84, 112 | 0 |
| 9 | W | 204/205 (99%) | -0.32 | 1 (0%) 87 87 | 46, 64, 85, 102 | 0 |
| 10 | J | 198/198 (100%) | -0.15 | 4 (2%) 65 63 | 48, 64, 90, 148 | 0 |
| 10 | X | 198/198 (100%) | -0.20 | 2 (1%) 79 79 | 51, 66, 90, 149 | 0 |
| 11 | K | 211/211 (100%) | -0.14 | 2 (0%) 81 81 | 48, 63, 90, 111 | 0 |
| 11 | Y | 211/211 (100%) | -0.12 | 3 (1%) 73 73 | 52, 63, 91, 109 | 0 |
| 12 | L | 222/222 (100%) | -0.21 | 0 100 100 | 48, 65, 88, 103 | 0 |
| 12 | Z | 222/222 (100%) | -0.18 | 0 100 100 | 50, 65, 92, 108 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|-----------------|--------|---------------|-----------------------|-------|
| 13 | M | 233/246 (94%) | -0.23 | 3 (1%) 75 75 | 48, 63, 84, 110 | 0 |
| 13 | a | 233/246 (94%) | -0.23 | 2 (0%) 81 81 | 46, 62, 82, 111 | 0 |
| 14 | N | 196/196 (100%) | -0.26 | 0 100 100 | 47, 59, 88, 113 | 0 |
| 14 | b | 196/196 (100%) | -0.29 | 0 100 100 | 46, 59, 86, 108 | 0 |
| All | All | 6364/6610 (96%) | -0.09 | 84 (1%) 75 75 | 46, 68, 108, 165 | 0 |

The worst 5 of 84 RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 3 | Q | 50 | LEU | 5.9 |
| 4 | D | 119 | ALA | 5.8 |
| 4 | R | 125 | LEU | 5.0 |
| 3 | C | 50 | LEU | 4.6 |
| 3 | Q | 49 | THR | 4.5 |

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|-------|-------|-----|-------|------|------|----------------------------|-------|
| 15 | MG | Z | 301 | 1/1 | 0.61 | 0.26 | 98,98,98,98 | 0 |
| 17 | MES | H | 301 | 12/12 | 0.88 | 0.24 | 60,64,68,68 | 12 |
| 18 | A1IFL | H | 302 | 51/51 | 0.90 | 0.15 | 50,65,74,76 | 51 |
| 18 | A1IFL | V | 303 | 51/51 | 0.90 | 0.15 | 55,66,83,84 | 51 |
| 15 | MG | G | 301 | 1/1 | 0.91 | 0.08 | 75,75,75,75 | 0 |
| 15 | MG | K | 301 | 1/1 | 0.92 | 0.13 | 89,89,89,89 | 0 |
| 16 | CL | N | 203 | 1/1 | 0.93 | 0.12 | 95,95,95,95 | 0 |

Continued on next page...

Continued from previous page...




| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|-------|-------|-----|-------|------|------|----------------------------|-------|
| 18 | A1IFL | Y | 303 | 51/51 | 0.93 | 0.10 | 46,56,72,72 | 0 |
| 18 | A1IFL | K | 304 | 51/51 | 0.94 | 0.09 | 48,52,76,76 | 0 |
| 15 | MG | V | 301 | 1/1 | 0.94 | 0.13 | 112,112,112,112 | 0 |
| 15 | MG | W | 301 | 1/1 | 0.94 | 0.12 | 78,78,78,78 | 0 |
| 19 | SO4 | V | 302 | 5/5 | 0.94 | 0.12 | 78,83,88,88 | 5 |
| 19 | SO4 | K | 303 | 5/5 | 0.95 | 0.16 | 100,107,110,110 | 0 |
| 17 | MES | K | 302 | 12/12 | 0.95 | 0.11 | 75,82,86,88 | 0 |
| 15 | MG | N | 201 | 1/1 | 0.96 | 0.09 | 74,74,74,74 | 0 |
| 15 | MG | Y | 301 | 1/1 | 0.96 | 0.06 | 84,84,84,84 | 0 |
| 17 | MES | X | 202 | 12/12 | 0.96 | 0.11 | 69,78,84,86 | 0 |
| 15 | MG | I | 301 | 1/1 | 0.97 | 0.06 | 75,75,75,75 | 0 |
| 16 | CL | G | 302 | 1/1 | 0.98 | 0.03 | 59,59,59,59 | 0 |
| 16 | CL | U | 301 | 1/1 | 0.98 | 0.05 | 66,66,66,66 | 0 |
| 16 | CL | b | 201 | 1/1 | 0.98 | 0.05 | 70,70,70,70 | 0 |
| 19 | SO4 | Y | 302 | 5/5 | 0.98 | 0.10 | 70,72,73,76 | 5 |
| 16 | CL | N | 202 | 1/1 | 0.99 | 0.06 | 70,70,70,70 | 0 |
| 15 | MG | X | 201 | 1/1 | 0.99 | 0.14 | 52,52,52,52 | 0 |

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around A1IFL V 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A1IFL Y 303:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.