



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:18 AM UTC

PDB ID : 2FU6 / pdb_00002fu6
Title : Zinc-beta-lactamase 11 from *Stenotrophomonas maltophilia* (apo form)
Authors : Nauton, L.; Garau, G.; Kahn, R.; Dideberg, O.
Deposited on : 2006-01-26
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	
1	B	269	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4611 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metallo-beta-lactamase L1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	266	Total	C	N	O	S	0	2	0
			2013	1265	363	377	8			
1	B	266	Total	C	N	O	S	0	2	0
			2013	1265	363	377	8			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	269	Total	O	0	0
			269	269		
4	B	257	Total	O	0	1
			258	258		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Metallo-beta-lactamase L1

Chain A:  92% 6% ..



• Molecule 1: Metallo-beta-lactamase L1

Chain B:  86% 11% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	105.05Å 105.05Å 196.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.89 – 2.05 19.89 – 2.05	Depositor EDS
% Data completeness (in resolution range)	98.9 (19.89-2.05) 98.7 (19.89-2.05)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.60 (at 2.06Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.160 , 0.209 0.173 , 0.174	Depositor DCC
R_{free} test set	2003 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	20.0	Xtrriage
Anisotropy	0.050	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4611	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 78.74 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.4901e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.46	7/2071 (0.3%)	1.01	5/2827 (0.2%)
1	B	1.43	8/2071 (0.4%)	1.02	7/2827 (0.2%)
All	All	1.44	15/4142 (0.4%)	1.02	12/5654 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	276	ARG	C-N	30.44	1.76	1.33
1	A	57	THR	C-N	29.34	1.74	1.34
1	A	275	ALA	C-N	21.03	1.61	1.33
1	A	157	GLY	C-N	20.72	1.63	1.33
1	B	157	GLY	C-N	18.56	1.57	1.33
1	B	57	THR	C-N	17.49	1.56	1.33
1	A	87	MET	C-N	14.55	1.52	1.33
1	B	166	PRO	C-N	14.10	1.51	1.33
1	B	87	MET	C-N	12.78	1.52	1.33
1	A	222	LEU	C-N	12.58	1.50	1.33
1	B	222	LEU	C-N	11.69	1.49	1.33
1	A	166	PRO	C-N	9.10	1.44	1.33
1	B	191	PHE	C-N	7.05	1.42	1.33
1	A	191	PHE	C-N	6.20	1.41	1.33
1	B	189	ALA	CA-CB	5.12	1.60	1.53

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	276	ARG	O-C-N	-18.03	98.80	122.42
1	A	57	THR	O-C-N	-7.82	112.03	122.59
1	B	166	PRO	CA-C-N	-6.22	112.13	122.21
1	B	166	PRO	C-N-CA	-6.22	112.13	122.21
1	B	87	MET	O-C-N	-5.92	114.12	121.64
1	B	48	ILE	CB-CA-C	-5.43	106.09	111.30
1	A	24	GLU	CA-C-N	5.40	131.41	121.70
1	A	24	GLU	C-N-CA	5.40	131.41	121.70
1	A	275	ALA	CA-C-N	-5.38	114.12	122.67
1	A	275	ALA	C-N-CA	-5.38	114.12	122.67
1	B	166	PRO	O-C-N	5.35	129.64	123.06
1	B	68	ASP	N-CA-C	5.12	118.84	112.59

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	222	LEU	Mainchain
1	A	275	ALA	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2013	0	1973	10	0
1	B	2013	0	1972	19	0
2	A	20	0	0	0	0
2	B	20	0	0	0	0
3	A	12	0	16	0	0
3	B	6	0	8	2	0
4	A	269	0	0	0	0
4	B	258	0	0	4	0
All	All	4611	0	3969	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:276:ARG:C	1:B:289:ALA:N	1.76	1.39
1:B:90:GLN:H	1:B:90:GLN:HE21	1.07	0.90
3:B:1:GOL:H11	4:B:383:HOH:O	1.87	0.72
1:B:163:THR:HB	4:B:574:HOH:O	1.93	0.69
1:B:276:ARG:O	1:B:289:ALA:N	2.30	0.61
3:B:1:GOL:C1	4:B:383:HOH:O	2.47	0.60
1:A:41:GLN:HB3	1:A:67[A]:GLU:HG3	1.82	0.60
1:A:24:GLU:N	1:A:25:VAL:HA	2.17	0.60
1:A:179:VAL:HG13	1:A:186:VAL:HG13	1.87	0.57
1:B:315:THR:C	1:B:317:GLY:H	2.12	0.57
1:B:221:SER:HB2	4:B:489:HOH:O	2.10	0.51
1:B:41:GLN:HB3	1:B:67[A]:GLU:HG3	1.93	0.50
1:B:90:GLN:H	1:B:90:GLN:NE2	1.90	0.50
1:A:179:VAL:HG13	1:A:186:VAL:CG1	2.44	0.47
1:B:76:THR:HB	1:B:77:PRO:HD2	1.95	0.47
1:B:315:THR:C	1:B:317:GLY:N	2.73	0.47
1:A:179:VAL:CG1	1:A:186:VAL:HG13	2.45	0.47
1:B:240:LEU:C	1:B:240:LEU:HD23	2.40	0.46
1:B:248:PHE:O	1:B:252[B]:ARG:HG3	2.14	0.46
1:B:219:ALA:O	1:B:262:PRO:HD3	2.16	0.45
1:B:270:ASP:O	1:B:289:ALA:HA	2.16	0.45
1:B:90:GLN:HE21	1:B:90:GLN:N	1.91	0.44
1:A:174:VAL:HG11	1:A:180:ILE:HD11	1.98	0.44
1:B:116:HIS:CE1	1:B:121:HIS:CD2	3.06	0.43
1:A:24:GLU:O	1:A:24:GLU:HG2	2.19	0.42
1:B:105:THR:OG1	1:B:107:ARG:HG2	2.20	0.42
1:A:138:ALA:O	1:A:173:ILE:HA	2.21	0.41
1:B:315:THR:O	1:B:317:GLY:N	2.54	0.41
1:B:41:GLN:CG	1:B:67[A]:GLU:HG3	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/269 (99%)	257 (97%)	9 (3%)	0	100	100
1	B	266/269 (99%)	254 (96%)	11 (4%)	1 (0%)	30	22
All	All	532/538 (99%)	511 (96%)	20 (4%)	1 (0%)	43	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	316	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/202 (100%)	201 (99%)	2 (1%)	68	72
1	B	203/202 (100%)	196 (97%)	7 (3%)	32	27
All	All	406/404 (100%)	397 (98%)	9 (2%)	48	44

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	THR
1	A	212	LYS
1	B	25	VAL
1	B	67[A]	GLU
1	B	67[B]	GLU
1	B	90	GLN
1	B	107	ARG
1	B	306	LYS
1	B	311	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
1	A	196	HIS
1	A	232	GLN
1	A	263	HIS
1	B	90	GLN
1	B	263	HIS
1	B	310	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	11	-	4,4,4	0.30	0	6,6,6	0.32	0
2	SO4	A	5	-	4,4,4	0.16	0	6,6,6	0.37	0
3	GOL	A	2	-	5,5,5	0.83	0	5,5,5	1.30	1 (20%)
2	SO4	B	4	-	4,4,4	0.66	0	6,6,6	0.67	0
2	SO4	B	9	-	4,4,4	0.24	0	6,6,6	0.23	0
2	SO4	B	10	-	4,4,4	0.28	0	6,6,6	0.36	0
3	GOL	B	1	-	5,5,5	0.45	0	5,5,5	1.03	0
3	GOL	A	3	-	5,5,5	0.42	0	5,5,5	0.57	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	7	-	4,4,4	0.21	0	6,6,6	0.37	0
2	SO4	A	8	-	4,4,4	0.36	0	6,6,6	0.69	0
2	SO4	A	6	-	4,4,4	0.36	0	6,6,6	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	1	-	-	3/4/4/4	-
3	GOL	A	3	-	-	2/4/4/4	-
3	GOL	A	2	-	-	4/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2	GOL	C3-C2-C1	2.20	119.87	111.80

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2	GOL	O1-C1-C2-O2
3	A	2	GOL	O1-C1-C2-C3
3	B	1	GOL	C1-C2-C3-O3
3	B	1	GOL	O2-C2-C3-O3
3	A	2	GOL	C1-C2-C3-O3
3	A	2	GOL	O2-C2-C3-O3
3	A	3	GOL	O1-C1-C2-C3
3	A	3	GOL	O1-C1-C2-O2
3	B	1	GOL	O1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	3
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	276:ARG	C	289:ALA	N	1.76
1	A	57:THR	C	67:GLU	N	1.74
1	A	157:GLY	C	160:ASP	N	1.63
1	A	275:ALA	C	288:ARG	N	1.61

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	266/269 (98%)	-0.56	1 (0%) 88 90	11, 18, 30, 47	2 (0%)
1	B	266/269 (98%)	-0.47	2 (0%) 82 84	12, 19, 32, 49	2 (0%)
All	All	532/538 (98%)	-0.51	3 (0%) 85 87	11, 19, 31, 49	4 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	317	GLY	3.0
1	B	316	ALA	2.8
1	A	317	GLY	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	1	6/6	0.77	0.18	45,48,51,54	0
3	GOL	A	2	6/6	0.81	0.15	37,42,43,44	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	B	9	5/5	0.83	0.09	48,49,49,49	5
2	SO4	B	10	5/5	0.83	0.19	36,39,40,41	5
2	SO4	A	8	5/5	0.85	0.14	44,45,49,49	5
2	SO4	B	7	5/5	0.85	0.13	27,29,32,32	5
3	GOL	A	3	6/6	0.89	0.12	42,43,44,44	0
2	SO4	A	11	5/5	0.91	0.11	41,41,42,44	5
2	SO4	A	6	5/5	0.92	0.11	37,38,41,42	5
2	SO4	A	5	5/5	0.97	0.06	22,29,33,33	5
2	SO4	B	4	5/5	0.98	0.06	22,23,27,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.