



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 10:44 AM UTC

PDB ID : 2FUT / pdb\_00002fut  
Title : Crystal Structure of Heparinase II Complexed with a Disaccharide Product  
Authors : Shaya, D.; Cygler, M.  
Deposited on : 2006-01-27  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	748	 86% 12% ..
1	B	748	 3% 85% 13% ..
2	C	2	 50% 50%
2	D	2	 100%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 12752 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called heparinase II protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	743	5949	3832	1003	1091	3	20	0	0	0
1	B	743	5948	3832	1003	1090	3	20	0	0	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	41	MSE	MET	modified residue	UNP Q46080
A	65	MSE	MET	modified residue	UNP Q46080
A	77	MSE	MET	modified residue	UNP Q46080
A	80	MSE	MET	modified residue	UNP Q46080
A	111	MSE	MET	modified residue	UNP Q46080
A	117	MSE	MET	modified residue	UNP Q46080
A	154	MSE	MET	modified residue	UNP Q46080
A	186	MSE	MET	modified residue	UNP Q46080
A	207	MSE	MET	modified residue	UNP Q46080
A	209	MSE	MET	modified residue	UNP Q46080
A	226	MSE	MET	modified residue	UNP Q46080
A	255	MSE	MET	modified residue	UNP Q46080
A	275	MSE	MET	modified residue	UNP Q46080
A	320	MSE	MET	modified residue	UNP Q46080
A	381	MSE	MET	modified residue	UNP Q46080
A	396	MSE	MET	modified residue	UNP Q46080
A	499	MSE	MET	modified residue	UNP Q46080
A	559	MSE	MET	modified residue	UNP Q46080
A	603	MSE	MET	modified residue	UNP Q46080
A	729	MSE	MET	modified residue	UNP Q46080
A	758	ALA	PRO	engineered mutation	UNP Q46080
B	41	MSE	MET	modified residue	UNP Q46080
B	65	MSE	MET	modified residue	UNP Q46080
B	77	MSE	MET	modified residue	UNP Q46080
B	80	MSE	MET	modified residue	UNP Q46080

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	111	MSE	MET	modified residue	UNP Q46080
B	117	MSE	MET	modified residue	UNP Q46080
B	154	MSE	MET	modified residue	UNP Q46080
B	186	MSE	MET	modified residue	UNP Q46080
B	207	MSE	MET	modified residue	UNP Q46080
B	209	MSE	MET	modified residue	UNP Q46080
B	226	MSE	MET	modified residue	UNP Q46080
B	255	MSE	MET	modified residue	UNP Q46080
B	275	MSE	MET	modified residue	UNP Q46080
B	320	MSE	MET	modified residue	UNP Q46080
B	381	MSE	MET	modified residue	UNP Q46080
B	396	MSE	MET	modified residue	UNP Q46080
B	499	MSE	MET	modified residue	UNP Q46080
B	559	MSE	MET	modified residue	UNP Q46080
B	603	MSE	MET	modified residue	UNP Q46080
B	729	MSE	MET	modified residue	UNP Q46080
B	758	ALA	PRO	engineered mutation	UNP Q46080

- Molecule 2 is an oligosaccharide called 4-deoxy-2-O-sulfo-alpha-L-threo-hex-4-enopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	2	Total	C	N	O	S	0	0	0
			35	12	1	19	3			
2	D	2	Total	C	N	O	S	0	0	0
			35	12	1	19	3			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	547	Total 547	O 547	0	0
4	B	236	Total 236	O 236	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: heparinase II protein



- Molecule 1: heparinase II protein



- Molecule 2: 4-deoxy-2-O-sulfo-alpha-L-threo-hex-4-enopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose



SGN1  
UAF2

- Molecule 2: 4-deoxy-2-O-sulfo-alpha-L-threo-hex-4-enopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose

Chain D:

100%

SGN1  
UAF2

## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.10Å 163.30Å 95.30Å 90.00° 105.50° 90.00°	Depositor
Resolution (Å)	91.67 – 2.30 91.67 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.0 (91.67-2.30) 95.9 (91.67-2.30)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.97 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.176 , 0.221 0.181 , 0.220	Depositor DCC
$R_{free}$ test set	3309 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.7	Xtrriage
Anisotropy	0.790	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.030 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12752	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SGN, UAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	1/6091 (0.0%)	0.81	3/8212 (0.0%)
1	B	0.48	0/6090	0.80	1/8210 (0.0%)
All	All	0.52	1/12181 (0.0%)	0.80	4/16422 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	58	VAL	CA-CB	5.14	1.57	1.54

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	388	GLY	CA-C-N	6.00	126.43	119.47
1	A	388	GLY	C-N-CA	6.00	126.43	119.47
1	A	207	MSE	N-CA-C	5.33	116.77	111.07
1	B	207	MSE	N-CA-C	5.33	117.09	111.28

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	91	GLU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5949	0	5859	62	0
1	B	5948	0	5859	61	0
2	C	35	0	13	1	0
2	D	35	0	14	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	547	0	0	6	0
4	B	236	0	0	1	0
All	All	12752	0	11745	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 119 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:296:ARG:HH11	1:A:296:ARG:HG2	1.30	0.96
1:B:717:PHE:HE1	1:B:729:MSE:HE1	1.29	0.95
1:B:606:ASN:HD21	1:B:669:ASN:HB3	1.29	0.94
1:B:473:HIS:H	1:B:473:HIS:CD2	1.85	0.94
1:B:717:PHE:CE1	1:B:729:MSE:HE1	2.04	0.93

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	741/748 (99%)	711 (96%)	29 (4%)	1 (0%)	48	60
1	B	741/748 (99%)	699 (94%)	39 (5%)	3 (0%)	30	38
All	All	1482/1496 (99%)	1410 (95%)	68 (5%)	4 (0%)	36	46

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	515	GLY
1	B	440	HIS
1	A	91	GLU
1	B	399	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	630/614 (103%)	609 (97%)	21 (3%)	33	50
1	B	630/614 (103%)	603 (96%)	27 (4%)	26	39
All	All	1260/1228 (103%)	1212 (96%)	48 (4%)	29	44

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	139	THR
1	B	295	LYS
1	B	171	GLU
1	B	250	ASN
1	B	303	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	81	GLN
1	B	253	GLN
1	B	606	ASN
1	B	250	ASN
1	B	285	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	SGN	C	1	2	20,20,20	2.01	4 (20%)	25,31,31	1.47	2 (8%)
2	UAP	C	2	2	15,15,16	2.08	3 (20%)	18,22,24	1.86	4 (22%)
2	SGN	D	1	2	20,20,20	2.24	4 (20%)	25,31,31	1.44	3 (12%)
2	UAP	D	2	2	15,15,16	2.16	3 (20%)	18,22,24	2.32	3 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SGN	C	1	2	-	1/11/31/31	0/1/1/1
2	UAP	C	2	2	-	0/9/22/25	0/1/1/1

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SGN	D	1	2	-	3/11/31/31	0/1/1/1
2	UAP	D	2	2	-	0/9/22/25	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	UAP	O5-C5	6.58	1.46	1.37
2	C	2	UAP	O5-C5	6.20	1.45	1.37
2	D	1	SGN	O2S-S1	5.51	1.48	1.42
2	D	1	SGN	S1-N2	-5.49	1.52	1.59
2	D	1	SGN	O1S-S1	5.30	1.48	1.42

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	UAP	O5-C5-C4	-7.18	118.34	124.94
2	C	1	SGN	O1S-S1-O2S	-5.38	108.38	120.36
2	C	2	UAP	O5-C5-C4	-4.91	120.43	124.94
2	D	1	SGN	O1S-S1-O2S	-4.63	110.05	120.36
2	D	2	UAP	O5-C5-C6	4.49	120.11	111.85

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	1	SGN	C4-C5-C6-O6
2	D	1	SGN	O5-C5-C6-O6
2	D	1	SGN	C6-O6-S2-O6S
2	C	1	SGN	C5-C6-O6-S2


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	UAP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	723/748 (96%)	-0.38	4 (0%) 85 86	2, 10, 25, 44	0
1	B	723/748 (96%)	0.43	21 (2%) 53 56	9, 28, 45, 54	0
All	All	1446/1496 (96%)	0.03	25 (1%) 69 70	2, 18, 40, 54	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	515	GLY	4.6
1	A	36	VAL	4.2
1	B	87	ALA	3.6
1	B	439	PRO	3.5
1	B	39	VAL	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	UAP	D	2	15/16	0.82	0.13	48,50,51,51	0
2	SGN	D	1	20/20	0.85	0.12	46,50,58,58	0
2	SGN	C	1	20/20	0.93	0.10	15,17,31,31	0
2	UAP	C	2	15/16	0.95	0.08	14,16,18,19	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ZN	B	2	1/1	0.97	0.07	37,37,37,37	0
3	ZN	A	1	1/1	0.99	0.09	10,10,10,10	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.