



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 08:14 AM UTC

PDB ID : 6FUP / pdb_00006fup
Title : F11 T-Cell Receptor Recognising PKYVKQNTLKLAT Peptide Presented by HLA-DR*0101
Authors : Rizkallah, P.J.; Cole, D.K.
Deposited on : 2018-02-27
Resolution : 1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	202	9% 87% 12%
2	B	240	4% 91% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	316	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3985 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Human F11 T-cell Receptor alpha Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	202	Total	C	N	O	S	0	5	0
			1607	1020	261	318	8			

- Molecule 2 is a protein called Human F11 T-Cell Receptor beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	240	Total	C	N	O	S	0	10	0
			1971	1242	340	379	10			

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		
4	A	1	Total	C	O	0	0
			7	4	3		
4	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is SULFATE ION (CCD ID: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	114	Total 114	O 114	0	0
7	B	155	Total 155	O 155	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Human F11 T-cell Receptor alpha Chain



- Molecule 2: Human F11 T-Cell Receptor beta chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.28Å 115.16Å 50.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.41 – 1.72 46.41 – 1.72	Depositor EDS
% Data completeness (in resolution range)	99.4 (46.41-1.72) 99.3 (46.41-1.72)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 1.72Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.176 , 0.215 0.185 , 0.221	Depositor DCC
R_{free} test set	2715 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3985	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, PEG, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	1/1653 (0.1%)	1.05	2/2242 (0.1%)
2	B	1.00	1/2045 (0.0%)	0.98	1/2779 (0.0%)
All	All	1.03	2/3698 (0.1%)	1.01	3/5021 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	151	VAL	N-CA	6.09	1.52	1.46
1	A	53	ALA	CA-C	5.98	1.60	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	2	ASP	N-CA-C	-12.23	91.54	109.15
1	A	160	CYS	N-CA-CB	-6.40	99.67	111.13
1	A	125	ARG	N-CA-C	5.94	119.50	109.76

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	130	SER	Peptide
2	B	1	ALA	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1607	0	1556	19	3
2	B	1971	0	1901	14	0
3	A	52	0	78	1	1
3	B	48	0	72	0	0
4	A	14	0	20	3	0
4	B	7	0	10	1	0
5	A	6	0	8	5	0
5	B	6	0	8	0	0
6	B	5	0	0	0	0
7	A	114	0	0	2	0
7	B	155	0	0	4	0
All	All	3985	0	3653	33	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1:ALA:HB1	7:B:413:HOH:O	1.75	0.87
1:A:26[A]:SER:O	1:A:26[A]:SER:OG	1.94	0.83
1:A:62:GLY:HA2	5:A:316:GOL:H12	1.64	0.80
1:A:156:ILE:O	4:A:314:PEG:H22	1.84	0.77
1:A:62:GLY:CA	5:A:316:GOL:H12	2.20	0.71
2:B:1:ALA:HB2	7:B:420:HOH:O	1.92	0.68
1:A:115:GLN:H	1:A:115:GLN:CD	2.09	0.61
1:A:2:GLN:HB3	7:A:409:HOH:O	2.00	0.61
2:B:11:ILE:HD12	2:B:212:GLY:HA2	1.84	0.58
1:A:63:PHE:O	5:A:316:GOL:O2	2.17	0.57
1:A:3:SER:N	1:A:26[A]:SER:OG	2.37	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:68:GLN:NE2	7:B:405:HOH:O	2.38	0.52
1:A:2:GLN:HA	1:A:26[A]:SER:O	2.11	0.50
2:B:150:HIS:CD2	4:B:313:PEG:H42	2.48	0.49
2:B:176:GLN:HB3	2:B:179:LEU:HD13	1.95	0.48
2:B:13[A]:LYS:HD2	2:B:14:THR:O	2.14	0.48
2:B:7:PRO:HD2	2:B:20[A]:LEU:HD22	1.96	0.47
2:B:5:GLN:HE21	2:B:102:GLY:HA3	1.81	0.46
1:A:126:ASP:HB3	1:A:130:SER:CB	2.45	0.46
2:B:215:GLU:H	2:B:215:GLU:CD	2.23	0.46
1:A:62:GLY:N	5:A:316:GOL:H12	2.32	0.45
1:A:65:ALA:HA	4:A:315:PEG:H32	1.98	0.45
3:A:302:EDO:H21	7:A:434:HOH:O	2.17	0.44
2:B:1:ALA:CB	7:B:413:HOH:O	2.51	0.44
1:A:62:GLY:H	5:A:316:GOL:C1	2.31	0.43
1:A:124:LEU:N	1:A:124:LEU:HD12	2.34	0.43
1:A:148:GLN:HA	4:A:314:PEG:H21	2.00	0.43
1:A:98:ASP:HB2	2:B:49:PHE:CZ	2.54	0.41
2:B:5:GLN:NE2	2:B:90[B]:CYS:H	2.18	0.41
1:A:2:GLN:HB2	1:A:95[A]:GLU:OE1	2.21	0.41
2:B:5:GLN:NE2	2:B:90[A]:CYS:H	2.19	0.41
1:A:161:VAL:HG22	1:A:172:ASN:OD1	2.22	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:ASN:OD1	1:A:145:ASN:OD1[2_775]	1.53	0.67
1:A:7:LEU:O	3:A:308:EDO:O2[2_775]	2.01	0.19
1:A:148:GLN:NE2	1:A:148:GLN:NE2[2_775]	2.04	0.16

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/202 (102%)	194 (95%)	11 (5%)	0	100	100
2	B	248/240 (103%)	242 (98%)	6 (2%)	0	100	100
All	All	453/442 (102%)	436 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/180 (103%)	178 (96%)	7 (4%)	29	8
2	B	220/210 (105%)	214 (97%)	6 (3%)	39	17
All	All	405/390 (104%)	392 (97%)	13 (3%)	35	11

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	GLU
1	A	115	GLN
1	A	125	ARG
1	A	133	SER
1	A	149	SER
1	A	165	ARG
1	A	202	SER
2	B	8	ARG
2	B	13[A]	LYS
2	B	13[B]	LYS
2	B	59	ILE
2	B	160	LYS
2	B	215	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	143	GLN
1	A	145	ASN
1	A	190	ASN
2	B	5	GLN
2	B	9	ASN
2	B	221	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

31 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	304	-	3,3,3	0.44	0	2,2,2	0.48	0
3	EDO	B	303	-	3,3,3	0.45	0	2,2,2	0.30	0
3	EDO	B	309	-	3,3,3	0.46	0	2,2,2	0.48	0
3	EDO	B	312	-	3,3,3	0.48	0	2,2,2	0.20	0
3	EDO	A	308	-	3,3,3	0.47	0	2,2,2	0.12	0
3	EDO	B	304	-	3,3,3	0.51	0	2,2,2	0.21	0
3	EDO	B	302	-	3,3,3	0.33	0	2,2,2	0.47	0
3	EDO	A	307	-	3,3,3	0.46	0	2,2,2	0.34	0
3	EDO	B	307	-	3,3,3	0.43	0	2,2,2	0.30	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	A	310	-	3,3,3	0.44	0	2,2,2	0.24	0
3	EDO	B	311	-	3,3,3	0.47	0	2,2,2	0.35	0
3	EDO	A	301	-	3,3,3	0.47	0	2,2,2	0.34	0
4	PEG	A	315	-	6,6,6	0.48	0	5,5,5	1.04	0
4	PEG	B	313	-	6,6,6	0.56	0	5,5,5	0.46	0
3	EDO	A	302	-	3,3,3	0.23	0	2,2,2	0.78	0
3	EDO	B	301	-	3,3,3	0.44	0	2,2,2	0.31	0
4	PEG	A	314	-	6,6,6	0.51	0	5,5,5	1.00	0
5	GOL	A	316	-	5,5,5	0.56	0	5,5,5	1.64	1 (20%)
3	EDO	A	309	-	3,3,3	0.47	0	2,2,2	0.22	0
3	EDO	B	306	-	3,3,3	0.32	0	2,2,2	0.68	0
3	EDO	A	313	-	3,3,3	0.43	0	2,2,2	0.36	0
3	EDO	B	305	-	3,3,3	0.30	0	2,2,2	0.50	0
3	EDO	A	311	-	3,3,3	0.37	0	2,2,2	0.45	0
3	EDO	A	305	-	3,3,3	0.49	0	2,2,2	0.19	0
3	EDO	A	312	-	3,3,3	0.37	0	2,2,2	0.59	0
3	EDO	B	310	-	3,3,3	0.44	0	2,2,2	0.28	0
3	EDO	A	303	-	3,3,3	0.32	0	2,2,2	0.56	0
6	SO4	B	315	-	4,4,4	0.49	0	6,6,6	0.18	0
3	EDO	A	306	-	3,3,3	0.33	0	2,2,2	0.44	0
3	EDO	B	308	-	3,3,3	0.53	0	2,2,2	0.24	0
5	GOL	B	314	-	5,5,5	0.54	0	5,5,5	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	304	-	-	0/1/1/1	-
3	EDO	B	303	-	-	1/1/1/1	-
3	EDO	B	309	-	-	1/1/1/1	-
3	EDO	B	312	-	-	1/1/1/1	-
3	EDO	A	308	-	-	1/1/1/1	-
3	EDO	B	304	-	-	1/1/1/1	-
3	EDO	B	302	-	-	0/1/1/1	-
3	EDO	A	307	-	-	0/1/1/1	-
3	EDO	B	307	-	-	0/1/1/1	-
3	EDO	A	310	-	-	0/1/1/1	-
3	EDO	B	311	-	-	0/1/1/1	-
3	EDO	A	301	-	-	1/1/1/1	-
4	PEG	A	315	-	-	3/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	B	313	-	-	1/4/4/4	-
3	EDO	A	302	-	-	1/1/1/1	-
3	EDO	B	301	-	-	0/1/1/1	-
4	PEG	A	314	-	-	2/4/4/4	-
5	GOL	A	316	-	-	3/4/4/4	-
3	EDO	A	309	-	-	1/1/1/1	-
3	EDO	B	306	-	-	0/1/1/1	-
3	EDO	A	313	-	-	0/1/1/1	-
3	EDO	B	305	-	-	1/1/1/1	-
3	EDO	A	311	-	-	0/1/1/1	-
3	EDO	A	305	-	-	1/1/1/1	-
3	EDO	A	312	-	-	1/1/1/1	-
3	EDO	B	310	-	-	1/1/1/1	-
3	EDO	A	303	-	-	0/1/1/1	-
3	EDO	A	306	-	-	0/1/1/1	-
3	EDO	B	308	-	-	0/1/1/1	-
5	GOL	B	314	-	-	3/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	316	GOL	O1-C1-C2	3.16	124.61	110.38

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	316	GOL	O1-C1-C2-O2
5	A	316	GOL	O1-C1-C2-C3
5	B	314	GOL	O1-C1-C2-C3
5	B	314	GOL	C1-C2-C3-O3
4	A	314	PEG	O1-C1-C2-O2
4	A	315	PEG	O1-C1-C2-O2
5	B	314	GOL	O1-C1-C2-O2
3	A	309	EDO	O1-C1-C2-O2
4	B	313	PEG	O2-C3-C4-O4
4	A	315	PEG	O2-C3-C4-O4
3	A	301	EDO	O1-C1-C2-O2
3	A	302	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	308	EDO	O1-C1-C2-O2
3	B	303	EDO	O1-C1-C2-O2
4	A	314	PEG	O2-C3-C4-O4
3	B	312	EDO	O1-C1-C2-O2
3	A	305	EDO	O1-C1-C2-O2
5	A	316	GOL	O2-C2-C3-O3
3	A	312	EDO	O1-C1-C2-O2
3	B	304	EDO	O1-C1-C2-O2
3	B	305	EDO	O1-C1-C2-O2
3	B	310	EDO	O1-C1-C2-O2
4	A	315	PEG	C1-C2-O2-C3
3	B	309	EDO	O1-C1-C2-O2

There are no ring outliers.

6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	308	EDO	0	1
4	A	315	PEG	1	0
4	B	313	PEG	1	0
3	A	302	EDO	1	0
4	A	314	PEG	2	0
5	A	316	GOL	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	202/202 (100%)	0.43	18 (8%)	15 20	15, 31, 61, 83	5 (2%)
2	B	240/240 (100%)	0.15	9 (3%)	44 52	15, 29, 50, 83	10 (4%)
All	All	442/442 (100%)	0.27	27 (6%)	27 33	15, 30, 57, 83	15 (3%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	ALA	4.7
1	A	52	SER	4.1
2	B	1	ALA	3.6
1	A	150	LYS	3.5
1	A	128	LYS	3.4
1	A	131	ASP	3.1
2	B	239	ALA	2.7
1	A	41	ASN	2.6
1	A	129	SER	2.5
1	A	28	SER	2.5
1	A	179	ASN	2.4
2	B	236[A]	TRP	2.4
1	A	39	TYR	2.4
1	A	133	SER	2.4
1	A	202	SER	2.4
2	B	240	ASP	2.4
1	A	125	ARG	2.3
1	A	115	GLN	2.3
2	B	51	VAL	2.3
1	A	132	LYS	2.2
1	A	183	PHE	2.2
1	A	53	ALA	2.1
2	B	180	ASN	2.1
1	A	126	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	19	MET	2.1
2	B	195[A]	THR	2.1
2	B	52	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	B	312	4/4	0.60	0.30	66,69,69,70	0
6	SO4	B	315	5/5	0.67	0.19	81,82,87,89	0
5	GOL	A	316	6/6	0.74	0.17	44,46,49,55	0
3	EDO	A	304	4/4	0.75	0.16	60,60,61,62	0
3	EDO	B	305	4/4	0.76	0.19	48,48,50,54	0
4	PEG	A	314	7/7	0.77	0.20	57,58,60,60	0
3	EDO	B	303	4/4	0.78	0.16	53,54,54,56	0
4	PEG	B	313	7/7	0.80	0.15	39,42,43,45	0
3	EDO	A	305	4/4	0.80	0.15	57,62,62,62	0
3	EDO	A	309	4/4	0.80	0.18	59,59,61,64	0
3	EDO	B	301	4/4	0.81	0.16	48,48,54,56	0
3	EDO	B	311	4/4	0.82	0.13	48,53,54,55	0
3	EDO	A	311	4/4	0.82	0.20	55,57,57,62	0
4	PEG	A	315	7/7	0.84	0.16	35,40,51,61	0
3	EDO	A	313	4/4	0.85	0.12	48,48,48,49	0
3	EDO	B	302	4/4	0.86	0.12	40,43,46,50	0
3	EDO	B	310	4/4	0.86	0.15	50,58,59,63	0
3	EDO	B	304	4/4	0.86	0.14	45,52,53,55	0
3	EDO	A	310	4/4	0.87	0.13	55,55,56,58	0
3	EDO	A	301	4/4	0.87	0.16	56,56,57,59	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	B	307	4/4	0.87	0.16	53,55,56,56	0
3	EDO	B	309	4/4	0.87	0.13	40,50,50,53	0
3	EDO	A	306	4/4	0.88	0.13	41,43,44,46	0
5	GOL	B	314	6/6	0.88	0.12	45,55,56,56	0
3	EDO	B	306	4/4	0.88	0.12	37,42,43,43	0
3	EDO	A	308	4/4	0.89	0.13	44,47,48,53	0
3	EDO	A	302	4/4	0.89	0.12	38,44,45,48	0
3	EDO	B	308	4/4	0.90	0.12	39,40,45,45	0
3	EDO	A	312	4/4	0.93	0.14	36,39,40,43	0
3	EDO	A	307	4/4	0.94	0.16	29,31,32,32	0
3	EDO	A	303	4/4	0.95	0.09	38,46,47,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.