



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:41 AM UTC

PDB ID : 6FUZ / pdb_00006fuz
Title : Crystal structure of the TPR domain of KLC1 in complex with the C-terminal peptide of JIP1
Authors : Pernigo, S.; Dodding, M.P.; Steiner, R.A.
Deposited on : 2018-02-28
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	4% 64% 14% • 21%
2	N	121	76% 16% • 7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kinesin light chain 1, Kinesin light chain 1, C-Jun-amino-terminal kinase-interacting protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2140	1340	381	411	8			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	201	GLY	ALA	conflict	UNP Q5UE59
A	202	SER	ALA	conflict	UNP Q5UE59
A	203	HIS	GLN	conflict	UNP Q5UE59
A	204	MET	GLN	conflict	UNP Q5UE59
A	671	THR	-	linker	UNP Q5UE59
A	672	GLY	-	linker	UNP Q5UE59
A	673	SER	-	linker	UNP Q5UE59
A	674	THR	-	linker	UNP Q5UE59
A	675	GLY	-	linker	UNP Q5UE59
A	676	SER	-	linker	UNP Q5UE59
A	677	THR	-	linker	UNP Q5UE59
A	678	GLY	-	linker	UNP Q5UE59
A	679	SER	-	linker	UNP Q5UE59
A	680	THR	-	linker	UNP Q5UE59
A	681	GLY	-	linker	UNP Q5UE59
A	682	SER	-	linker	UNP Q5UE59
A	683	THR	-	linker	UNP Q5UE59
A	684	GLY	-	linker	UNP Q5UE59
A	685	SER	-	linker	UNP Q5UE59
A	686	THR	-	linker	UNP Q5UE59
A	687	GLY	-	linker	UNP Q5UE59
A	688	SER	-	linker	UNP Q5UE59
A	689	THR	-	linker	UNP Q5UE59
A	690	GLY	-	linker	UNP Q5UE59
A	691	SER	-	linker	UNP Q5UE59
A	692	THR	-	linker	UNP Q5UE59

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	693	GLY	-	linker	UNP Q5UE59
A	694	SER	-	linker	UNP Q5UE59
A	695	THR	-	linker	UNP Q5UE59
A	696	GLY	-	linker	UNP Q5UE59
A	697	SER	-	linker	UNP Q5UE59
A	698	THR	-	linker	UNP Q5UE59
A	699	GLY	-	linker	UNP Q5UE59
A	700	SER	-	linker	UNP Q5UE59

- Molecule 2 is a protein called nanobody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	N	112	Total	C	N	O	S	0	0	0
			861	541	147	168	5			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	N	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	7	Total	O	0	0
			7	7		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	N	6	Total	O	0	0
			6	6		

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	107.47Å 90.36Å 51.70Å 90.00° 99.80° 90.00°	Depositor
Resolution (Å)	45.18 – 2.70 45.18 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.4 (45.18-2.70) 98.3 (45.18-2.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 2.69Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.233 , 0.257 0.244 , 0.283	Depositor DCC
R_{free} test set	669 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	84.9	Xtriage
Anisotropy	0.271	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 81.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3020	wwPDB-VP
Average B, all atoms (Å ²)	113.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/2174	1.52	7/2929 (0.2%)
2	N	0.67	0/880	1.18	3/1191 (0.3%)
All	All	0.76	0/3054	1.43	10/4120 (0.2%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	N	68	PHE	CA-CB-CG	6.11	119.91	113.80
1	A	312	LYS	CA-C-N	5.76	128.27	120.38
1	A	312	LYS	C-N-CA	5.76	128.27	120.38
2	N	33	TRP	N-CA-C	-5.68	101.30	110.32
1	A	242	ASP	CA-CB-CG	5.65	118.25	112.60
1	A	494	ARG	CA-C-N	5.20	131.06	121.70
1	A	494	ARG	C-N-CA	5.20	131.06	121.70
2	N	99	ASP	CA-CB-CG	5.16	117.76	112.60
1	A	254	VAL	N-CA-C	-5.07	105.76	110.53
1	A	480	PHE	N-CA-C	5.06	116.87	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2140	0	2141	27	0
2	N	861	0	824	9	0
3	N	6	0	8	1	0
4	A	7	0	0	0	0
4	N	6	0	0	0	0
All	All	3020	0	2973	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:475:ARG:HH21	1:A:487:GLU:HG3	1.63	0.64
1:A:271:TYR:HB3	1:A:310:ARG:HD2	1.80	0.62
2:N:83:MET:HE2	2:N:86:LEU:HD21	1.82	0.61
1:A:226:GLN:HB3	1:A:228:ARG:HG3	1.83	0.61
1:A:294:PRO:HG3	1:A:330:VAL:HG11	1.81	0.60
1:A:475:ARG:NH2	1:A:487:GLU:HG3	2.20	0.57
2:N:39:GLN:HB2	2:N:45:LEU:HD12	1.87	0.56
1:A:468:LYS:HE3	1:A:494:ARG:HH21	1.71	0.55
1:A:399:GLN:HG2	3:N:201:GOL:H32	1.89	0.54
1:A:370:GLN:OE1	2:N:33:TRP:HB2	2.10	0.52
1:A:406:GLU:HB3	1:A:410:ARG:HH12	1.78	0.48
1:A:336:PRO:HD3	1:A:372:LYS:HG2	1.94	0.48
1:A:343:ASN:HD21	1:A:386:ASN:ND2	2.11	0.47
1:A:251:HIS:CD2	1:A:253:ASP:H	2.32	0.47
2:N:37:VAL:HG12	2:N:47:TRP:HA	1.97	0.47
1:A:362:TYR:CE1	1:A:386:ASN:HB3	2.51	0.46
1:A:350:GLN:HG3	1:A:393:LYS:HE2	1.98	0.45
1:A:340:LYS:HG3	1:A:710:LEU:HB2	1.98	0.45
1:A:216:LEU:O	1:A:220:VAL:HG23	2.17	0.45
2:N:102:GLY:HA3	2:N:105:ARG:HD3	1.98	0.44
1:A:312:LYS:HG2	1:A:315:GLU:HB2	1.99	0.44
1:A:298:ALA:HA	1:A:327:ARG:HH11	1.82	0.44
1:A:331:LEU:HD13	1:A:337:ASP:HB2	2.00	0.44
1:A:340:LYS:HA	1:A:710:LEU:HD12	1.99	0.44
1:A:327:ARG:HH22	1:A:340:LYS:HD2	1.83	0.43
2:N:50:THR:HB	2:N:59:TYR:HB2	2.00	0.43
1:A:425:PRO:HD2	1:A:428:MET:HG3	2.00	0.43
1:A:275:ALA:HB2	1:A:306:LEU:HD21	2.00	0.43
1:A:217:HIS:CE1	1:A:260:ILE:HD11	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:421:ASP:HA	1:A:424:LYS:HE2	2.01	0.42
2:N:95:TYR:HB3	2:N:105:ARG:HB2	2.02	0.42
2:N:22:CYS:HB3	2:N:79:LEU:HB3	2.02	0.41
1:A:475:ARG:HG3	1:A:475:ARG:HH11	1.86	0.41
1:A:375:PRO:HG3	2:N:32:TYR:CE1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/337 (77%)	253 (97%)	7 (3%)	1 (0%)	30	54
2	N	110/121 (91%)	106 (96%)	3 (3%)	1 (1%)	14	35
All	All	371/458 (81%)	359 (97%)	10 (3%)	2 (0%)	24	48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	417	GLY
2	N	41	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/275 (82%)	213 (94%)	13 (6%)	18	42
2	N	90/99 (91%)	88 (98%)	2 (2%)	45	74
All	All	316/374 (84%)	301 (95%)	15 (5%)	23	51

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	216	LEU
1	A	221	ILE
1	A	226	GLN
1	A	312	LYS
1	A	315	GLU
1	A	350	GLN
1	A	357	GLU
1	A	393	LYS
1	A	416	PHE
1	A	426	ILE
1	A	433	ARG
1	A	465	THR
1	A	467	LEU
2	N	12	VAL
2	N	112	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	217	HIS
1	A	259	ASN
1	A	268	GLN
1	A	302	ASN
1	A	343	ASN
1	A	379	ASN
1	A	386	ASN
1	A	429	HIS
2	N	82	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	N	201	-	5,5,5	0.32	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	N	201	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	N	201	GOL	O1-C1-C2-C3
3	N	201	GOL	C1-C2-C3-O3
3	N	201	GOL	O2-C2-C3-O3
3	N	201	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	N	201	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/337 (79%)	0.34	12 (4%) 38 34	79, 112, 153, 180	0
2	N	112/121 (92%)	-0.10	0 100 100	82, 104, 124, 132	0
All	All	379/458 (82%)	0.21	12 (3%) 50 46	79, 108, 149, 180	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	235	LEU	3.6
1	A	219	LEU	3.1
1	A	223	TYR	3.0
1	A	216	LEU	2.9
1	A	271	TYR	2.9
1	A	229	TYR	2.5
1	A	704	PRO	2.4
1	A	236	CYS	2.3
1	A	243	LEU	2.2
1	A	221	ILE	2.1
1	A	306	LEU	2.1
1	A	214	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	N	201	6/6	0.92	0.07	101,102,103,103	0

6.5 Other polymers [i](#)

There are no such residues in this entry.