



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:27 AM UTC

PDB ID : 3FVU / pdb_00003fvu
Title : Crystal Structure of Human Kynurenine Aminotransferase I in Complex with Indole-3-acetic Acid
Authors : Han, Q.; Robinson, H.; Cai, T.; Tagle, D.A.; Li, J.
Deposited on : 2009-01-16
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	6% 88% 10% ..
1	B	422	4% 94% 5% ..

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 7386 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kynurenine--oxoglutarate transaminase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	419	Total	C	N	O	P	S	0	0	0
			3369	2175	565	609	1	19			
1	B	419	Total	C	N	O	P	S	0	0	0
			3369	2175	565	609	1	19			

- Molecule 2 is 1H-INDOL-3-YLACETIC ACID (CCD ID: IAC) (formula: $C_{10}H_9NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			13	10	1	2		
2	B	1	Total	C	N	O	0	0
			13	10	1	2		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		
4	B	1	Total	Na	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	277	Total	O	0	0
			277	277		
5	B	313	Total	O	0	0
			313	313		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kynurenine--oxoglutarate transaminase 1



- Molecule 1: Kynurenine--oxoglutarate transaminase 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	106.36Å 108.45Å 81.87Å 90.00° 114.20° 90.00°	Depositor
Resolution (Å)	29.22 – 1.55 29.22 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.22-1.55) 99.7 (29.22-1.55)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.44 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.185 , 0.210 (Not available) , 0.208	Depositor DCC
R_{free} test set	6120 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	18.9	Xtriage
Anisotropy	0.019	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7386	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IAC, GOL, LLP, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/3441	0.81	1/4671 (0.0%)
1	B	0.61	0/3441	0.79	1/4671 (0.0%)
All	All	0.61	0/6882	0.80	2/9342 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	70	PRO	N-CA-C	6.09	118.13	110.70
1	B	70	PRO	N-CA-C	5.50	117.41	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3369	0	3295	34	0
1	B	3369	0	3295	15	0
2	A	13	0	8	1	0
2	B	13	0	8	0	0
3	A	18	0	24	4	0
3	B	12	0	16	2	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
5	A	277	0	0	3	0
5	B	313	0	0	1	0
All	All	7386	0	6646	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:327:VAL:HG11	1:A:418:TRP:CE2	2.00	0.96
1:A:16:ASN:O	1:A:19:VAL:HG22	1.67	0.92
1:A:89:ASP:H	1:A:93:ASN:HD22	1.19	0.90
1:B:89:ASP:H	1:B:93:ASN:HD22	1.30	0.79
1:A:327:VAL:HG12	1:A:327:VAL:O	1.86	0.76
1:B:117:GLU:CD	1:B:174:ARG:HG3	2.17	0.69
1:A:421:GLU:O	1:A:422:LEU:HB2	1.94	0.67
1:A:150:GLN:HG2	1:A:152:GLY:H	1.63	0.63
1:B:417:LYS:O	1:B:420:VAL:HG22	2.00	0.61
1:A:350:LYS:N	1:A:350:LYS:HD3	2.18	0.59
1:B:89:ASP:H	1:B:93:ASN:ND2	1.99	0.58
1:A:163:ASP:HB3	1:A:166:GLU:HB2	1.85	0.57
1:A:327:VAL:HG11	1:A:418:TRP:CD2	2.40	0.56
1:A:416:ARG:O	1:A:420:VAL:HG22	2.07	0.55
1:A:88:ILE:O	3:A:424:GOL:H11	2.06	0.55
1:A:346:ASP:OD1	1:A:349:ARG:NH2	2.37	0.55
1:A:356:GLY:HA3	1:A:365:ARG:HD3	1.89	0.54
1:A:89:ASP:H	1:A:93:ASN:ND2	1.98	0.52
1:A:267:LYS:NZ	5:A:651:HOH:O	2.43	0.52
1:A:149:ILE:HG21	1:A:155:GLY:HA2	1.90	0.52
1:B:172:THR:OG1	1:B:174:ARG:HG2	2.10	0.52
1:A:327:VAL:O	1:A:327:VAL:CG1	2.55	0.51
1:A:279:HIS:HD2	1:B:255:LYS:NZ	2.09	0.50
1:A:255:LYS:NZ	1:B:279:HIS:HD2	2.10	0.49
1:A:252:THR:H	1:B:62:GLN:NE2	2.11	0.49
1:A:89:ASP:N	1:A:93:ASN:HD22	2.00	0.48
1:A:273:HIS:HD2	5:A:490:HOH:O	1.96	0.48
1:A:43:PRO:CB	3:A:426:GOL:H2	2.44	0.47
1:B:103:ALA:HB1	1:B:259:VAL:HG22	1.97	0.47
1:A:279:HIS:HE1	3:B:424:GOL:O3	1.99	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:273:HIS:HD2	5:B:523:HOH:O	1.99	0.46
1:B:282:THR:HG21	3:B:425:GOL:H11	1.98	0.45
1:A:182:THR:HA	1:A:183:PRO:C	2.41	0.45
1:A:391:LYS:HG3	5:A:443:HOH:O	2.16	0.45
1:A:338:TYR:CD1	1:A:402:VAL:HG21	2.51	0.45
1:B:150:GLN:HE21	1:B:150:GLN:HB2	1.63	0.45
1:A:62:GLN:NE2	1:B:252:THR:H	2.16	0.44
1:A:160:TRP:O	1:A:189:LYS:HE3	2.19	0.43
1:A:327:VAL:HG11	1:A:418:TRP:NE1	2.30	0.43
3:A:425:GOL:O1	1:B:279:HIS:HE1	2.02	0.42
1:B:182:THR:HA	1:B:183:PRO:C	2.43	0.42
1:A:18:TRP:CH2	2:A:423:IAC:H8	2.54	0.42
1:A:90:PRO:HD3	3:A:424:GOL:H32	2.01	0.42
1:A:103:ALA:HB1	1:A:259:VAL:HG22	2.02	0.42
1:A:21:PHE:CZ	1:A:380:PRO:HG3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	416/422 (99%)	406 (98%)	9 (2%)	1 (0%)	43	24
1	B	416/422 (99%)	406 (98%)	10 (2%)	0	100	100
All	All	832/844 (99%)	812 (98%)	19 (2%)	1 (0%)	48	26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	337	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/367 (100%)	360 (99%)	5 (1%)	59	33
1	B	365/367 (100%)	357 (98%)	8 (2%)	45	18
All	All	730/734 (100%)	717 (98%)	13 (2%)	51	24

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	149	ILE
1	A	166	GLU
1	A	350	LYS
1	A	413	GLU
1	A	422	LEU
1	B	4	GLN
1	B	138	ARG
1	B	150	GLN
1	B	151	ASN
1	B	174	ARG
1	B	237	GLU
1	B	279	HIS
1	B	417	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	93	ASN
1	A	109	GLN
1	A	204	GLN
1	A	273	HIS
1	A	279	HIS
1	A	325	GLN
1	A	395	HIS
1	A	409	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	4	GLN
1	B	62	GLN
1	B	93	ASN
1	B	109	GLN
1	B	150	GLN
1	B	151	ASN
1	B	273	HIS
1	B	279	HIS
1	B	283	GLN
1	B	395	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	B	247	1	23,24,25	1.72	5 (21%)	25,32,34	1.85	6 (24%)
1	LLP	A	247	1	23,24,25	1.68	4 (17%)	25,32,34	1.96	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	B	247	1	-	0/16/17/19	0/1/1/1
1	LLP	A	247	1	-	0/16/17/19	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	247	LLP	O3-C3	-4.71	1.26	1.36
1	A	247	LLP	O3-C3	-4.67	1.26	1.36
1	B	247	LLP	C2-N1	3.12	1.39	1.33
1	A	247	LLP	CE-NZ	3.00	1.53	1.46
1	A	247	LLP	C4-C4'	2.92	1.52	1.46
1	B	247	LLP	CE-NZ	2.85	1.53	1.46
1	B	247	LLP	C4-C4'	2.60	1.52	1.46
1	A	247	LLP	C2-N1	2.58	1.38	1.33
1	B	247	LLP	C6-N1	2.11	1.38	1.34

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	247	LLP	OP4-C5'-C5	6.13	120.85	109.36
1	B	247	LLP	OP4-C5'-C5	4.88	118.51	109.36
1	B	247	LLP	C5-C4-C4'	-4.67	114.26	121.47
1	A	247	LLP	C5-C4-C4'	-4.64	114.32	121.47
1	A	247	LLP	CE-NZ-C4'	-3.20	108.47	118.72
1	B	247	LLP	CE-NZ-C4'	-2.36	111.16	118.72
1	A	247	LLP	C5-C6-N1	-2.33	120.04	123.83
1	B	247	LLP	C5-C6-N1	-2.20	120.25	123.83
1	B	247	LLP	C5'-C5-C6	-2.11	115.92	119.36
1	B	247	LLP	OP3-P-OP2	2.04	115.47	107.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	425	-	5,5,5	0.35	0	5,5,5	0.61	0
2	IAC	A	423	-	14,14,14	1.39	2 (14%)	19,19,19	1.77	4 (21%)
3	GOL	A	424	-	5,5,5	0.53	0	5,5,5	0.73	0
3	GOL	A	426	-	5,5,5	0.31	0	5,5,5	0.98	0
2	IAC	B	423	-	14,14,14	1.35	2 (14%)	19,19,19	1.52	2 (10%)
3	GOL	B	424	-	5,5,5	0.32	0	5,5,5	0.53	0
3	GOL	A	425	-	5,5,5	0.37	0	5,5,5	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	425	-	-	4/4/4/4	-
2	IAC	A	423	-	-	2/4/4/4	0/2/2/2
3	GOL	A	424	-	-	2/4/4/4	-
3	GOL	A	426	-	-	0/4/4/4	-
2	IAC	B	423	-	-	2/4/4/4	0/2/2/2
3	GOL	B	424	-	-	0/4/4/4	-
3	GOL	A	425	-	-	0/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	423	IAC	C1-C7	-3.09	1.38	1.44
2	B	423	IAC	C1-C7	-2.47	1.39	1.44
2	A	423	IAC	C8-C7	2.30	1.40	1.36
2	B	423	IAC	C8-C7	2.08	1.39	1.36

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	423	IAC	C18-C17-C7	-5.27	106.84	114.93
2	B	423	IAC	C18-C17-C7	-4.63	107.82	114.93
2	A	423	IAC	C2-C1-C7	-2.79	129.69	133.85
2	A	423	IAC	C2-C1-C	2.72	121.66	118.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	423	IAC	C17-C7-C1	2.20	129.29	126.43
2	A	423	IAC	C5-C-C1	-2.10	120.15	122.19

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	424	GOL	C1-C2-C3-O3
3	B	425	GOL	C1-C2-C3-O3
3	B	425	GOL	O1-C1-C2-C3
3	A	424	GOL	O2-C2-C3-O3
3	B	425	GOL	O2-C2-C3-O3
2	A	423	IAC	C7-C17-C18-O2
2	A	423	IAC	C7-C17-C18-O3
2	B	423	IAC	C7-C17-C18-O2
2	B	423	IAC	C7-C17-C18-O3
3	B	425	GOL	O1-C1-C2-O2

There are no ring outliers.

6 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	425	GOL	1	0
2	A	423	IAC	1	0
3	A	424	GOL	2	0
3	A	426	GOL	1	0
3	B	424	GOL	1	0
3	A	425	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	418/422 (99%)	0.30	24 (5%) 29 35	12, 19, 34, 51	0
1	B	418/422 (99%)	0.11	16 (3%) 44 52	11, 18, 32, 42	0
All	All	836/844 (99%)	0.21	40 (4%) 35 43	11, 18, 33, 51	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	358	VAL	6.4
1	B	152	GLY	6.1
1	A	21	PHE	5.6
1	B	151	ASN	5.3
1	B	422	LEU	5.1
1	A	149	ILE	5.1
1	A	152	GLY	4.8
1	A	151	ASN	4.7
1	B	357	ALA	4.7
1	B	359	ASP	4.6
1	A	22	VAL	4.4
1	A	358	VAL	4.4
1	A	18	TRP	4.4
1	A	19	VAL	4.1
1	A	150	GLN	3.5
1	A	15	TYR	3.3
1	A	17	PRO	3.1
1	A	115	GLY	2.9
1	A	422	LEU	2.8
1	A	148	PRO	2.7
1	A	359	ASP	2.6
1	B	297	LEU	2.6
1	A	114	GLU	2.6
1	A	357	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	420	VAL	2.4
1	B	29	ASP	2.3
1	B	26	SER	2.3
1	A	232	LEU	2.2
1	B	149	ILE	2.2
1	A	392	HIS	2.2
1	B	19	VAL	2.2
1	A	153	GLU	2.2
1	A	87	GLU	2.2
1	A	234	GLY	2.1
1	B	24	LEU	2.1
1	A	201	SER	2.1
1	A	302	PRO	2.1
1	B	303	SER	2.1
1	B	25	ALA	2.0
1	B	22	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	A	247	24/25	0.98	0.05	12,14,17,18	0
1	LLP	B	247	24/25	0.98	0.04	11,13,16,18	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	425	6/6	0.77	0.14	25,31,32,35	0
3	GOL	A	424	6/6	0.78	0.13	30,36,36,37	0
2	IAC	A	423	13/13	0.81	0.11	20,24,25,25	0
3	GOL	A	426	6/6	0.81	0.14	25,32,35,35	0
3	GOL	B	425	6/6	0.83	0.14	27,33,33,37	0
4	NA	A	427	1/1	0.91	0.23	34,34,34,34	0
2	IAC	B	423	13/13	0.95	0.06	16,17,17,17	0
3	GOL	B	424	6/6	0.95	0.06	17,19,20,21	0
4	NA	B	426	1/1	0.97	0.13	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.