



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:16 PM UTC

PDB ID : 4FVK / pdb_00004fvk
Title : Structural and functional characterization of neuraminidase-like molecule N10 derived from bat influenza A virus
Authors : Li, Q.; Sun, X.M.; Li, Z.X.; Liu, Y.; Vavricka, C.J.; Qi, J.X.; Gao, G.F.
Deposited on : 2012-06-29
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	3% 80% 15% ...
1	B	376	7% 83% 13% ..
2	C	2	100%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6108 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	367	Total	C	N	O	S	0	0	0
			2875	1816	489	550	20			
1	B	367	Total	C	N	O	S	0	0	0
			2875	1816	489	550	20			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	74	GLY	-	expression tag	UNP H6QM75
A	75	SER	-	expression tag	UNP H6QM75
A	76	PRO	-	expression tag	UNP H6QM75
A	77	SER	-	expression tag	UNP H6QM75
A	78	ARG	-	expression tag	UNP H6QM75
A	79	SER	-	expression tag	UNP H6QM75
A	80	PRO	-	expression tag	UNP H6QM75
A	81	GLU	-	expression tag	UNP H6QM75
A	82	PHE	-	expression tag	UNP H6QM75
B	74	GLY	-	expression tag	UNP H6QM75
B	75	SER	-	expression tag	UNP H6QM75
B	76	PRO	-	expression tag	UNP H6QM75
B	77	SER	-	expression tag	UNP H6QM75
B	78	ARG	-	expression tag	UNP H6QM75
B	79	SER	-	expression tag	UNP H6QM75
B	80	PRO	-	expression tag	UNP H6QM75
B	81	GLU	-	expression tag	UNP H6QM75
B	82	PHE	-	expression tag	UNP H6QM75

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			


- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	B	1	Total	Ca	0	0
			1	1		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Zn	0	0
			3	3		
4	B	3	Total	Zn	0	0
			3	3		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	184	Total	O	0	0
			184	184		
6	B	96	Total	O	0	0
			96	96		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Neuraminidase



• Molecule 1: Neuraminidase



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	122.50Å 122.50Å 110.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.03 – 2.20 41.03 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (41.03-2.20) 99.8 (41.03-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.70 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.207 , 0.246 0.209 , 0.248	Depositor DCC
R_{free} test set	2176 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	37.0	Xtriage
Anisotropy	0.530	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6108	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	3/2944 (0.1%)	0.90	3/3990 (0.1%)
1	B	0.47	0/2944	0.81	3/3990 (0.1%)
All	All	0.57	3/5888 (0.1%)	0.86	6/7980 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	155	GLU	C-N	-7.86	1.22	1.33
1	A	458	TRP	C-N	-7.51	1.24	1.34
1	A	259	ASN	C-O	-5.77	1.16	1.24

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	163	ILE	CB-CA-C	-7.01	102.68	112.14
1	A	153	GLU	O-C-N	-5.67	116.22	122.07
1	B	379	ILE	CA-C-N	5.33	125.40	119.32
1	B	379	ILE	C-N-CA	5.33	125.40	119.32
1	B	163	ILE	CB-CA-C	-5.25	104.03	112.16
1	A	180	SER	N-CA-C	5.07	116.00	108.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2875	0	2775	56	0
1	B	2875	0	2778	39	1
2	C	28	0	25	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
5	A	28	0	26	0	0
5	B	14	0	13	0	0
6	A	184	0	0	31	0
6	B	96	0	0	17	1
All	All	6108	0	5617	96	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:247:VAL:HG22	6:B:655:HOH:O	1.41	1.19
1:A:264:ARG:CZ	6:A:723:HOH:O	1.92	1.16
1:B:395:SER:HB2	6:B:641:HOH:O	1.48	1.14
1:A:219:ARG:NH1	6:A:733:HOH:O	1.84	1.09
1:A:264:ARG:NH1	6:A:723:HOH:O	1.85	1.06
1:B:395:SER:CB	6:B:641:HOH:O	2.01	1.06
1:A:197:ARG:CA	6:A:727:HOH:O	2.04	1.04
1:B:395:SER:CA	6:B:641:HOH:O	2.09	0.98
1:A:143:HIS:ND1	6:A:742:HOH:O	2.00	0.94
1:A:459:ASP:O	6:A:690:HOH:O	1.86	0.92
1:B:394:LEU:O	6:B:641:HOH:O	1.88	0.92
1:A:451:SER:OG	6:A:738:HOH:O	1.90	0.89
1:A:143:HIS:CG	6:A:742:HOH:O	2.25	0.89
1:A:197:ARG:O	6:A:727:HOH:O	1.91	0.88
1:B:163:ILE:O	6:B:636:HOH:O	1.92	0.88
1:A:197:ARG:CB	6:A:727:HOH:O	2.25	0.85
1:A:143:HIS:HB2	6:A:742:HOH:O	1.78	0.84
1:B:324:ASP:OD2	6:B:645:HOH:O	1.95	0.84
1:A:197:ARG:HG2	6:A:727:HOH:O	1.79	0.82
1:B:454:GLN:NE2	6:B:688:HOH:O	2.10	0.82
1:A:264:ARG:CD	6:A:723:HOH:O	2.30	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:264:ARG:NE	6:A:723:HOH:O	2.08	0.77
1:A:197:ARG:HA	6:A:727:HOH:O	1.70	0.76
1:B:292:THR:OG1	6:B:627:HOH:O	2.04	0.74
1:A:219:ARG:CG	6:A:725:HOH:O	2.35	0.73
1:A:197:ARG:CG	6:A:727:HOH:O	2.37	0.71
1:A:219:ARG:HG2	6:A:725:HOH:O	1.87	0.71
1:A:264:ARG:HD2	6:A:723:HOH:O	1.92	0.67
1:A:231:ILE:HG13	1:A:287:ILE:HD13	1.77	0.66
1:B:187:LEU:HD13	1:B:188:LYS:HG3	1.78	0.66
1:A:198:ASP:OD2	6:A:724:HOH:O	2.13	0.65
1:B:231:ILE:HG13	1:B:287:ILE:HD13	1.79	0.63
1:B:395:SER:HA	6:B:641:HOH:O	1.82	0.63
1:B:394:LEU:C	6:B:641:HOH:O	2.36	0.61
1:B:165:ALA:O	1:B:170:THR:HG21	2.02	0.60
1:A:249:GLN:HA	6:A:657:HOH:O	2.02	0.59
1:A:143:HIS:CB	6:A:742:HOH:O	2.37	0.58
1:B:163:ILE:HD11	6:B:671:HOH:O	2.04	0.58
1:A:370:SER:HB3	1:A:372:ASN:OD1	2.04	0.57
1:A:246:ASN:HB3	1:A:248:ASN:HB2	1.87	0.57
1:B:359:PRO:HB3	1:B:380:PRO:HA	1.87	0.57
1:A:187:LEU:HD13	1:A:188:LYS:HG2	1.87	0.57
1:A:102:ARG:NH2	1:A:458:TRP:O	2.38	0.56
1:A:197:ARG:C	6:A:727:HOH:O	2.23	0.56
1:B:247:VAL:HG11	1:B:294:LEU:HB2	1.89	0.54
1:A:287:ILE:HG12	1:A:305:PHE:CZ	2.44	0.53
1:B:377:GLU:HG2	6:B:641:HOH:O	2.07	0.52
1:B:153:GLU:HB3	6:B:642:HOH:O	2.09	0.52
1:A:129:CYS:HB2	1:A:163:ILE:HG12	1.93	0.51
1:A:165:ALA:O	1:A:170:THR:HG21	2.10	0.50
1:B:386:GLU:HG2	6:B:635:HOH:O	2.11	0.50
1:A:129:CYS:O	1:A:163:ILE:HG13	2.11	0.50
1:B:454:GLN:H	1:B:454:GLN:CD	2.20	0.50
1:A:280:CYS:HB3	1:A:287:ILE:CD1	2.43	0.49
1:A:280:CYS:HB3	1:A:287:ILE:HD11	1.96	0.48
1:A:145:LEU:O	6:A:627:HOH:O	2.20	0.47
6:A:617:HOH:O	2:C:1:NAG:O6	2.20	0.46
1:A:330:LEU:H	1:A:330:LEU:HD12	1.80	0.46
1:A:202:PHE:HE1	1:A:204:MET:HG3	1.81	0.46
1:B:95:LYS:HG3	1:B:450:ASN:O	2.16	0.46
1:A:153:GLU:HB3	6:A:637:HOH:O	2.15	0.45
1:B:203:ILE:HG13	1:B:223:LEU:HD23	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:LEU:HB3	1:A:150:THR:HG23	1.98	0.45
1:A:454:GLN:NE2	6:A:762:HOH:O	2.48	0.45
1:B:254:ILE:HD11	1:B:268:LEU:HD21	2.00	0.44
1:A:120:ALA:HA	1:A:132:PHE:O	2.17	0.44
1:A:451:SER:CB	6:A:738:HOH:O	2.58	0.44
1:B:378:GLN:NE2	6:B:682:HOH:O	2.39	0.44
1:A:122:VAL:HG22	1:A:410:PHE:CG	2.53	0.43
1:B:143:HIS:CE1	6:B:652:HOH:O	2.71	0.43
1:B:243:ASP:OD1	1:B:244:GLY:N	2.46	0.43
1:A:95:LYS:HG3	1:A:450:ASN:O	2.19	0.43
1:A:84:TYR:O	1:A:86:ARG:NH2	2.52	0.43
1:B:163:ILE:H	1:B:163:ILE:HG13	1.65	0.42
1:B:213:ASP:OD1	1:B:214:THR:N	2.52	0.42
1:A:181:SER:HB2	1:A:226:PRO:HD2	2.02	0.42
1:B:231:ILE:O	1:B:237:CYS:HA	2.18	0.42
1:A:302:LEU:CD2	1:A:383:ILE:HG21	2.49	0.42
1:A:83:PHE:N	6:A:683:HOH:O	2.51	0.42
1:B:144:LEU:HD12	1:B:144:LEU:HA	1.86	0.42
1:B:427:GLU:OE2	1:B:441:GLN:HG3	2.19	0.42
1:A:248:ASN:ND2	6:A:661:HOH:O	2.53	0.42
1:A:359:PRO:HB3	1:A:380:PRO:HA	2.01	0.41
1:A:83:PHE:N	1:A:86:ARG:HH22	2.18	0.41
1:B:287:ILE:HG12	1:B:305:PHE:CZ	2.55	0.41
1:A:202:PHE:CE1	1:A:204:MET:HG3	2.56	0.41
1:A:188:LYS:HE3	1:A:208:GLY:HA3	2.02	0.41
1:A:219:ARG:HD3	6:A:725:HOH:O	2.20	0.41
1:A:331:THR:OG1	1:A:332:THR:N	2.54	0.41
1:B:280:CYS:HB3	1:B:287:ILE:HD11	2.03	0.41
1:A:144:LEU:HD12	1:A:144:LEU:HA	1.91	0.40
1:B:122:VAL:HG22	1:B:410:PHE:CG	2.55	0.40
1:B:206:LYS:HE2	1:B:209:GLU:HA	2.03	0.40
1:B:283:ILE:HD12	1:B:283:ILE:HA	1.95	0.40
1:B:302:LEU:HD22	1:B:383:ILE:HG21	2.03	0.40
1:B:226:PRO:HB2	1:B:278:SER:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:199:ARG:O	6:B:688:HOH:O[4_445]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	365/376 (97%)	349 (96%)	16 (4%)	0	100	100
1	B	365/376 (97%)	352 (96%)	13 (4%)	0	100	100
All	All	730/752 (97%)	701 (96%)	29 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/328 (98%)	300 (94%)	20 (6%)	16	19
1	B	320/328 (98%)	304 (95%)	16 (5%)	22	28
All	All	640/656 (98%)	604 (94%)	36 (6%)	19	24

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	PHE
1	A	90	GLN
1	A	122	VAL
1	A	126	LEU
1	A	144	LEU
1	A	145	LEU
1	A	153	GLU
1	A	163	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	187	LEU
1	A	199	ARG
1	A	223	LEU
1	A	286	LEU
1	A	287	ILE
1	A	362	THR
1	A	368	GLU
1	A	427	GLU
1	A	433	LEU
1	A	449	ILE
1	A	456	ILE
1	A	460	SER
1	B	86	ARG
1	B	110	LEU
1	B	122	VAL
1	B	126	LEU
1	B	144	LEU
1	B	163	ILE
1	B	187	LEU
1	B	223	LEU
1	B	264	ARG
1	B	278	SER
1	B	286	LEU
1	B	287	ILE
1	B	346	GLU
1	B	362	THR
1	B	449	ILE
1	B	454	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	249	GLN
1	A	258	GLN
1	A	263	GLN
1	B	90	GLN
1	B	284	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	1,2	14,14,15	0.65	0	17,19,21	0.70	0
2	NAG	C	2	2	14,14,15	0.39	0	17,19,21	1.37	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	NAG	C1-O5-C5	2.98	116.17	112.19
2	C	2	NAG	C4-C3-C2	-2.64	107.15	111.02
2	C	2	NAG	C2-N2-C7	-2.18	119.98	122.90

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	2	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	O7-C7-N2-C2
2	C	2	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

Of 11 ligands modelled in this entry, 8 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	NAG	B	505	1	14,14,15	0.36	0	17,19,21	1.55	5 (29%)
5	NAG	A	505	1	14,14,15	0.38	0	17,19,21	1.75	4 (23%)
5	NAG	A	506	1	14,14,15	0.55	0	17,19,21	1.02	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	505	1	-	2/6/23/26	0/1/1/1
5	NAG	A	505	1	-	0/6/23/26	0/1/1/1
5	NAG	A	506	1	-	5/6/23/26	0/1/1/1

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	505	NAG	C1-O5-C5	5.07	118.98	112.19
5	B	505	NAG	O5-C5-C6	3.01	113.53	107.66
5	B	505	NAG	C6-C5-C4	-2.97	105.73	113.02
5	A	505	NAG	C3-C4-C5	2.86	115.42	110.23
5	A	505	NAG	C6-C5-C4	-2.77	106.21	113.02
5	A	506	NAG	C1-O5-C5	2.61	115.69	112.19
5	B	505	NAG	C3-C4-C5	2.55	114.86	110.23
5	B	505	NAG	O5-C1-C2	-2.11	108.03	111.29
5	B	505	NAG	C1-O5-C5	2.09	114.98	112.19
5	A	505	NAG	O5-C5-C4	2.02	115.73	110.83

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	506	NAG	O5-C5-C6-O6
5	A	506	NAG	C8-C7-N2-C2
5	A	506	NAG	O7-C7-N2-C2
5	B	505	NAG	O5-C5-C6-O6
5	A	506	NAG	C4-C5-C6-O6
5	B	505	NAG	C4-C5-C6-O6
5	A	506	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	367/376 (97%)	0.05	10 (2%) 56 53	15, 28, 50, 74	0
1	B	367/376 (97%)	0.84	28 (7%) 20 17	33, 55, 87, 112	0
All	All	734/752 (97%)	0.45	38 (5%) 33 29	15, 40, 79, 112	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	83	PHE	5.0
1	B	197	ARG	3.6
1	A	83	PHE	3.6
1	B	451	SER	3.5
1	B	330	LEU	3.5
1	B	247	VAL	3.5
1	B	413(C)	LEU	3.3
1	B	251	VAL	3.2
1	A	451	SER	3.1
1	B	332	THR	3.1
1	A	249	GLN	2.8
1	B	417	CYS	2.8
1	A	343	GLU	2.8
1	B	337	CYS	2.8
1	A	452	PRO	2.7
1	B	331	THR	2.5
1	B	383	ILE	2.5
1	A	330	LEU	2.5
1	B	287	ILE	2.4
1	B	163	ILE	2.4
1	B	233	ILE	2.4
1	B	416	GLY	2.3
1	B	249	GLN	2.3
1	B	328	GLY	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	435	MET	2.2
1	A	307(A)	GLU	2.2
1	A	248	ASN	2.2
1	A	337	CYS	2.2
1	A	197	ARG	2.2
1	B	295	TRP	2.2
1	B	326	PRO	2.1
1	B	452	PRO	2.1
1	B	170	THR	2.1
1	B	285	ASN	2.1
1	B	456	ILE	2.1
1	B	187	LEU	2.0
1	B	84	TYR	2.0
1	B	302	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	C	2	14/15	0.79	0.12	45,51,56,63	0
2	NAG	C	1	14/15	0.91	0.08	36,43,46,48	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	506	14/15	0.63	0.16	48,55,64,68	0
5	NAG	A	505	14/15	0.85	0.09	38,43,50,52	0
5	NAG	B	505	14/15	0.87	0.10	39,46,53,58	0
4	ZN	B	504	1/1	0.93	0.10	61,61,61,61	0
4	ZN	A	504	1/1	0.96	0.05	49,49,49,49	0
4	ZN	B	503	1/1	0.96	0.07	86,86,86,86	0
4	ZN	A	503	1/1	0.96	0.06	53,53,53,53	0
3	CA	B	501	1/1	0.98	0.04	54,54,54,54	0
4	ZN	B	502	1/1	0.99	0.07	25,25,25,25	0
3	CA	A	501	1/1	0.99	0.03	32,32,32,32	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ZN	A	502	1/1	0.99	0.09	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.