



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:05 AM UTC

PDB ID : 6FV0 / pdb_00006fv0
Title : Crystal structure of the TPR domain of KLC1 in complex with the C-terminal peptide of torsinA
Authors : Pernigo, S.; Dodding, M.P.; Steiner, R.A.
Deposited on : 2018-02-28
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	338	7% 76% • 20%
2	F	121	2% 83% 11% • 6%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3137 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kinesin light chain 1,Torsin-1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	1	0
			2166	1359	384	415	8			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	201	GLY	-	expression tag	UNP Q5UE59
A	202	SER	-	expression tag	UNP Q5UE59
A	203	HIS	-	expression tag	UNP Q5UE59
A	204	MET	-	expression tag	UNP Q5UE59
A	671	THR	-	linker	UNP Q5UE59
A	672	GLY	-	linker	UNP Q5UE59
A	673	SER	-	linker	UNP Q5UE59
A	674	THR	-	linker	UNP Q5UE59
A	675	GLY	-	linker	UNP Q5UE59
A	676	SER	-	linker	UNP Q5UE59
A	677	THR	-	linker	UNP Q5UE59
A	678	GLY	-	linker	UNP Q5UE59
A	679	SER	-	linker	UNP Q5UE59
A	680	THR	-	linker	UNP Q5UE59
A	681	GLY	-	linker	UNP Q5UE59
A	682	SER	-	linker	UNP Q5UE59
A	683	THR	-	linker	UNP Q5UE59
A	684	GLY	-	linker	UNP Q5UE59
A	685	SER	-	linker	UNP Q5UE59
A	686	THR	-	linker	UNP Q5UE59
A	687	GLY	-	linker	UNP Q5UE59
A	688	SER	-	linker	UNP Q5UE59
A	689	THR	-	linker	UNP Q5UE59
A	690	GLY	-	linker	UNP Q5UE59
A	691	SER	-	linker	UNP Q5UE59
A	692	THR	-	linker	UNP Q5UE59
A	693	GLY	-	linker	UNP Q5UE59

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	694	SER	-	linker	UNP Q5UE59
A	695	THR	-	linker	UNP Q5UE59
A	696	GLY	-	linker	UNP Q5UE59
A	697	SER	-	linker	UNP Q5UE59
A	698	THR	-	linker	UNP Q5UE59
A	699	GLY	-	linker	UNP Q5UE59
A	700	SER	-	linker	UNP Q5UE59

- Molecule 2 is a protein called nanobody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	114	Total	C	N	O	S	0	0	0
			873	547	149	172	5			

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	36	Total	O	0	0
			36	36		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	F	54	Total	O	0	1
			55	55		

- Molecule 1: Kinesin light chain 1,Torsin-1A



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	106.04Å 89.68Å 50.99Å 90.00° 98.00° 90.00°	Depositor
Resolution (Å)	52.50 – 2.29 52.50 – 2.29	Depositor EDS
% Data completeness (in resolution range)	98.4 (52.50-2.29) 98.4 (52.50-2.29)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.29Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.208 , 0.245 (Not available) , 0.235	Depositor DCC
R_{free} test set	1070 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	41.7	Xtriage
Anisotropy	0.395	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 57.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3137	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/2201	1.41	0/2967
2	F	0.63	0/892	1.04	1/1207 (0.1%)
All	All	0.69	0/3093	1.31	1/4174 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	68	PHE	CA-CB-CG	6.09	119.89	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2166	0	2152	5	0
2	F	873	0	834	6	0
3	A	7	0	10	0	0
4	A	36	0	0	0	0
4	F	55	0	0	0	0
All	All	3137	0	2996	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:83:MET:HE2	2:F:86:LEU:HD21	1.74	0.69
1:A:220:VAL:HG11	1:A:236:CYS:SG	2.49	0.53
2:F:29:PHE:HB3	2:F:77:ASN:HD22	1.75	0.51
2:F:39:GLN:HB2	2:F:45:LEU:HD12	1.94	0.50
1:A:366:LEU:HD11	1:A:384:LYS:HG3	1.94	0.49
2:F:30:SER:O	2:F:53:THR:HB	2.16	0.45
2:F:50:THR:HB	2:F:59:TYR:HB2	1.98	0.45
2:F:22:CYS:HB3	2:F:79:LEU:HB3	1.99	0.44
1:A:346:ALA:HB1	1:A:358:VAL:HG13	1.99	0.44
1:A:426:ILE:H	1:A:426:ILE:HG13	1.76	0.41
1:A:340:LYS:HG3	1:A:710:LEU:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/338 (78%)	259 (98%)	5 (2%)	0	100	100
2	F	112/121 (93%)	111 (99%)	1 (1%)	0	100	100
All	All	376/459 (82%)	370 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/276 (83%)	223 (98%)	5 (2%)	45	65
2	F	92/99 (93%)	90 (98%)	2 (2%)	45	65
All	All	320/375 (85%)	313 (98%)	7 (2%)	45	65

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	216	LEU
1	A	306	LEU
1	A	347	LEU
1	A	422	GLU
1	A	705	LYS
2	F	45	LEU
2	F	112	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	238	GLN
1	A	268	GLN
1	A	394	GLN
1	A	412	HIS
1	A	423	ASN
2	F	77	ASN
2	F	82	GLN
2	F	84	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PEG	A	801	-	6,6,6	0.47	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	A	801	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	801	PEG	O2-C3-C4-O4
3	A	801	PEG	C1-C2-O2-C3
3	A	801	PEG	C4-C3-O2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/338 (79%)	0.70	24 (8%) 15 17	35, 68, 109, 160	0
2	F	114/121 (94%)	0.11	3 (2%) 57 59	33, 51, 75, 113	0
All	All	384/459 (83%)	0.53	27 (7%) 22 24	33, 60, 107, 160	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	704	THR	4.2
1	A	219	LEU	4.1
1	A	436	CYS	3.8
1	A	223	TYR	3.8
1	A	221	ILE	3.7
1	A	217[A]	HIS	3.2
1	A	216	LEU	3.1
1	A	215	THR	3.0
1	A	493	SER	3.0
1	A	703	PHE	3.0
2	F	104	TYR	2.9
1	A	461	PRO	2.6
1	A	486	LEU	2.6
1	A	227	GLY	2.6
1	A	702	VAL	2.6
1	A	495	LYS	2.6
1	A	426	ILE	2.6
1	A	220	VAL	2.5
1	A	462	THR	2.5
1	A	229	TYR	2.5
1	A	231	VAL	2.4
2	F	115	SER	2.4
1	A	334	ASP	2.3
1	A	214	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	F	41	PRO	2.2
1	A	224	ALA	2.2
1	A	330	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PEG	A	801	7/7	0.83	0.18	61,62,63,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.