



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:14 PM UTC

PDB ID : 6FVN / pdb_00006fvn
Title : DNA polymerase sliding clamp from Mycobacterium tuberculosis with bound P7 peptide
Authors : Martiel, I.; Andre, C.; Olieric, V.; Guichard, G.; Burnouf, D.
Deposited on : 2018-03-04
Resolution : 3.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2351 (3.18-3.10)
Clashscore	190562	2452 (3.18-3.10)
Ramachandran outliers	187476	2324 (3.18-3.10)
Sidechain outliers	187428	2324 (3.18-3.10)
RSRZ outliers	180081	2351 (3.18-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	3% 86% 7% 6%
1	B	402	3% 82% 10% 7%
1	C	402	5% 83% 9% 7%
1	D	402	5% 84% 10% 5%
2	H	6	67% 33%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	I	6	67% 33%
2	J	6	67% 17% 17%
2	K	6	17% 83% 17%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10676 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta sliding clamp.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	379	Total	C	N	O	S	0	0	0
			2659	1691	446	516	6			
1	B	372	Total	C	N	O	S	0	0	0
			2555	1637	430	482	6			
1	C	372	Total	C	N	O	S	0	0	0
			2563	1631	439	487	6			
1	D	380	Total	C	N	O	S	0	0	0
			2636	1679	441	510	6			

- Molecule 2 is a protein called P7 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	J	6	Total	C	N	O	0	0	0
			51	35	6	10			
2	K	6	Total	C	N	O	0	0	0
			48	32	6	10			
2	H	6	Total	C	N	O	0	0	0
			44	29	6	9			
2	I	6	Total	C	N	O	0	0	0
			51	35	6	10			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	25	Total	O	0	0
			25	25		
3	J	2	Total	O	0	0
			2	2		
3	B	10	Total	O	0	0
			10	10		
3	C	9	Total	O	0	0
			9	9		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	22	Total 22	O 22	0	0
3	I	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp




- Molecule 2: P7 peptide




- Molecule 2: P7 peptide



- Molecule 2: P7 peptide



- Molecule 2: P7 peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.45Å 126.76Å 171.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	101.84 – 3.14 101.84 – 3.14	Depositor EDS
% Data completeness (in resolution range)	88.0 (101.84-3.14) 88.0 (101.84-3.14)	Depositor EDS
R_{merge}	0.38	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 3.13Å)	Xtriage
Refinement program	BUSTER, PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.256 , 0.291 0.259 , 0.286	Depositor DCC
R_{free} test set	1351 reflections (4.38%)	wwPDB-VP
Wilson B-factor (Å ²)	80.3	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 54.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	10676	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ALC, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.17	0/2703	0.44	0/3705
1	B	0.13	0/2595	0.39	0/3558
1	C	0.27	1/2599 (0.0%)	0.44	2/3555 (0.1%)
1	D	0.16	0/2680	0.46	1/3680 (0.0%)
2	H	0.11	0/29	0.23	0/37
2	I	0.10	0/37	0.18	0/46
2	J	0.10	0/37	0.17	0/46
2	K	0.09	0/34	0.17	0/42
All	All	0.19	1/10714 (0.0%)	0.43	3/14669 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	195	ALA	CA-C	-5.55	1.47	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	62	TYR	N-CA-C	-7.60	104.51	114.31
1	C	60	PHE	N-CA-C	6.06	118.36	108.55
1	C	64	VAL	N-CA-C	5.52	117.17	109.55

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2659	0	2558	30	0
1	B	2555	0	2454	21	0
1	C	2563	0	2436	30	0
1	D	2636	0	2511	29	0
2	H	44	0	42	1	0
2	I	51	0	49	1	0
2	J	51	0	49	2	0
2	K	48	0	40	0	0
3	A	25	0	0	0	0
3	B	10	0	0	0	0
3	C	9	0	0	0	0
3	D	22	0	0	0	0
3	I	1	0	0	0	0
3	J	2	0	0	0	0
All	All	10676	0	10139	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:198:PRO:HA	1:C:199:ASP:HB3	1.24	1.18
1:C:197:SER:CB	1:C:198:PRO:CB	2.22	1.17
1:C:60:PHE:CD2	1:C:64:VAL:HG12	1.83	1.14
1:C:197:SER:CB	1:C:198:PRO:CA	2.26	1.12
1:D:61:ASP:OD1	1:D:62:TYR:N	1.89	1.05
1:C:197:SER:CB	1:C:198:PRO:HA	1.94	0.96
1:C:60:PHE:CG	1:C:64:VAL:HG12	2.05	0.91
1:A:197:SER:CB	1:A:198:PRO:CD	2.49	0.90
1:A:198:PRO:HB2	1:A:199:ASP:CB	2.02	0.89
1:A:198:PRO:HB2	1:A:199:ASP:CA	2.03	0.88
1:D:61:ASP:CG	1:D:62:TYR:H	1.82	0.88
1:C:60:PHE:CD2	1:C:64:VAL:CG1	2.56	0.87
1:A:198:PRO:HB2	1:A:199:ASP:C	2.03	0.83
1:C:198:PRO:CA	1:C:199:ASP:HB3	2.08	0.80
1:A:197:SER:CB	1:A:198:PRO:HD3	2.13	0.79
1:D:60:PHE:CD1	1:D:61:ASP:N	2.53	0.76
1:C:58:SER:HB3	1:C:129:LEU:HD11	1.68	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:58:SER:HB3	1:B:129:LEU:HD11	1.67	0.74
1:A:35:PRO:O	1:A:36:ALA:HB2	1.86	0.73
1:A:62:TYR:HA	1:D:39:ALA:HB2	1.71	0.72
1:D:64:VAL:HG11	1:D:130:PRO:HD2	1.72	0.70
1:D:60:PHE:HD1	1:D:61:ASP:H	1.41	0.67
1:A:228:LEU:H	1:A:229:GLY:HA2	1.59	0.67
1:C:198:PRO:HA	1:C:199:ASP:CB	2.05	0.67
1:A:198:PRO:HB2	1:A:199:ASP:O	1.96	0.63
1:D:61:ASP:CG	1:D:62:TYR:N	2.49	0.62
1:B:145:ALA:HA	1:B:215:ALA:HB1	1.82	0.61
1:B:195:ALA:HB1	1:B:200:ILE:HD11	1.82	0.61
1:A:39:ALA:HA	1:A:40:VAL:C	2.27	0.59
1:A:58:SER:HB3	1:A:129:LEU:HD11	1.86	0.57
1:A:35:PRO:O	1:A:36:ALA:CB	2.53	0.57
1:C:64:VAL:HG11	1:C:130:PRO:HD2	1.87	0.57
1:D:182:ASP:O	1:D:183:ARG:HB2	2.06	0.55
1:B:182:ASP:O	1:B:183:ARG:HB2	2.07	0.54
1:D:133:PRO:HD3	1:D:239:LEU:HD11	1.88	0.54
1:C:102:VAL:HG22	1:C:107:VAL:HG22	1.88	0.54
1:C:353:SER:HB3	1:C:366:ARG:HG3	1.90	0.53
1:D:145:ALA:HA	1:D:215:ALA:HB1	1.91	0.53
1:A:36:ALA:O	1:A:37:ARG:CB	2.57	0.53
1:D:58:SER:HB3	1:D:129:LEU:HD11	1.89	0.53
1:A:394:LEU:HD21	2:J:471:LEU:HD21	1.91	0.52
1:B:136:THR:OG1	1:B:228:LEU:HB2	2.09	0.52
1:D:306:VAL:HB	1:D:324:VAL:HG21	1.90	0.52
1:D:220:GLY:HA3	1:D:221:GLY:O	2.08	0.52
1:B:80:LEU:HD23	1:B:121:MET:HB2	1.91	0.52
1:B:354:PHE:HB3	1:B:356:PHE:CE2	2.44	0.52
1:A:182:ASP:O	1:A:183:ARG:HB2	2.08	0.52
1:D:394:LEU:HD21	2:I:471:LEU:HD21	1.91	0.52
1:A:198:PRO:CB	1:A:199:ASP:CB	2.83	0.52
1:C:85:LEU:O	1:C:89:ILE:HG12	2.10	0.52
1:A:85:LEU:O	1:A:89:ILE:HG12	2.10	0.51
1:C:80:LEU:HD23	1:C:121:MET:HB2	1.93	0.51
1:B:306:VAL:HB	1:B:324:VAL:HG21	1.93	0.51
1:B:239:LEU:HD23	1:B:252:ARG:HA	1.92	0.51
1:B:85:LEU:O	1:B:89:ILE:HG12	2.11	0.50
1:D:85:LEU:O	1:D:89:ILE:HG12	2.11	0.49
1:D:354:PHE:HB3	1:D:356:PHE:CE2	2.47	0.49
1:B:315:VAL:HG13	1:B:316:GLY:H	1.78	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:315:VAL:HG13	1:D:316:GLY:H	1.77	0.48
1:C:189:ARG:NH1	1:C:190:GLU:O	2.44	0.48
1:C:60:PHE:CB	1:C:64:VAL:HG12	2.44	0.48
1:D:80:LEU:HD23	1:D:121:MET:HB2	1.96	0.48
1:A:228:LEU:N	1:A:229:GLY:HA2	2.23	0.47
1:B:277:VAL:HG13	1:B:344:LEU:HB3	1.95	0.47
1:A:125:ASP:OD1	1:D:128:THR:HG22	2.14	0.47
1:B:196:SER:HA	1:B:197:SER:HA	1.67	0.47
1:C:263:GLN:OE1	1:C:263:GLN:N	2.42	0.47
1:A:354:PHE:HB3	1:A:356:PHE:CE2	2.50	0.46
1:B:182:ASP:OD1	1:B:185:ARG:HG2	2.16	0.46
1:A:197:SER:CB	1:A:198:PRO:HD2	2.40	0.46
1:C:182:ASP:OD1	1:C:185:ARG:HG2	2.16	0.46
1:C:197:SER:CA	1:C:198:PRO:CB	2.89	0.46
1:C:198:PRO:CB	1:C:200:ILE:H	2.29	0.45
1:C:396:MET:HG2	2:H:471:LEU:HG	1.99	0.45
1:D:343:GLY:O	1:D:347:LEU:HG	2.17	0.45
1:D:364:LEU:HD11	1:D:394:LEU:HD12	1.99	0.44
1:C:206:VAL:HG21	1:C:240:LEU:HD22	1.99	0.44
1:A:150:GLN:HB3	1:A:391:TYR:CE1	2.52	0.44
1:C:145:ALA:HA	1:C:215:ALA:HB1	1.99	0.44
1:A:61:ASP:OD1	1:A:62:TYR:N	2.48	0.44
1:A:314:ASP:OD2	1:A:315:VAL:N	2.50	0.44
1:C:186:LEU:HD21	1:C:261:PHE:HB2	1.98	0.44
1:B:102:VAL:HG22	1:B:107:VAL:HG22	2.00	0.44
1:D:20:ARG:HD3	1:D:94:PRO:O	2.18	0.44
1:A:182:ASP:OD1	1:A:185:ARG:HG2	2.18	0.43
2:J:469:ALC:HD23	2:J:469:ALC:HA	1.88	0.43
1:C:239:LEU:HD23	1:C:252:ARG:HA	1.99	0.43
1:D:182:ASP:OD1	1:D:185:ARG:HG2	2.18	0.43
1:A:136:THR:OG1	1:A:228:LEU:HB2	2.19	0.43
1:B:186:LEU:HD21	1:B:261:PHE:HB2	2.01	0.43
1:B:343:GLY:O	1:B:347:LEU:HG	2.19	0.43
1:D:314:ASP:OD2	1:D:315:VAL:N	2.52	0.43
1:A:198:PRO:CB	1:A:199:ASP:C	2.84	0.42
1:D:196:SER:HA	1:D:197:SER:HA	1.70	0.42
1:C:354:PHE:HB3	1:C:356:PHE:CE2	2.55	0.42
1:D:60:PHE:CD1	1:D:64:VAL:HG12	2.54	0.42
1:B:155:ALA:HB2	1:B:180:ALA:HB1	2.03	0.41
1:A:197:SER:O	1:A:198:PRO:O	2.37	0.41
1:C:89:ILE:HD12	1:C:116:PHE:CE2	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:306:VAL:HB	1:C:324:VAL:HG21	2.03	0.41
1:D:140:PRO:HA	1:D:223:ASP:CG	2.45	0.41
1:B:298:ARG:HE	1:B:332:THR:HG21	1.85	0.41
1:A:20:ARG:HD3	1:A:94:PRO:O	2.20	0.40
1:D:206:VAL:HG21	1:D:240:LEU:HD22	2.03	0.40
1:C:382:GLY:HA2	1:C:383:PRO:HD3	1.91	0.40
1:D:299:MET:HE3	1:D:308:LEU:HD11	2.03	0.40
1:B:20:ARG:HD3	1:B:94:PRO:O	2.21	0.40
1:A:273:ALA:HB2	1:A:326:TYR:HD1	1.86	0.40
1:B:75:SER:HA	1:B:76:PRO:HD3	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	371/402 (92%)	354 (95%)	11 (3%)	6 (2%)	7	28
1	B	364/402 (90%)	353 (97%)	10 (3%)	1 (0%)	36	64
1	C	360/402 (90%)	344 (96%)	14 (4%)	2 (1%)	21	50
1	D	376/402 (94%)	361 (96%)	14 (4%)	1 (0%)	36	64
2	H	3/6 (50%)	3 (100%)	0	0	100	100
2	I	3/6 (50%)	3 (100%)	0	0	100	100
2	J	3/6 (50%)	3 (100%)	0	0	100	100
2	K	3/6 (50%)	3 (100%)	0	0	100	100
All	All	1483/1632 (91%)	1424 (96%)	49 (3%)	10 (1%)	18	47

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	197	SER
1	A	198	PRO
1	C	197	SER
1	C	198	PRO
1	A	36	ALA
1	A	38	PRO
1	B	222	SER
1	D	223	ASP
1	A	37	ARG
1	A	35	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/316 (82%)	258 (100%)	0	100	100
1	B	236/316 (75%)	236 (100%)	0	100	100
1	C	236/316 (75%)	236 (100%)	0	100	100
1	D	249/316 (79%)	249 (100%)	0	100	100
2	H	3/4 (75%)	3 (100%)	0	100	100
2	I	4/4 (100%)	4 (100%)	0	100	100
2	J	4/4 (100%)	4 (100%)	0	100	100
2	K	3/4 (75%)	3 (100%)	0	100	100
All	All	993/1280 (78%)	993 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	33	ASN
1	C	269	HIS
1	C	336	ASN
1	D	95	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	296	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ALC	J	469	2	9,11,12	0.18	0	11,13,15	1.37	2 (18%)
2	ALC	H	469	2	9,11,12	0.20	0	11,13,15	1.24	2 (18%)
2	ALC	I	469	2	9,11,12	0.20	0	11,13,15	1.28	2 (18%)
2	ALC	K	469	2	9,11,12	0.19	0	11,13,15	1.73	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALC	J	469	2	-	0/5/14/16	0/1/1/1
2	ALC	H	469	2	-	0/5/14/16	0/1/1/1
2	ALC	I	469	2	-	1/5/14/16	0/1/1/1
2	ALC	K	469	2	-	0/5/14/16	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	K	469	ALC	CB-CA-C	-4.80	103.61	110.99
2	J	469	ALC	CB-CA-C	-3.69	105.31	110.99
2	K	469	ALC	CB-CA-N	3.07	117.19	110.48
2	I	469	ALC	CB-CA-N	3.01	117.06	110.48
2	H	469	ALC	CB-CA-N	3.01	117.05	110.48
2	I	469	ALC	CB-CA-C	-2.95	106.44	110.99
2	H	469	ALC	CB-CA-C	-2.74	106.77	110.99
2	J	469	ALC	CB-CA-N	2.62	116.21	110.48

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	I	469	ALC	CA-CB-CG-CD2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	J	469	ALC	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	379/402 (94%)	0.26	14 (3%)	45	25	37, 56, 86, 103	0
1	B	372/402 (92%)	0.46	12 (3%)	50	29	52, 76, 99, 114	0
1	C	372/402 (92%)	0.49	22 (5%)	28	14	47, 65, 96, 119	0
1	D	380/402 (94%)	0.40	21 (5%)	30	16	37, 61, 93, 117	0
2	H	4/6 (66%)	0.78	0	100	100	69, 74, 78, 83	3 (75%)
2	I	4/6 (66%)	0.51	0	100	100	71, 72, 80, 81	0
2	J	4/6 (66%)	-0.23	0	100	100	62, 65, 65, 73	0
2	K	4/6 (66%)	1.61	1 (25%)	2	1	64, 72, 80, 97	2 (50%)
All	All	1519/1632 (93%)	0.40	70 (4%)	37	19	37, 65, 95, 119	5 (0%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	39	ALA	5.7
1	D	382	GLY	4.0
1	B	42	VAL	3.8
1	B	313	ASP	3.4
1	C	239	LEU	3.4
1	D	43	LEU	3.3
1	C	269	HIS	3.2
1	A	63	GLU	3.2
1	A	387	VAL	3.1
1	B	68	GLN	3.0
1	A	62	TYR	3.0
1	D	386	ALA	3.0
1	B	237	ASP	3.0
2	K	471	LEU	2.9
1	A	41	PRO	2.8
1	C	251	THR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	52	ASP	2.8
1	D	379	ASN	2.8
1	A	40	VAL	2.8
1	B	43	LEU	2.8
1	C	237	ASP	2.8
1	C	263	GLN	2.7
1	C	383	PRO	2.7
1	A	231	GLY	2.7
1	A	355	GLY	2.7
1	C	61	ASP	2.7
1	D	38	PRO	2.6
1	D	232	PRO	2.6
1	A	313	ASP	2.6
1	C	270	THR	2.6
1	D	387	VAL	2.6
1	C	275	MET	2.6
1	D	383	PRO	2.5
1	C	315	VAL	2.5
1	D	385	PRO	2.5
1	A	385	PRO	2.4
1	A	158	ASP	2.4
1	B	70	GLY	2.4
1	C	335	PHE	2.4
1	C	357	THR	2.3
1	D	381	ASN	2.3
1	B	41	PRO	2.3
1	C	53	ASN	2.3
1	A	388	SER	2.3
1	A	219	ILE	2.2
1	D	62	TYR	2.2
1	D	37	ARG	2.2
1	B	62	TYR	2.2
1	D	41	PRO	2.2
1	C	128	THR	2.2
1	B	312	ALA	2.2
1	A	237	ASP	2.2
1	B	263	GLN	2.1
1	C	266	PRO	2.1
1	D	220	GLY	2.1
1	A	60	PHE	2.1
1	D	94	PRO	2.1
1	C	284	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	63	GLU	2.1
1	B	239	LEU	2.1
1	D	52	ASP	2.1
1	C	301	PHE	2.1
1	C	265	LEU	2.1
1	C	277	VAL	2.0
1	D	42	VAL	2.0
1	D	231	GLY	2.0
1	D	384	PHE	2.0
1	C	268	GLU	2.0
1	C	34	LEU	2.0
1	D	219	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ALC	K	469	11/12	0.59	0.20	79,89,94,98	0
2	ALC	H	469	11/12	0.82	0.21	86,89,93,99	11
2	ALC	I	469	11/12	0.83	0.14	68,71,76,83	0
2	ALC	J	469	11/12	0.89	0.13	56,62,70,75	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.