



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 12:26 AM UTC

PDB ID : 8FV6 / pdb_00008fv6
Title : E coli. CTP synthase in complex with dF-dCTP
Authors : Holyoak, T.; McLeod, M.J.; Tran, N.
Deposited on : 2023-01-18
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	545	15% 94% ..
1	BBB	545	8% 95% ..

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 8994 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CTP synthase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	535	Total	C	N	O	S	0	8	0
			4212	2665	732	792	23			
1	BBB	535	Total	C	N	O	S	0	8	0
			4208	2663	730	793	22			

- Molecule 2 is MALONIC ACID (CCD ID: MLA) (formula: $C_3H_4O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	C	O	0	0
			7	3	4		
2	BBB	1	Total	C	O	0	0
			7	3	4		


- Molecule 3 is 2'-deoxy-2',2'-difluorocytidine 5'-(tetrahydrogen triphosphate) (CCD ID: GTF) (formula: $C_9H_{14}F_2N_3O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).




3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CTP synthase



- Molecule 1: CTP synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	158.40Å 108.95Å 129.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.90 – 2.25 48.90 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.90-2.25) 100.0 (48.90-2.25)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.190 , 0.210 0.193 , 0.211	Depositor DCC
R_{free} test set	5316 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	38.9	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8994	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLA, GTF, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.01	0/4313	1.43	3/5841 (0.1%)
1	BBB	1.00	0/4309	1.43	0/5836
All	All	1.01	0/8622	1.43	3/11677 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	246	LYS	CA-C-N	5.54	123.73	120.24
1	AAA	246	LYS	C-N-CA	5.54	123.73	120.24
1	AAA	165	GLY	CA-C-O	-5.34	118.55	122.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	4212	0	4242	11	0
1	BBB	4208	0	4238	7	0
2	AAA	7	0	2	0	0
2	BBB	7	0	2	0	0
3	AAA	30	0	10	0	0
3	BBB	30	0	10	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	AAA	1	0	0	0	0
4	BBB	1	0	0	0	0
5	AAA	241	0	0	2	0
5	BBB	257	0	0	0	0
All	All	8994	0	8504	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:382:MET:HA	1:AAA:511:ALA:HB1	1.89	0.54
1:BBB:355:TYR:CD1	1:BBB:404:PHE:HB3	2.43	0.53
1:AAA:355:TYR:CD1	1:AAA:404:PHE:HB3	2.45	0.52
1:BBB:382:MET:HA	1:BBB:511:ALA:HB1	1.94	0.50
1:AAA:396:MET:HE2	1:AAA:480:GLN:CD	2.37	0.49
1:BBB:71:LEU:HD12	1:BBB:71:LEU:C	2.37	0.49
1:AAA:183:SER:OG	1:AAA:185[A]:GLU:HG2	2.14	0.48
1:AAA:177:VAL:HG12	1:AAA:186:VAL:HB	1.96	0.47
1:AAA:285:ASN:ND2	5:AAA:709:HOH:O	2.48	0.46
1:BBB:174:LEU:HG	1:BBB:209:ILE:HB	1.98	0.45
1:BBB:374:PRO:HA	1:BBB:508:TRP:O	2.19	0.42
1:AAA:458:TYR:CD1	1:AAA:463:ILE:HG21	2.55	0.41
1:AAA:114:GLN:NE2	5:AAA:718:HOH:O	2.53	0.41
1:BBB:177:VAL:HG12	1:BBB:186:VAL:HB	2.02	0.41
1:AAA:189:LYS:HB3	1:AAA:190:PRO:HD3	2.02	0.41
1:AAA:174:LEU:HG	1:AAA:209:ILE:HB	2.01	0.41
1:AAA:454:VAL:HG11	1:AAA:500:ILE:HG21	2.02	0.40
1:BBB:377:GLY:O	1:BBB:511:ALA:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	539/545 (99%)	531 (98%)	8 (2%)	0	100	100
1	BBB	539/545 (99%)	526 (98%)	13 (2%)	0	100	100
All	All	1078/1090 (99%)	1057 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	461/461 (100%)	460 (100%)	1 (0%)	87	92
1	BBB	461/461 (100%)	455 (99%)	6 (1%)	61	72
All	All	922/922 (100%)	915 (99%)	7 (1%)	73	80

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	470	ARG
1	BBB	55	ILE
1	BBB	428	VAL
1	BBB	439	MET
1	BBB	450	ASP
1	BBB	459	ASN
1	BBB	505	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GTF	AAA	602	4	28,31,31	0.60	0	39,50,50	0.68	1 (2%)
3	GTF	BBB	701	4	28,31,31	0.61	0	39,50,50	0.75	0
2	MLA	BBB	703	-	6,6,6	1.12	0	7,7,7	0.94	0
2	MLA	AAA	601	-	6,6,6	1.06	0	7,7,7	1.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GTF	AAA	602	4	-	5/22/42/42	0/2/2/2
3	GTF	BBB	701	4	-	6/22/42/42	0/2/2/2
2	MLA	BBB	703	-	-	2/4/4/4	-
2	MLA	AAA	601	-	-	4/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	AAA	602	GTF	F2-C2'-F1	2.18	108.17	105.55

There are no chirality outliers.


All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	BBB	701	GTF	PB-O3B-PG-O2G
3	BBB	701	GTF	PA-O3A-PB-O1B
2	AAA	601	MLA	O1B-C1-C2-C3
2	BBB	703	MLA	O1A-C1-C2-C3
3	AAA	602	GTF	PB-O3B-PG-O2G
2	AAA	601	MLA	O1A-C1-C2-C3
3	AAA	602	GTF	PA-O3A-PB-O1B
3	BBB	701	GTF	PB-O3A-PA-O1A
3	BBB	701	GTF	PB-O3B-PG-O3G
3	BBB	701	GTF	PA-O3A-PB-O2B
2	BBB	703	MLA	O1B-C1-C2-C3
3	AAA	602	GTF	PB-O3A-PA-O2A
2	AAA	601	MLA	C1-C2-C3-O3B
3	AAA	602	GTF	PB-O3A-PA-O1A
3	AAA	602	GTF	PA-O3A-PB-O2B
3	BBB	701	GTF	PB-O3A-PA-O2A
2	AAA	601	MLA	C1-C2-C3-O3A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	535/545 (98%)	0.74	82 (15%) 5 4	18, 49, 83, 99	8 (1%)
1	BBB	535/545 (98%)	0.52	43 (8%) 18 16	18, 47, 82, 96	8 (1%)
All	All	1070/1090 (98%)	0.63	125 (11%) 9 9	18, 48, 82, 99	16 (1%)

All (125) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	428	VAL	10.1
1	AAA	428	VAL	6.4
1	AAA	1	MET	6.1
1	BBB	1	MET	5.5
1	AAA	438	THR	5.4
1	BBB	437	GLY	5.1
1	AAA	342	LYS	4.8
1	BBB	438	THR	4.8
1	BBB	57	HIS	4.3
1	AAA	504	PRO	4.3
1	BBB	342	LYS	4.2
1	AAA	437	GLY	3.9
1	AAA	484	ALA	3.9
1	BBB	407	ASP	3.8
1	AAA	476	MET	3.6
1	AAA	481	ILE	3.5
1	BBB	451	ASP	3.5
1	BBB	101	ARG	3.4
1	AAA	446[A]	CYS	3.3
1	AAA	56	GLN	3.3
1	BBB	479	LYS	3.2
1	BBB	335	THR	3.2
1	AAA	479	LYS	3.2
1	AAA	487	ARG	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	BBB	340	ILE	3.2
1	AAA	501	ILE	3.2
1	AAA	395	ASN	3.1
1	AAA	505	ASN	3.1
1	AAA	507	PRO	3.1
1	AAA	477	LEU	3.1
1	AAA	333	VAL	3.0
1	AAA	338	VAL	3.0
1	AAA	335	THR	3.0
1	BBB	55	ILE	3.0
1	AAA	394	ALA	2.9
1	AAA	454	VAL	2.9
1	AAA	339	GLU	2.9
1	AAA	341	LEU	2.9
1	AAA	439	MET	2.9
1	AAA	456	GLN	2.9
1	BBB	505	ASN	2.9
1	AAA	375	TYR	2.8
1	AAA	542	ARG	2.8
1	BBB	456	GLN	2.8
1	AAA	393	VAL	2.8
1	AAA	449	VAL	2.8
1	BBB	493	GLY	2.8
1	AAA	57	HIS	2.8
1	AAA	290	VAL	2.8
1	BBB	495	ASP	2.7
1	AAA	397	GLU	2.7
1	AAA	540	GLN	2.7
1	AAA	470	ARG	2.7
1	BBB	267[A]	ASN	2.7
1	BBB	423	ASN	2.7
1	AAA	55	ILE	2.7
1	AAA	340	ILE	2.7
1	AAA	416	ILE	2.7
1	AAA	101	ARG	2.6
1	AAA	396	MET	2.6
1	BBB	394	ALA	2.6
1	AAA	451	ASP	2.6
1	AAA	503	VAL	2.6
1	AAA	508	TRP	2.6
1	AAA	389	TYR	2.6
1	AAA	410	TYR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	BBB	480	GLN	2.5
1	AAA	287	VAL	2.5
1	AAA	498	VAL	2.5
1	BBB	543	GLN	2.5
1	AAA	480	GLN	2.5
1	BBB	129	GLU	2.5
1	AAA	543	GLN	2.5
1	BBB	540	GLN	2.5
1	BBB	410	TYR	2.4
1	AAA	407	ASP	2.4
1	BBB	133	GLY	2.4
1	AAA	495	ASP	2.4
1	AAA	390	ALA	2.4
1	AAA	292	ILE	2.4
1	BBB	541	LYS	2.4
1	AAA	458	TYR	2.4
1	BBB	105	ARG	2.4
1	BBB	324	ASN	2.4
1	AAA	448	LEU	2.4
1	BBB	397	GLU	2.4
1	BBB	393	VAL	2.3
1	AAA	369	ARG	2.3
1	AAA	491	ARG	2.3
1	BBB	487	ARG	2.3
1	AAA	488	VAL	2.3
1	BBB	439	MET	2.3
1	AAA	105	ARG	2.3
1	BBB	427	GLU	2.3
1	AAA	460	ALA	2.2
1	AAA	483	ASP	2.2
1	BBB	504	PRO	2.2
1	AAA	531	GLY	2.2
1	AAA	185[A]	GLU	2.2
1	AAA	346	ALA	2.2
1	AAA	412	VAL	2.2
1	AAA	464	VAL	2.2
1	BBB	338	VAL	2.2
1	AAA	324	ASN	2.2
1	AAA	133	GLY	2.2
1	BBB	83	LYS	2.2
1	AAA	457	LEU	2.2
1	AAA	463	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	541	LYS	2.2
1	BBB	408	CYS	2.2
1	AAA	399	ALA	2.2
1	BBB	484	ALA	2.2
1	AAA	293	GLY	2.1
1	BBB	424	GLY	2.1
1	BBB	491	ARG	2.1
1	AAA	419	TRP	2.1
1	BBB	447	GLN	2.1
1	AAA	535	ALA	2.1
1	AAA	450	ASP	2.1
1	AAA	462	THR	2.1
1	BBB	355	TYR	2.1
1	AAA	267	ASN	2.0
1	AAA	298	TYR	2.0
1	AAA	447	GLN	2.0
1	AAA	299	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MLA	AAA	601	7/7	0.86	0.15	51,55,59,64	0
2	MLA	BBB	703	7/7	0.93	0.10	51,53,57,61	0
4	NA	AAA	603	1/1	0.96	0.09	46,46,46,46	0
4	NA	BBB	702	1/1	0.98	0.05	41,41,41,41	0
3	GTF	AAA	602	30/30	0.99	0.04	24,25,30,30	0
3	GTF	BBB	701	30/30	0.99	0.04	22,25,29,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around GTF AAA 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GTF BBB 701:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.