



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 04:58 AM UTC

PDB ID : 8FVA / pdb_00008fva
Title : E coli. CTP synthase in complex with F-araCTP
Authors : Holyoak, T.; McLeod, M.J.; Tran, N.
Deposited on : 2023-01-18
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	545	9% 91% 7%
1	BBB	545	15% 92% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MLA	AAA	601	-	-	-	X

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 8684 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CTP synthase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	534	Total	C	N	O	S	0	9	0
			4221	2671	737	791	22			
1	BBB	534	Total	C	N	O	S	0	3	0
			4183	2646	730	786	21			

- Molecule 2 is MALONIC ACID (CCD ID: MLA) (formula: $C_3H_4O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	C	O	0	0
			7	3	4		

- Molecule 3 is 4-amino-1-{2-deoxy-2-fluoro-5-O-[(R)-hydroxy{[(R)-hydroxy(phosphonooxy)phosphoryl]oxy}phosphoryl]-beta-D-arabinofuranosyl}pyrimidin-2(1H)-one (CCD ID: 1TK) (formula: $C_9H_{15}FN_3O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	AAA	1	Total	C	F	N	O	P	0	0
			29	9	1	3	13	3		
3	BBB	1	Total	C	F	N	O	P	0	0
			29	9	1	3	13	3		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	1	Total	Na	0	0
			1	1		
4	BBB	1	Total	Na	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	116	Total	O	0	0
			116	116		
5	BBB	97	Total	O	0	0
			97	97		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CTP synthase



• Molecule 1: CTP synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	158.29Å 108.07Å 128.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.81 – 2.40 48.81 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.81-2.40) 99.8 (48.81-2.40)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.205 , 0.229 0.206 , 0.227	Depositor DCC
R_{free} test set	4335 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	48.6	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 31.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8684	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 1TK, MLA, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.97	0/4314	1.39	1/5844 (0.0%)
1	BBB	0.98	0/4269	1.41	3/5783 (0.1%)
All	All	0.98	0/8583	1.40	4/11627 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	165	GLY	CA-C-O	-6.05	118.28	122.22
1	AAA	165	GLY	CA-C-O	-5.43	118.48	122.23
1	BBB	246	LYS	CA-C-N	5.33	123.60	120.24
1	BBB	246	LYS	C-N-CA	5.33	123.60	120.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	4221	0	4249	19	0
1	BBB	4183	0	4207	17	0
2	AAA	7	0	2	0	0
3	AAA	29	0	0	1	0
3	BBB	29	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	AAA	1	0	0	0	0
4	BBB	1	0	0	0	0
5	AAA	116	0	0	0	0
5	BBB	97	0	0	1	0
All	All	8684	0	8458	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:BBB:701:1TK:O4'	3:BBB:701:1TK:C1'	1.66	1.18
3:AAA:602:1TK:O4'	3:AAA:602:1TK:C1'	1.65	1.16
1:BBB:382:MET:HA	1:BBB:511:ALA:HB1	1.72	0.70
1:BBB:154:LEU:HD13	1:BBB:201:ILE:CD1	2.25	0.67
1:AAA:382:MET:HA	1:AAA:511:ALA:HB1	1.85	0.58
1:AAA:186:VAL:HG12	1:AAA:220:GLU:HG2	1.85	0.58
1:BBB:454:VAL:HG21	1:BBB:500:ILE:HG21	1.85	0.57
1:BBB:355:TYR:CD1	1:BBB:404:PHE:HB3	2.40	0.57
1:BBB:455:ARG:HG2	1:BBB:455:ARG:HH21	1.73	0.54
1:BBB:285:ASN:ND2	5:BBB:804:HOH:O	2.40	0.54
1:AAA:396:MET:HE1	1:AAA:481:ILE:HG13	1.91	0.52
1:BBB:450:ASP:HA	1:BBB:455:ARG:HD3	1.91	0.52
1:BBB:455:ARG:HH21	1:BBB:455:ARG:CG	2.22	0.51
1:AAA:177:VAL:HG12	1:AAA:186:VAL:HG22	1.92	0.51
1:AAA:11:GLY:HA3	1:AAA:194:SER:OG	2.12	0.49
1:BBB:458:TYR:CD1	1:BBB:463:ILE:HG21	2.48	0.49
1:AAA:45:ILE:O	1:AAA:93:GLY:HA3	2.13	0.48
1:BBB:154:LEU:HB3	1:BBB:201:ILE:HD12	1.96	0.48
1:AAA:32:ARG:NH2	1:AAA:258:ASP:OD1	2.46	0.48
1:AAA:86:ARG:O	1:AAA:86:ARG:HD3	2.13	0.47
1:AAA:355:TYR:CD1	1:AAA:404:PHE:HB3	2.48	0.47
1:AAA:241:VAL:HG11	1:AAA:250:LEU:CD1	2.45	0.47
1:BBB:189:LYS:HB3	1:BBB:190:PRO:HD3	1.98	0.45
1:BBB:177:VAL:HG12	1:BBB:186:VAL:HB	1.99	0.44
1:BBB:396:MET:HE1	1:BBB:481:ILE:HG13	1.99	0.44
1:AAA:52:MET:HE2	1:AAA:57[B]:HIS:CG	2.53	0.44
1:AAA:401:SER:HA	1:AAA:413:VAL:O	2.18	0.43
1:AAA:443:ALA:HB1	1:AAA:464:VAL:CG1	2.48	0.43
1:AAA:28:ILE:HD11	1:AAA:251:LEU:HB2	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:401:SER:HA	1:BBB:413:VAL:O	2.19	0.43
1:BBB:374:PRO:HA	1:BBB:508:TRP:O	2.20	0.42
1:AAA:124:LYS:HG2	1:AAA:160:MET:HG3	2.02	0.41
1:AAA:377:GLY:O	1:AAA:511:ALA:HA	2.20	0.41
1:AAA:55[B]:ILE:HG23	1:AAA:298:TYR:HB3	2.01	0.41
1:BBB:247:ILE:N	1:BBB:248:PRO:CD	2.83	0.41
1:AAA:172:MET:SD	1:AAA:209:ILE:HD11	2.61	0.41
1:BBB:449:VAL:HG21	1:BBB:488:VAL:O	2.20	0.41
1:AAA:374:PRO:HA	1:AAA:508:TRP:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	539/545 (99%)	526 (98%)	12 (2%)	1 (0%)	43	58
1	BBB	533/545 (98%)	523 (98%)	9 (2%)	1 (0%)	43	58
All	All	1072/1090 (98%)	1049 (98%)	21 (2%)	2 (0%)	43	58

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	BBB	379	CYS
1	AAA	379	CYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	461/461 (100%)	460 (100%)	1 (0%)	87	94
1	BBB	456/461 (99%)	453 (99%)	3 (1%)	76	88
All	All	917/922 (100%)	913 (100%)	4 (0%)	84	92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	186	VAL
1	BBB	388	ASP
1	BBB	450	ASP
1	BBB	455	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	1TK	BBB	701	4	29,30,30	4.60	17 (58%)	43,47,47	1.14	4 (9%)
2	MLA	AAA	601	-	6,6,6	1.40	0	7,7,7	1.02	0
3	1TK	AAA	602	4	29,30,30	4.52	17 (58%)	43,47,47	1.25	5 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1TK	BBB	701	4	-	8/22/38/38	0/2/2/2
2	MLA	AAA	601	-	-	0/4/4/4	-
3	1TK	AAA	602	4	-	5/22/38/38	0/2/2/2

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	BBB	701	1TK	C2'-C1'	-15.57	1.34	1.53
3	AAA	602	1TK	C2'-C1'	-15.47	1.34	1.53
3	BBB	701	1TK	O4'-C1'	10.40	1.66	1.42
3	AAA	602	1TK	O4'-C1'	10.12	1.65	1.42
3	AAA	602	1TK	PB-O3B	7.18	1.67	1.59
3	BBB	701	1TK	PB-O3B	7.11	1.67	1.59
3	BBB	701	1TK	O4'-C4'	-6.95	1.29	1.45
3	AAA	602	1TK	O4'-C4'	-6.85	1.29	1.45
3	BBB	701	1TK	C4-N4	6.70	1.50	1.33
3	AAA	602	1TK	C4-N4	6.61	1.49	1.33
3	BBB	701	1TK	PB-O3A	4.56	1.64	1.59
3	AAA	602	1TK	PB-O3A	4.27	1.64	1.59
3	BBB	701	1TK	O3'-C3'	-3.26	1.34	1.43
3	BBB	701	1TK	O2-C2	-3.25	1.17	1.23
3	AAA	602	1TK	O3'-C3'	-3.19	1.35	1.43
3	BBB	701	1TK	C2-N1	-3.17	1.33	1.40
3	AAA	602	1TK	O2-C2	-3.09	1.18	1.23
3	AAA	602	1TK	C2-N1	-3.09	1.33	1.40
3	BBB	701	1TK	PA-O3A	2.79	1.62	1.59
3	AAA	602	1TK	C5-C4	-2.72	1.36	1.42
3	BBB	701	1TK	C3'-C4'	2.70	1.59	1.53
3	BBB	701	1TK	C5-C4	-2.62	1.36	1.42
3	BBB	701	1TK	C6-C5	2.61	1.41	1.35
3	AAA	602	1TK	C3'-C4'	2.54	1.59	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	AAA	602	1TK	C6-C5	2.51	1.40	1.35
3	BBB	701	1TK	C2-N3	-2.51	1.31	1.36
3	BBB	701	1TK	C2'-C3'	2.51	1.56	1.52
3	AAA	602	1TK	C2'-C3'	2.48	1.56	1.52
3	AAA	602	1TK	PA-O3A	2.26	1.61	1.59
3	BBB	701	1TK	F1-C2'	2.21	1.44	1.40
3	AAA	602	1TK	C2-N3	-2.17	1.32	1.36
3	AAA	602	1TK	F1-C2'	2.11	1.44	1.40
3	AAA	602	1TK	C6-N1	-2.01	1.33	1.38
3	BBB	701	1TK	C6-N1	-2.00	1.33	1.38

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	BBB	701	1TK	C4'-O4'-C1'	-3.32	102.14	109.47
3	BBB	701	1TK	O2-C2-N3	-3.15	117.37	122.33
3	AAA	602	1TK	O2-C2-N3	-2.91	117.74	122.33
3	AAA	602	1TK	C4'-O4'-C1'	-2.90	103.05	109.47
3	BBB	701	1TK	O4'-C1'-C2'	-2.41	103.30	105.84
3	AAA	602	1TK	C2'-C1'-N1	2.38	117.93	114.27
3	AAA	602	1TK	F1-C2'-C1'	2.26	113.41	108.88
3	AAA	602	1TK	O4'-C1'-C2'	-2.10	103.62	105.84
3	BBB	701	1TK	N1-C2-N3	2.09	122.42	118.80

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	AAA	602	1TK	PB-O3B-PG-O3G
3	BBB	701	1TK	C3'-C4'-C5'-O5'
3	BBB	701	1TK	PB-O3B-PG-O1G
3	BBB	701	1TK	C5'-O5'-PA-O1A
3	AAA	602	1TK	PB-O3A-PA-O1A
3	AAA	602	1TK	PA-O3A-PB-O1B
3	BBB	701	1TK	PA-O3A-PB-O2B
3	AAA	602	1TK	PB-O3B-PG-O1G
3	AAA	602	1TK	PB-O3B-PG-O2G
3	BBB	701	1TK	PB-O3B-PG-O2G
3	BBB	701	1TK	PB-O3B-PG-O3G
3	BBB	701	1TK	O4'-C4'-C5'-O5'
3	BBB	701	1TK	PA-O3A-PB-O1B


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	BBB	701	1TK	1	0
3	AAA	602	1TK	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	534/545 (97%)	0.76	47 (8%) 15 12	24, 52, 84, 102	9 (1%)
1	BBB	534/545 (97%)	0.98	81 (15%) 5 4	24, 55, 87, 106	3 (0%)
All	All	1068/1090 (97%)	0.87	128 (11%) 9 6	24, 54, 86, 106	12 (1%)

All (128) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	428	VAL	11.6
1	BBB	1	MET	6.7
1	AAA	57[A]	HIS	6.3
1	AAA	1	MET	6.1
1	BBB	342	LYS	5.3
1	AAA	397	GLU	5.3
1	BBB	438	THR	5.2
1	AAA	240	ASP	5.2
1	BBB	57	HIS	4.9
1	AAA	438	THR	4.8
1	BBB	504	PRO	4.7
1	BBB	479	LYS	4.6
1	AAA	267	ASN	4.5
1	AAA	182	ALA	4.5
1	BBB	487	ARG	4.4
1	BBB	446	CYS	4.3
1	AAA	494	ASP	4.3
1	AAA	56[A]	GLN	4.3
1	AAA	428	VAL	4.2
1	BBB	462	THR	4.1
1	BBB	397	GLU	3.7
1	BBB	133	GLY	3.6
1	AAA	487	ARG	3.6
1	BBB	240	ASP	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	BBB	543	GLN	3.5
1	AAA	493	GLY	3.5
1	AAA	329	ASP	3.5
1	BBB	495	ASP	3.4
1	BBB	181	ALA	3.4
1	BBB	183	SER	3.3
1	BBB	182	ALA	3.3
1	BBB	451	ASP	3.2
1	BBB	395	ASN	3.1
1	AAA	125	GLU	3.0
1	BBB	324	ASN	3.0
1	BBB	505	ASN	3.0
1	BBB	456	GLN	3.0
1	AAA	504	PRO	3.0
1	BBB	502	GLU	2.9
1	AAA	335	THR	2.9
1	AAA	342	LYS	2.9
1	BBB	289	GLU	2.9
1	BBB	267	ASN	2.8
1	AAA	541	LYS	2.8
1	BBB	447	GLN	2.8
1	AAA	450	ASP	2.8
1	BBB	449	VAL	2.8
1	BBB	339	GLU	2.8
1	BBB	460	ALA	2.7
1	BBB	56	GLN	2.7
1	AAA	183	SER	2.7
1	BBB	288	SER	2.7
1	BBB	300	GLU	2.6
1	BBB	493	GLY	2.6
1	BBB	492	SER	2.6
1	AAA	213	ASP	2.6
1	BBB	101	ARG	2.6
1	AAA	147	ASP	2.6
1	BBB	470	ARG	2.6
1	BBB	481	ILE	2.6
1	AAA	492	SER	2.6
1	BBB	476	MET	2.5
1	AAA	55[A]	ILE	2.5
1	BBB	480	GLN	2.5
1	BBB	340	ILE	2.5
1	AAA	408	CYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	BBB	542	ARG	2.5
1	AAA	300	GLU	2.4
1	BBB	541	LYS	2.4
1	BBB	484	ALA	2.4
1	BBB	335	THR	2.4
1	AAA	451	ASP	2.4
1	AAA	395	ASN	2.4
1	BBB	384	VAL	2.4
1	AAA	421	ASP	2.4
1	BBB	494	ASP	2.4
1	BBB	454	VAL	2.4
1	BBB	489	ALA	2.4
1	BBB	464	VAL	2.4
1	BBB	185	GLU	2.4
1	BBB	483	ASP	2.3
1	AAA	289	GLU	2.3
1	BBB	538	GLU	2.3
1	BBB	51	THR	2.3
1	BBB	452	SER	2.3
1	BBB	478	LEU	2.3
1	AAA	479	LYS	2.3
1	BBB	450	ASP	2.3
1	AAA	393	VAL	2.3
1	AAA	505	ASN	2.2
1	BBB	345	ASP	2.2
1	BBB	338	VAL	2.2
1	BBB	179	TYR	2.2
1	AAA	543	GLN	2.2
1	BBB	417	THR	2.2
1	AAA	407	ASP	2.2
1	AAA	339	GLU	2.2
1	AAA	480	GLN	2.2
1	AAA	476	MET	2.1
1	BBB	448	LEU	2.1
1	BBB	416	ILE	2.1
1	BBB	463	ILE	2.1
1	AAA	129	GLU	2.1
1	BBB	427	GLU	2.1
1	BBB	52	MET	2.1
1	AAA	214	ARG	2.1
1	BBB	167	GLU	2.1
1	BBB	461	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	BBB	535	ALA	2.1
1	BBB	372	ASN	2.1
1	BBB	491	ARG	2.1
1	BBB	42	ASP	2.1
1	BBB	184	GLY	2.1
1	AAA	220	GLU	2.1
1	BBB	98	ASP	2.1
1	BBB	213	ASP	2.1
1	AAA	439	MET	2.1
1	BBB	455	ARG	2.0
1	AAA	301	LEU	2.0
1	AAA	133	GLY	2.0
1	BBB	531	GLY	2.0
1	AAA	409	LYS	2.0
1	BBB	368	ALA	2.0
1	BBB	132	GLU	2.0
1	AAA	340	ILE	2.0
1	BBB	439	MET	2.0
1	AAA	355	TYR	2.0
1	BBB	410	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MLA	AAA	601	7/7	0.67	0.48	102,112,116,117	0
4	NA	BBB	702	1/1	0.79	0.23	77,77,77,77	0
4	NA	AAA	603	1/1	0.86	0.14	73,73,73,73	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	1TK	BBB	701	29/29	0.93	0.13	38,45,52,53	29
3	1TK	AAA	602	29/29	0.94	0.11	33,43,48,49	29

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 1TK BBB 701:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 1TK AAA 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.