



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:00 PM UTC

PDB ID : 8FVN / pdb_00008fvn
Title : PCSK9 in complex with an inhibitor
Authors : Xu, M.; Chopra, R.
Deposited on : 2023-01-19
Resolution : 2.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4348 (2.74-2.70)
Clashscore	190562	4665 (2.74-2.70)
Ramachandran outliers	187476	4584 (2.74-2.70)
Sidechain outliers	187428	4585 (2.74-2.70)
RSRZ outliers	180081	4348 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	152	
2	B	540	
3	L	11	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4431 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	92	Total	C	N	O	S	0	0	0
			740	474	133	131	2			

- Molecule 2 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	477	Total	C	N	O	S	0	0	0
			3537	2185	651	670	31			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	474	ILE	VAL	variant	UNP Q8NBP7
B	670	GLU	GLY	variant	UNP Q8NBP7

- Molecule 3 is a protein called MCR-ALO-MAA-ALA-MAA-NLE-7T2-SER-7T2-ALA-NH2 inhibitor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	L	11	Total	C	Cl	N	O	S	0	0	1
			74	49	2	10	12	1			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	19	Total	O	0	0
			19	19		
4	B	59	Total	O	0	0
			59	59		
4	L	2	Total	O	0	0
			2	2		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.42Å 70.50Å 149.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.15 – 2.71 47.15 – 2.71	Depositor EDS
% Data completeness (in resolution range)	99.2 (47.15-2.71) 99.2 (47.15-2.71)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.34 (at 2.73Å)	Xtriage
Refinement program	PHENIX 1	Depositor
R, R_{free}	0.181 , 0.257 0.184 , 0.258	Depositor DCC
R_{free} test set	954 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	34.0	Xtriage
Anisotropy	0.016	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4431	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAA, NLE, ALO, NH2, MCR, 7T2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/757	0.50	0/1023
2	B	0.37	0/3599	0.54	1/4882 (0.0%)
3	L	0.47	0/13	2.11	1/13 (7.7%)
All	All	0.38	0/4369	0.55	2/5918 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	4	ALA	CA-C-O	-5.54	111.39	120.80
2	B	179	LEU	N-CA-C	-5.22	106.88	113.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	740	0	750	3	0
2	B	3537	0	3451	39	0
3	L	74	0	46	2	0
4	A	19	0	0	0	0
4	B	59	0	0	1	0
4	L	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4431	0	4247	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:523:ILE:HD13	2:B:648:TYR:HB3	1.79	0.63
2:B:551:HIS:HB3	2:B:587:GLN:HG3	1.84	0.60
2:B:176:GLY:HA3	2:B:398:MET:HE3	1.86	0.58
2:B:163:PRO:HG2	2:B:166:TYR:HB3	1.85	0.57
2:B:345:PRO:HD3	2:B:424:ILE:HG23	1.85	0.57
2:B:469:ARG:HD2	2:B:515:PHE:HA	1.87	0.56
2:B:638:LEU:HB2	2:B:673:THR:HB	1.88	0.55
2:B:351:LEU:HD22	3:L:5:MAA:HB3	1.88	0.54
2:B:488:SER:OG	2:B:495:ARG:NH1	2.43	0.52
2:B:549:ARG:CA	2:B:596:ILE:HD11	2.41	0.51
2:B:568:VAL:HG12	2:B:569:GLU:H	1.75	0.51
2:B:343:ASP:HB3	2:B:423:VAL:HG13	1.91	0.51
2:B:186:ASP:OD2	2:B:230:LEU:HD12	2.12	0.50
2:B:418:PHE:CE2	2:B:445:PRO:HB3	2.49	0.48
2:B:418:PHE:CD2	2:B:445:PRO:HB3	2.49	0.47
2:B:490:SER:HB2	2:B:520:VAL:HG12	1.97	0.47
2:B:200:VAL:HG22	2:B:247:MET:HB2	1.97	0.47
2:B:464:HIS:CD2	2:B:466:GLY:H	2.33	0.46
2:B:622:VAL:HG13	2:B:656:VAL:HB	1.97	0.46
2:B:185:LEU:HD11	2:B:271:ILE:HD11	1.97	0.45
2:B:185:LEU:HD22	2:B:287:LEU:HD21	1.98	0.45
1:A:97:ARG:O	1:A:101:GLN:HG3	2.17	0.44
1:A:82:LEU:HG	1:A:121:GLY:HA2	2.00	0.44
2:B:557:HIS:CD2	2:B:602:HIS:HB2	2.53	0.43
2:B:311:LEU:HB2	2:B:333:VAL:HG22	2.01	0.43
2:B:468:THR:HG23	4:B:749:HOH:O	2.18	0.43
2:B:194:ARG:HG2	2:B:376:SER:O	2.18	0.42
2:B:183:TYR:HA	2:B:248:ARG:O	2.19	0.42
2:B:474:ILE:HD11	2:B:510:ARG:NH2	2.34	0.42
3:L:2:ALO:HA	3:L:3:MAA:HA	1.54	0.42
2:B:395:ILE:HG22	2:B:399:MET:HE3	2.02	0.42
2:B:585:PRO:HB2	2:B:586:ASN:H	1.67	0.42
1:A:144:GLU:OE2	2:B:294:SER:OG	2.32	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:312:VAL:HG12	2:B:389:ALA:HB1	2.01	0.41
2:B:549:ARG:HA	2:B:596:ILE:HD11	2.02	0.41
2:B:485:SER:HB3	2:B:633:THR:O	2.21	0.41
2:B:418:PHE:O	2:B:443:ALA:HB3	2.21	0.41
2:B:199:ARG:HD3	2:B:199:ARG:HA	1.83	0.41
2:B:464:HIS:NE2	2:B:519:GLY:HA3	2.35	0.40
2:B:558:VAL:HG12	2:B:601:CYS:O	2.21	0.40
2:B:398:MET:HE2	2:B:398:MET:HB3	1.97	0.40
2:B:605:GLY:O	2:B:681:SER:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	90/152 (59%)	87 (97%)	3 (3%)	0	100	100
2	B	457/540 (85%)	446 (98%)	8 (2%)	3 (1%)	18	39
3	L	3/11 (27%)	2 (67%)	1 (33%)	0	100	100
All	All	550/703 (78%)	535 (97%)	12 (2%)	3 (0%)	24	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	244	GLY
2	B	245	ALA
2	B	280	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	79/127 (62%)	78 (99%)	1 (1%)	61	81
2	B	381/431 (88%)	364 (96%)	17 (4%)	24	50
3	L	1/1 (100%)	1 (100%)	0	100	100
All	All	461/559 (82%)	443 (96%)	18 (4%)	28	56

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	ASP
2	B	197	GLU
2	B	278	GLN
2	B	283	LEU
2	B	381	SER
2	B	401	SER
2	B	423	VAL
2	B	449	HIS
2	B	470	MET
2	B	501	GLU
2	B	550	VAL
2	B	557	HIS
2	B	567	GLU
2	B	568	VAL
2	B	606	LEU
2	B	623	THR
2	B	645	LEU
2	B	680	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	GLN
2	B	557	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ALO	L	2	3	5,6,7	0.65	0	5,7,9	1.54	1 (20%)
3	7T2	L	9	3	12,13,14	0.92	0	15,16,18	1.81	4 (26%)
3	7T2	L	7	3	12,13,14	1.16	0	15,16,18	1.40	3 (20%)
3	NLE	L	6	3	6,7,8	1.02	0	2,7,9	0.16	0
3	MAA	L	3	3	4,5,6	0.80	0	2,5,7	1.25	0
3	MAA	L	5	3	4,5,6	0.61	0	2,5,7	1.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ALO	L	2	3	-	4/5/6/8	-
3	7T2	L	9	3	-	0/5/8/10	0/1/1/1
3	7T2	L	7	3	-	0/5/8/10	0/1/1/1
3	NLE	L	6	3	-	0/5/6/8	-
3	MAA	L	3	3	-	0/2/4/6	-
3	MAA	L	5	3	-	1/2/4/6	-

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	9	7T2	CG-CB-CA	-3.50	108.53	113.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	9	7T2	O-C-CA	-3.22	116.49	124.77
3	L	9	7T2	CE1-CD1-CG	-2.77	117.36	121.00
3	L	9	7T2	CD2-CG-CD1	2.63	122.15	118.23
3	L	7	7T2	CG-CB-CA	-2.59	109.83	113.51
3	L	2	ALO	O-C-CA	-2.56	118.19	124.77
3	L	7	7T2	CD2-CG-CD1	2.27	121.61	118.23
3	L	7	7T2	CE1-CD1-CG	-2.05	118.30	121.00

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	L	2	ALO	C-CA-CB-CG2
3	L	2	ALO	C-CA-CB-OG1
3	L	2	ALO	O-C-CA-CB
3	L	5	MAA	CB-CA-N-CM
3	L	2	ALO	N-CA-CB-CG2

There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	L	2	ALO	1	0
3	L	3	MAA	1	0
3	L	5	MAA	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	92/152 (60%)	-0.40	0 100 100	12, 19, 39, 53	0
2	B	477/540 (88%)	-0.11	22 (4%) 37 34	10, 22, 59, 95	0
3	L	3/11 (27%)	-0.30	0 100 100	16, 16, 19, 20	0
All	All	572/703 (81%)	-0.16	22 (3%) 44 40	10, 22, 56, 95	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	571	LEU	4.9
2	B	448	THR	3.7
2	B	515	PHE	3.7
2	B	177	GLY	3.7
2	B	449	HIS	3.6
2	B	616	PRO	3.5
2	B	178	SER	3.3
2	B	176	GLY	3.2
2	B	585	PRO	2.8
2	B	569	GLU	2.8
2	B	568	VAL	2.8
2	B	553	HIS	2.5
2	B	452	GLY	2.5
2	B	642	SER	2.5
2	B	557	HIS	2.5
2	B	643	HIS	2.4
2	B	628	GLU	2.4
2	B	220	ALA	2.2
2	B	602	HIS	2.2
2	B	604	PRO	2.1
2	B	167	ARG	2.1
2	B	531	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ALO	L	2	7/8	0.87	0.11	20,24,29,42	0
3	7T2	L	9	13/14	0.88	0.10	10,15,19,41	0
3	7T2	L	7	13/14	0.91	0.09	14,19,23,56	0
3	MAA	L	5	6/7	0.95	0.07	13,13,18,19	0
3	MAA	L	3	6/7	0.96	0.08	18,19,22,23	0
3	NLE	L	6	8/9	0.96	0.08	13,17,20,22	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.