



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:28 PM UTC

PDB ID : 1FWC / pdb_00001fwc
Title : KLEBSIELLA AEROGENES UREASE, C319A VARIANT AT PH 8.5
Authors : Pearson, M.A.; Karplus, P.A.
Deposited on : 1997-04-23
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	100	 95% 5%
2	B	106	 81% 13% • 5%
3	C	567	 86% 12% •

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6071 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UREASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	0	0	0
			776	491	134	146	5			

- Molecule 2 is a protein called UREASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	101	Total	C	N	O	S	0	0	0
			784	496	150	135	3			

- Molecule 3 is a protein called UREASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	566	Total	C	N	O	S	0	0	0
			4228	2652	741	813	22			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	217	KCX	LYS	modified residue	UNP P18314
C	319	ALA	CYS	engineered mutation	UNP P18314

- Molecule 4 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total	Ni	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	28	Total	O	0	0
			28	28		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	23	Total 23	O 23	0	0
5	C	230	Total 230	O 230	0	0

- Molecule 1: UREASE

```

graph LR
    M1[M1] --- K10[K10]
    K10 --- M44[M44]
    M44 --- V53[V53]
    V53 --- M57[M57]
    M57 --- I100[I100]
    style M1 fill:#f00
    style K10 fill:#00f
    style M44 fill:#00f
    style V53 fill:#00f
    style M57 fill:#00f
    style I100 fill:#00f
  
```

Chain B:  81% 13% 5%

M1		G4		H7		S38		F42		K50		R53		E72		Q75		A83		R88		A89		V90		V97		P100		L101		GLU		VAL		ASN		ASP		GLU
----	--	----	--	----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	------	--	------	--	-----	--	-----	--	-----	--	-----	--	-----

Chain C: 86% 12% .

[illegible]

4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	170.80Å 170.80Å 170.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.00 10.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.0 (10.00-2.00) 95.0 (10.00-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.00Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.168 , (Not available) 0.159 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 62.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.023 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6071	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/787	0.80	0/1061
2	B	0.38	0/804	0.88	2/1087 (0.2%)
3	C	0.44	0/4300	1.01	22/5859 (0.4%)
All	All	0.43	0/5891	0.97	24/8007 (0.3%)

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	312	HIS	N-CA-C	10.30	122.59	111.36
3	C	436	SER	N-CA-C	-8.93	98.56	109.72
3	C	173	GLY	CA-C-N	8.50	128.08	119.24
3	C	173	GLY	C-N-CA	8.50	128.08	119.24
3	C	299	ASN	N-CA-C	7.92	127.31	109.81
3	C	283	ASP	N-CA-C	7.54	122.21	112.26
3	C	298	THR	N-CA-C	-7.21	99.79	110.46
3	C	247	SER	N-CA-C	7.17	119.07	110.19
3	C	275	GLY	N-CA-C	7.13	121.21	113.58
3	C	50	VAL	N-CA-C	6.35	117.92	111.00
3	C	361	SER	N-CA-C	6.16	119.75	110.64
3	C	409	ILE	N-CA-C	6.03	116.81	110.72
3	C	172	PRO	N-CA-C	5.68	119.42	110.50
3	C	562	GLN	N-CA-C	5.53	119.51	112.87
3	C	245	LEU	N-CA-C	5.44	118.26	109.40
3	C	355	SER	N-CA-C	5.39	120.01	113.38
3	C	299	ASN	CA-C-N	-5.34	114.11	119.56
3	C	299	ASN	C-N-CA	-5.34	114.11	119.56
3	C	201	GLN	CA-C-N	5.25	125.30	119.32
3	C	201	GLN	C-N-CA	5.25	125.30	119.32
2	B	42	PHE	N-CA-C	5.20	117.62	111.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	38	SER	N-CA-C	5.12	116.86	111.28
3	C	467	THR	N-CA-C	5.06	120.98	109.81
3	C	565	PHE	N-CA-C	5.03	118.01	109.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	776	0	807	2	0
2	B	784	0	775	11	0
3	C	4228	0	4176	35	0
4	C	2	0	0	0	0
5	A	28	0	0	0	0
5	B	23	0	0	2	0
5	C	230	0	0	2	0
All	All	6071	0	5758	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:545:VAL:O	3:C:548:GLU:HG2	1.90	0.72
3:C:297:SER:OG	3:C:349:HIS:HE1	1.83	0.61
3:C:317:MET:HE3	3:C:333:ALA:HB2	1.83	0.60
3:C:300:PRO:HD3	3:C:365:GLY:HA2	1.84	0.59
3:C:236:ALA:O	3:C:518:ARG:HD2	2.04	0.58
3:C:300:PRO:HG2	5:C:703:HOH:O	2.03	0.57
3:C:218:ILE:HD11	3:C:232:ALA:CB	2.36	0.55
3:C:218:ILE:HD13	3:C:243:VAL:HG13	1.90	0.54
3:C:201:GLN:HE21	3:C:203:ASP:H	1.54	0.54
3:C:460:ASP:HB3	3:C:463:ALA:HB2	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:72:GLU:H	2:B:75:GLN:NE2	2.06	0.53
3:C:298:THR:CG2	3:C:360:ASP:HB2	2.40	0.52
2:B:100:PRO:O	2:B:101:LEU:HB2	2.09	0.52
1:A:53:VAL:O	1:A:57:MET:HG3	2.11	0.51
3:C:317:MET:HE3	3:C:333:ALA:CB	2.40	0.51
3:C:296:SER:HB3	3:C:356:LEU:HB2	1.94	0.49
3:C:363:ALA:O	3:C:364:MET:HB2	2.12	0.49
2:B:4:GLY:HA2	3:C:22:ARG:O	2.13	0.48
2:B:90:VAL:HB	2:B:97:VAL:HG11	1.96	0.47
1:A:10:LYS:HD2	1:A:44:MET:SD	2.54	0.47
3:C:36:LEU:HD23	3:C:97:GLY:HA2	1.96	0.47
2:B:83:ALA:HB1	2:B:88:ARG:HH21	1.80	0.47
2:B:50:LYS:HB2	5:B:114:HOH:O	2.15	0.47
3:C:75:ILE:O	3:C:81:ILE:HA	2.16	0.46
3:C:395:ASN:C	3:C:395:ASN:HD22	2.23	0.46
3:C:65:CYS:HA	3:C:111:PRO:HG2	1.97	0.46
3:C:246:HIS:CE1	3:C:278:GLY:HA3	2.51	0.46
3:C:395:ASN:HD22	3:C:397:ASN:H	1.64	0.45
2:B:89:ALA:HA	2:B:100:PRO:HA	1.98	0.45
3:C:507:LEU:HG	5:C:672:HOH:O	2.16	0.45
3:C:237:ASP:OD1	3:C:518:ARG:HD3	2.18	0.44
2:B:100:PRO:O	2:B:101:LEU:CB	2.65	0.43
3:C:246:HIS:C	3:C:246:HIS:CD2	2.97	0.42
3:C:271:PHE:O	3:C:272:HIS:C	2.61	0.42
3:C:188:PRO:O	3:C:449:LYS:HD2	2.20	0.42
3:C:355:SER:HB2	3:C:356:LEU:HD12	2.02	0.42
3:C:418:ALA:O	3:C:428:LYS:HE2	2.20	0.42
2:B:90:VAL:HB	2:B:97:VAL:CG1	2.50	0.41
2:B:7:HIS:HB3	3:C:20:LYS:HB2	2.02	0.41
3:C:135:ILE:HG13	3:C:154:MET:HB3	2.02	0.41
2:B:53:ARG:HD3	5:B:113:HOH:O	2.19	0.41
3:C:173:GLY:O	3:C:177:ILE:HG13	2.21	0.41
3:C:419:HIS:H	3:C:419:HIS:CD2	2.39	0.41
3:C:442:VAL:HG12	3:C:443:LYS:HG3	2.02	0.41
3:C:177:ILE:HD13	3:C:213:VAL:HG13	2.02	0.41
3:C:285:ILE:HD11	3:C:348:LEU:HD12	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/100 (98%)	98 (100%)	0	0	100	100
2	B	99/106 (93%)	94 (95%)	5 (5%)	0	100	100
3	C	563/567 (99%)	533 (95%)	27 (5%)	3 (0%)	24	21
All	All	760/773 (98%)	725 (95%)	32 (4%)	3 (0%)	30	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	360	ASP
3	C	364	MET
3	C	481	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/85 (100%)	84 (99%)	1 (1%)	63	70
2	B	78/83 (94%)	77 (99%)	1 (1%)	61	68
3	C	441/442 (100%)	427 (97%)	14 (3%)	34	35
All	All	604/610 (99%)	588 (97%)	16 (3%)	40	44

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	ILE
2	B	97	VAL
3	C	179	ARG
3	C	199	VAL
3	C	246	HIS
3	C	282	PRO
3	C	316	LEU
3	C	332	PHE
3	C	349	HIS
3	C	392	THR
3	C	395	ASN
3	C	449	LYS
3	C	465	ILE
3	C	507	LEU
3	C	508	ARG
3	C	520	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	GLN
2	B	12	GLN
2	B	16	ASN
2	B	29	HIS
2	B	75	GLN
3	C	142	GLN
3	C	201	GLN
3	C	349	HIS
3	C	362	GLN
3	C	395	ASN
3	C	419	HIS
3	C	469	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	KCX	C	217	3,4	10,11,12	0.95	1 (10%)	6,12,14	1.50	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KCX	C	217	3,4	-	0/9/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	217	KCX	OQ1-CX	2.19	1.25	1.21

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	217	KCX	OQ1-CX-NZ	-3.30	119.91	124.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/100 (100%)	-0.96	0 100 100	8, 14, 31, 44	0
2	B	101/106 (95%)	-0.53	1 (0%) 79 79	13, 22, 37, 51	0
3	C	565/567 (99%)	-0.92	0 100 100	7, 14, 30, 51	0
All	All	766/773 (99%)	-0.87	1 (0%) 92 91	7, 15, 33, 51	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	101	LEU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	KCX	C	217	12/13	0.97	0.04	7,12,13,14	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NI	C	574	1/1	0.99	0.05	17,17,17,17	1
4	NI	C	575	1/1	0.99	0.07	17,17,17,17	1

6.5 Other polymers [i](#)

There are no such residues in this entry.