



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:39 AM UTC

PDB ID : 1FWE / pdb_00001fwe
Title : KLEBSIELLA AEROGENES UREASE, C319A VARIANT WITH ACETO-HYDROXAMIC ACID (AHA) BOUND
Authors : Pearson, M.A.; Karplus, P.A.
Deposited on : 1997-04-23
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	100	93% 7%
2	B	106	77% 16% • 5%
3	C	567	80% 15% • •

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5967 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UREASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	0	0	0
			776	491	134	146	5			

- Molecule 2 is a protein called UREASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	101	Total	C	N	O	S	0	0	0
			784	496	150	135	3			

- Molecule 3 is a protein called UREASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	552	Total	C	N	O	S	0	0	0
			4123	2587	723	791	22			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	217	KCX	LYS	modified residue	UNP P18314
C	319	ALA	CYS	conflict	UNP P18314

- Molecule 4 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total	Ni	0	0
			2	2		

- Molecule 5 is ACETOHYDROXAMIC ACID (CCD ID: HAE) (formula: C₂H₅NO₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	C	N	O	0	0
			5	2	1	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	28	Total	O	0	0
			28	28		
6	B	23	Total	O	0	0
			23	23		
6	C	226	Total	O	0	0
			226	226		

- Molecule 1: UREASE

Chain B:  77% 16% 5%

M1	G4	H7	N16	T21	S38	H39	F42	R53	R60	E72	Q75	L81	V82	A83	R88	A89	V90	V97	P100	L101	GLU	VAL	ASN	ASP	GU
----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	-----	-----	-----	-----	----

Chain C:  80% 15% . . .

[illegible]

4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	170.80Å 170.80Å 170.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.00 10.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	92.0 (10.00-2.00) 91.2 (10.00-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 2.00Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.204 , (Not available) 0.199 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	13.3	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 66.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.036 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5967	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, NI, HAE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/787	0.88	0/1061
2	B	0.41	0/804	0.95	2/1087 (0.2%)
3	C	0.49	0/4191	1.04	31/5706 (0.5%)
All	All	0.48	0/5782	1.01	33/7854 (0.4%)

There are no bond length outliers.

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	436	SER	N-CA-C	-8.62	99.20	110.07
3	C	173	GLY	CA-C-N	7.74	127.29	119.24
3	C	173	GLY	C-N-CA	7.74	127.29	119.24
3	C	299	ASN	N-CA-C	7.38	126.13	109.81
3	C	283	ASP	N-CA-C	7.10	121.64	112.26
3	C	312	HIS	N-CA-C	7.06	121.15	112.54
3	C	50	VAL	N-CA-C	6.76	118.37	111.00
3	C	290	HIS	N-CA-C	6.64	118.17	109.93
2	B	38	SER	N-CA-C	6.58	119.29	111.33
3	C	247	SER	N-CA-C	6.46	118.20	110.19
3	C	298	THR	N-CA-C	-6.42	101.02	110.52
3	C	562	GLN	N-CA-C	6.17	119.91	112.38
3	C	355	SER	N-CA-C	6.14	120.93	113.38
3	C	299	ASN	CA-C-N	-6.13	113.37	119.56
3	C	299	ASN	C-N-CA	-6.13	113.37	119.56
3	C	306	LEU	N-CA-C	5.98	118.59	111.71
3	C	245	LEU	N-CA-C	5.88	118.97	109.40
3	C	367	VAL	N-CA-C	5.68	119.00	111.17
3	C	275	GLY	N-CA-C	5.66	119.63	113.58
3	C	201	GLN	CA-C-N	5.63	125.74	119.32
3	C	201	GLN	C-N-CA	5.63	125.74	119.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	493	SER	N-CA-C	-5.44	103.20	110.55
3	C	409	ILE	N-CA-C	5.35	116.13	110.72
3	C	467	THR	N-CA-C	5.32	121.56	109.81
3	C	172	PRO	N-CA-C	5.27	118.78	110.50
2	B	42	PHE	N-CA-C	5.24	117.68	111.33
3	C	443	LYS	CA-C-N	5.23	125.17	119.78
3	C	443	LYS	C-N-CA	5.23	125.17	119.78
3	C	101	ASN	CA-C-N	5.21	124.87	119.56
3	C	101	ASN	C-N-CA	5.21	124.87	119.56
3	C	252	GLU	N-CA-C	5.13	116.88	111.28
3	C	565	PHE	N-CA-C	5.03	117.69	109.59
3	C	173	GLY	N-CA-C	5.01	122.56	112.34

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	776	0	807	4	0
2	B	784	0	775	11	0
3	C	4123	0	4080	44	0
4	C	2	0	0	0	0
5	C	5	0	4	0	0
6	A	28	0	0	1	0
6	B	23	0	0	2	0
6	C	226	0	0	3	0
All	All	5967	0	5666	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:545:VAL:O	3:C:548:GLU:HG2	1.96	0.66
3:C:297:SER:OG	3:C:349:HIS:HE1	1.81	0.64
3:C:236:ALA:O	3:C:518:ARG:HD2	1.98	0.63
2:B:83:ALA:HB1	2:B:88:ARG:HH21	1.65	0.61
3:C:275:GLY:HA3	3:C:338:ARG:HH22	1.64	0.61
3:C:298:THR:CG2	3:C:360:ASP:HB2	2.33	0.58
3:C:395:ASN:HD22	3:C:397:ASN:H	1.52	0.57
3:C:300:PRO:HG2	6:C:1118:HOH:O	2.06	0.56
3:C:300:PRO:HD3	3:C:365:GLY:HA2	1.90	0.54
1:A:6:ARG:O	1:A:10:LYS:HG2	2.08	0.53
3:C:201:GLN:HE21	3:C:203:ASP:H	1.57	0.52
2:B:16:ASN:HA	6:B:119:HOH:O	2.09	0.51
1:A:43:ILE:HD11	1:A:95:VAL:HG21	1.91	0.51
3:C:363:ALA:O	3:C:364:MET:HB2	2.11	0.51
3:C:246:HIS:CE1	3:C:278:GLY:HA3	2.46	0.50
3:C:218:ILE:HD13	3:C:243:VAL:HG13	1.93	0.49
3:C:469:GLN:HE21	3:C:470:PRO:HA	1.77	0.49
3:C:298:THR:HG22	3:C:360:ASP:HB2	1.95	0.48
3:C:203:ASP:O	3:C:207:GLU:HG3	2.13	0.47
3:C:138:ILE:HD12	3:C:362:GLN:HB2	1.96	0.47
3:C:505:LEU:HB2	3:C:507:LEU:HD13	1.95	0.47
3:C:170:CYS:O	3:C:172:PRO:HD3	2.14	0.47
3:C:460:ASP:HB3	3:C:463:ALA:HB2	1.96	0.46
2:B:4:GLY:HA2	3:C:22:ARG:O	2.14	0.46
3:C:373:ARG:O	3:C:377:VAL:HG23	2.16	0.46
1:A:83:GLU:HG2	1:A:92:LEU:HA	1.98	0.46
2:B:7:HIS:HB3	3:C:20:LYS:HB2	1.98	0.46
3:C:380:ARG:NE	3:C:555:ALA:HB2	2.30	0.46
3:C:418:ALA:O	3:C:428:LYS:HE2	2.17	0.46
3:C:218:ILE:HD11	3:C:232:ALA:CB	2.47	0.45
3:C:465:ILE:HG13	3:C:468:PRO:HD3	1.97	0.45
2:B:39:HIS:HA	2:B:60:ARG:HD2	1.98	0.45
1:A:22:ARG:HD3	6:A:122:HOH:O	2.15	0.45
3:C:507:LEU:HG	6:C:1087:HOH:O	2.17	0.45
3:C:296:SER:HB3	3:C:356:LEU:HB2	1.97	0.45
3:C:395:ASN:HD22	3:C:395:ASN:C	2.25	0.45
3:C:300:PRO:HG3	3:C:364:MET:O	2.18	0.44
2:B:90:VAL:HB	2:B:97:VAL:HG11	1.99	0.44
3:C:72:ASN:HB2	3:C:122:GLU:HB3	2.00	0.44
3:C:304:TYR:OH	3:C:339:ARG:HG3	2.17	0.44
3:C:484:ARG:O	3:C:488:ARG:HG2	2.18	0.44
3:C:75:ILE:O	3:C:81:ILE:HA	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:89:ALA:HA	2:B:100:PRO:HA	2.00	0.43
2:B:72:GLU:H	2:B:75:GLN:NE2	2.17	0.43
3:C:526:VAL:HG12	3:C:527:HIS:CD2	2.53	0.42
3:C:271:PHE:O	3:C:272:HIS:C	2.61	0.42
3:C:349:HIS:HD2	3:C:406:LYS:NZ	2.18	0.42
3:C:65:CYS:HA	3:C:111:PRO:HG2	2.00	0.42
3:C:540:THR:O	3:C:541:TYR:HB2	2.20	0.42
2:B:21:THR:HA	2:B:81:LEU:O	2.19	0.42
2:B:53:ARG:HD3	6:B:112:HOH:O	2.20	0.41
3:C:246:HIS:C	3:C:246:HIS:CD2	2.98	0.41
3:C:297:SER:OG	3:C:349:HIS:CE1	2.68	0.41
2:B:100:PRO:O	2:B:101:LEU:CB	2.68	0.41
3:C:419:HIS:CD2	6:C:1060:HOH:O	2.74	0.41
3:C:246:HIS:NE2	3:C:278:GLY:HA3	2.37	0.40
3:C:419:HIS:CD2	3:C:419:HIS:H	2.40	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
2	B	99/106 (93%)	95 (96%)	4 (4%)	0	100	100
3	C	547/567 (96%)	519 (95%)	25 (5%)	3 (0%)	24	21
All	All	744/773 (96%)	711 (96%)	30 (4%)	3 (0%)	30	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	360	ASP
3	C	481	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	364	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/85 (100%)	85 (100%)	0	100	100
2	B	78/83 (94%)	76 (97%)	2 (3%)	40	44
3	C	430/442 (97%)	417 (97%)	13 (3%)	36	38
All	All	593/610 (97%)	578 (98%)	15 (2%)	42	45

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	81	LEU
2	B	97	VAL
3	C	179	ARG
3	C	199	VAL
3	C	282	PRO
3	C	311	GLU
3	C	316	LEU
3	C	332	PHE
3	C	336	ARG
3	C	349	HIS
3	C	395	ASN
3	C	449	LYS
3	C	507	LEU
3	C	508	ARG
3	C	520	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	29	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	201	GLN
3	C	349	HIS
3	C	395	ASN
3	C	419	HIS
3	C	469	GLN
3	C	486	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	KCX	C	217	4,3	10,11,12	0.94	1 (10%)	6,12,14	1.48	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KCX	C	217	4,3	-	0/9/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	217	KCX	OQ1-CX	2.21	1.25	1.21

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	217	KCX	OQ1-CX-NZ	-3.27	119.95	124.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	HAE	C	989	4	4,4,4	0.91	0	2,4,4	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HAE	C	989	4	-	0/1/2/2	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/100 (100%)	-0.68	0 100 100	4, 13, 27, 41	0
2	B	101/106 (95%)	-0.30	1 (0%) 79 79	8, 19, 33, 51	0
3	C	551/567 (97%)	-0.63	6 (1%) 78 77	3, 11, 27, 75	0
All	All	752/773 (97%)	-0.60	7 (0%) 81 80	3, 12, 30, 75	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	313	LEU	5.6
3	C	317	MET	5.5
3	C	314	ASP	4.2
3	C	316	LEU	4.0
3	C	333	ALA	2.8
2	B	101	LEU	2.4
3	C	332	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	KCX	C	217	12/13	0.96	0.05	2,9,13,14	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	HAE	C	989	5/5	0.98	0.06	23,25,28,29	0
4	NI	C	575	1/1	1.00	0.05	22,22,22,22	0
4	NI	C	574	1/1	1.00	0.05	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.