



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:33 AM UTC

PDB ID : 5FWY / pdb_00005fwy
Title : Crystal structure of the AMPA receptor GluA2/A3 N-terminal domain heterodimer
Authors : Herguedas, B.; Garcia-Nafria, J.; Greger, I.H.
Deposited on : 2016-02-21
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	385	 3% 87% 9% . .
1	C	385	 4% 92% . 5%
2	B	390	 2% 92% 5% .
2	D	390	 2% 89% 7% . .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12549 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUTAMATE RECEPTOR 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	371	Total	C	N	O	S	0	1	0
			2956	1885	501	561	9			
1	C	367	Total	C	N	O	S	0	1	0
			2927	1870	495	553	9			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	380	ARG	-	expression tag	UNP P19491
A	381	THR	-	expression tag	UNP P19491
A	382	LYS	-	expression tag	UNP P19491
A	383	HIS	-	expression tag	UNP P19491
A	384	HIS	-	expression tag	UNP P19491
A	385	HIS	-	expression tag	UNP P19491
A	386	HIS	-	expression tag	UNP P19491
A	387	HIS	-	expression tag	UNP P19491
A	388	HIS	-	expression tag	UNP P19491
C	380	ARG	-	expression tag	UNP P19491
C	381	THR	-	expression tag	UNP P19491
C	382	LYS	-	expression tag	UNP P19491
C	383	HIS	-	expression tag	UNP P19491
C	384	HIS	-	expression tag	UNP P19491
C	385	HIS	-	expression tag	UNP P19491
C	386	HIS	-	expression tag	UNP P19491
C	387	HIS	-	expression tag	UNP P19491
C	388	HIS	-	expression tag	UNP P19491

- Molecule 2 is a protein called GLUTAMATE RECEPTOR 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	379	Total	C	N	O	S	0	0	0
			3069	1952	536	565	16			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	377	Total	C	N	O	S	0	1	0
			3059	1949	533	560	17			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	382	GLY	-	expression tag	UNP P19492
B	383	THR	-	expression tag	UNP P19492
B	384	LYS	-	expression tag	UNP P19492
B	385	HIS	-	expression tag	UNP P19492
B	386	HIS	-	expression tag	UNP P19492
B	387	HIS	-	expression tag	UNP P19492
B	388	HIS	-	expression tag	UNP P19492
B	389	HIS	-	expression tag	UNP P19492
B	390	HIS	-	expression tag	UNP P19492
D	382	GLY	-	expression tag	UNP P19492
D	383	THR	-	expression tag	UNP P19492
D	384	LYS	-	expression tag	UNP P19492
D	385	HIS	-	expression tag	UNP P19492
D	386	HIS	-	expression tag	UNP P19492
D	387	HIS	-	expression tag	UNP P19492
D	388	HIS	-	expression tag	UNP P19492
D	389	HIS	-	expression tag	UNP P19492
D	390	HIS	-	expression tag	UNP P19492

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 14 8 1 5	0	0
3	A	1	Total C N O 14 8 1 5	0	0
3	B	1	Total C N O 14 8 1 5	0	0
3	B	1	Total C N O 14 8 1 5	0	0
3	B	1	Total C N O 14 8 1 5	0	0
3	C	1	Total C N O 14 8 1 5	0	0
3	C	1	Total C N O 14 8 1 5	0	0
3	D	1	Total C N O 14 8 1 5	0	0
3	D	1	Total C N O 14 8 1 5	0	0
3	D	1	Total C N O 14 8 1 5	0	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	88	Total	O	0	0
			88	88		
6	B	102	Total	O	0	0
			102	102		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	87	Total	O	0	0
			87	87		
6	D	75	Total	O	0	0
			75	75		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: GLUTAMATE RECEPTOR 2




• Molecule 1: GLUTAMATE RECEPTOR 2




• Molecule 2: GLUTAMATE RECEPTOR 3



• Molecule 2: GLUTAMATE RECEPTOR 3





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	66.27Å 78.52Å 107.85Å 83.36° 87.40° 64.94°	Depositor
Resolution (Å)	41.76 – 2.12 41.76 – 2.12	Depositor EDS
% Data completeness (in resolution range)	90.3 (41.76-2.12) 90.3 (41.76-2.12)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.12Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.192 , 0.231 0.198 , 0.233	Depositor DCC
R_{free} test set	4984 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.310	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.077 for h,h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12549	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.08	4/3021 (0.1%)	1.05	3/4088 (0.1%)
1	C	1.12	4/2991 (0.1%)	1.06	3/4045 (0.1%)
2	B	1.08	0/3140	1.04	3/4250 (0.1%)
2	D	1.04	1/3133 (0.0%)	1.04	3/4241 (0.1%)
All	All	1.08	9/12285 (0.1%)	1.05	12/16624 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	372	ASP	CA-C	9.13	1.58	1.53
1	A	372	ASP	N-CA	7.52	1.50	1.46
1	C	351	THR	C-O	-5.66	1.16	1.24
1	A	238	GLY	C-O	5.54	1.30	1.23
1	C	295	LYS	C-O	-5.32	1.17	1.24
1	C	372	ASP	CA-C	5.30	1.60	1.52
1	A	136	GLY	N-CA	5.23	1.50	1.45
1	C	374	MET	C-O	-5.19	1.17	1.24
2	D	136	LEU	C-O	5.08	1.30	1.24

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	75	GLY	N-CA-C	7.33	120.44	110.69
1	C	372	ASP	N-CA-C	6.95	121.61	113.20
2	D	213	ARG	N-CA-C	6.82	119.58	111.33
2	D	75	GLY	N-CA-C	6.59	120.09	110.80
1	C	69	GLY	N-CA-C	6.33	119.73	110.80
1	C	92	THR	N-CA-C	6.15	118.19	108.23
2	B	213	ARG	N-CA-C	6.06	118.67	111.71
1	A	69	GLY	N-CA-C	6.05	119.96	111.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	374	MET	N-CA-C	5.84	118.18	109.25
2	D	106	ASP	CB-CA-C	-5.70	101.76	110.88
2	B	272	ASN	N-CA-C	5.44	119.91	113.28
1	A	372	ASP	CB-CA-C	-5.37	109.34	117.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2956	0	2897	17	0
1	C	2927	0	2875	6	0
2	B	3069	0	2964	12	0
2	D	3059	0	2965	13	0
3	A	28	0	26	0	0
3	B	42	0	39	0	0
3	C	28	0	26	0	0
3	D	42	0	39	0	0
4	A	5	0	0	0	0
4	B	20	0	0	0	0
4	C	5	0	0	0	0
4	D	10	0	0	0	0
5	B	6	0	8	0	0
6	A	88	0	0	0	0
6	B	102	0	0	2	0
6	C	87	0	0	0	0
6	D	75	0	0	0	0
All	All	12549	0	11839	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:190:LEU:HD13	2:D:218:MET:HE2	1.57	0.87
1:C:27:MET:HE2	1:C:283:VAL:HG13	1.72	0.71
1:C:27:MET:HE1	1:C:37:LEU:HD12	1.81	0.62
1:A:27:MET:HE2	1:A:283:VAL:HG13	1.82	0.61
2:B:232:VAL:CG1	2:B:239:ILE:HD11	2.32	0.58
1:A:269:THR:HG23	1:A:271:THR:O	2.04	0.58
2:D:13:MET:HE2	2:D:76:PHE:HB3	1.86	0.56
2:D:125:LEU:HG	2:D:218:MET:HE1	1.87	0.56
2:D:106:ASP:OD1	2:D:349:ARG:NH2	2.40	0.55
1:C:27:MET:HE1	1:C:37:LEU:CD1	2.36	0.55
1:A:17:ASP:HB3	1:A:265:PRO:HB2	1.88	0.54
2:B:134:VAL:HG21	2:B:181:MET:HG3	1.88	0.54
1:A:241:ILE:HD12	1:A:242:VAL:HG23	1.93	0.51
2:B:256:ILE:HD12	6:B:2029:HOH:O	2.11	0.50
1:A:137:LEU:HD12	2:B:143:PHE:CD1	2.46	0.50
1:A:70:PHE:CE2	1:A:93:PRO:HG2	2.47	0.50
2:B:349:ARG:HD2	6:B:2098:HOH:O	2.11	0.50
1:A:221:PHE:CD1	1:A:238:GLY:HA3	2.47	0.49
1:A:91:ILE:HD13	1:A:286:MET:HE3	1.95	0.49
2:D:121:ILE:HG23	2:D:218:MET:CE	2.43	0.48
1:C:27:MET:CE	1:C:283:VAL:HG13	2.42	0.48
2:B:103:THR:OG1	2:B:104:ASP:N	2.46	0.48
1:A:68:PHE:CZ	1:A:279:THR:HG23	2.49	0.48
2:B:13:MET:HE2	2:B:76:PHE:HB3	1.96	0.47
2:D:121:ILE:HG23	2:D:218:MET:HE3	1.96	0.47
2:D:76:PHE:CE2	2:D:99:PRO:HG2	2.50	0.47
1:A:241:ILE:HD11	1:A:274:TYR:HB2	1.97	0.46
1:A:179:GLU:OE2	1:A:212:TYR:OH	2.28	0.45
1:C:27:MET:HE2	1:C:283:VAL:CG1	2.44	0.45
2:D:99:PRO:HB3	2:D:281:LEU:HB3	1.99	0.45
2:B:232:VAL:HG11	2:B:239:ILE:HD11	2.00	0.44
1:A:250:SER:O	1:A:254:GLU:HG3	2.18	0.44
1:A:372:ASP:O	1:A:373:LYS:HD3	2.19	0.43
1:A:184:ARG:HD2	1:A:212:TYR:CE2	2.54	0.43
2:D:111:ILE:HD13	2:D:330:LEU:HD13	2.00	0.43
1:A:44:LEU:N	1:A:44:LEU:HD23	2.34	0.42
2:D:132:LYS:HG3	2:D:159:GLN:HB2	2.00	0.42
2:B:161:THR:HG21	2:B:181:MET:HE1	2.00	0.42
2:D:230:GLU:HA	2:D:233[B]:MET:HE2	2.02	0.42
2:D:140:GLU:OE2	2:D:197:ARG:NH1	2.52	0.42
1:C:174:LEU:C	1:C:174:LEU:HD23	2.44	0.42
2:D:121:ILE:HD11	2:D:220:ALA:HB1	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:LEU:HD12	2:B:143:PHE:CG	2.55	0.41
1:A:155:THR:HG23	2:B:154:VAL:HG22	2.02	0.41
2:B:324:ILE:HD13	2:B:324:ILE:HG21	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	368/385 (96%)	358 (97%)	10 (3%)	0	100	100
1	C	362/385 (94%)	353 (98%)	9 (2%)	0	100	100
2	B	375/390 (96%)	367 (98%)	8 (2%)	0	100	100
2	D	374/390 (96%)	369 (99%)	5 (1%)	0	100	100
All	All	1479/1550 (95%)	1447 (98%)	32 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/334 (96%)	315 (98%)	5 (2%)	55	63
1	C	317/334 (95%)	314 (99%)	3 (1%)	70	78

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	328/341 (96%)	324 (99%)	4 (1%)	63	71
2	D	328/341 (96%)	318 (97%)	10 (3%)	36	40
All	All	1293/1350 (96%)	1271 (98%)	22 (2%)	53	61

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	VAL
1	A	173	SER
1	A	205	ILE
1	A	300	ILE
1	A	321	VAL
2	B	124	LEU
2	B	318	VAL
2	B	324	ILE
2	B	335	VAL
1	C	135	ARG
1	C	216	ILE
1	C	370	GLU
2	D	24	ARG
2	D	39	LYS
2	D	111	ILE
2	D	129	LYS
2	D	230	GLU
2	D	324	ILE
2	D	361	LYS
2	D	362	VAL
2	D	365	SER
2	D	378	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	GLN
1	A	77	ASN
1	A	163	ASN
1	A	284	GLN
2	B	49	HIS
2	B	159	GLN
2	B	204	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	253	GLN
2	B	254	GLN
2	B	336	GLN
1	C	4	ASN
1	C	353	ASN
2	D	42	HIS
2	D	65	GLN
2	D	147	GLN
2	D	234	HIS
2	D	243	GLN
2	D	253	GLN
2	D	336	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	D	1385	-	4,4,4	0.47	0	6,6,6	0.13	0
5	GOL	B	1388	-	5,5,5	0.39	0	5,5,5	0.25	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	1379	1	14,14,15	0.55	0	17,19,21	1.60	4 (23%)
4	SO4	C	1380	-	4,4,4	0.48	0	6,6,6	0.17	0
3	NAG	D	1382	2	14,14,15	0.61	0	17,19,21	0.93	0
3	NAG	B	1385	2	14,14,15	0.52	0	17,19,21	2.39	5 (29%)
3	NAG	B	1384	2	14,14,15	0.62	0	17,19,21	1.18	1 (5%)
4	SO4	A	1380	-	4,4,4	0.51	0	6,6,6	0.36	0
3	NAG	C	1379	1	14,14,15	0.47	0	17,19,21	1.65	3 (17%)
4	SO4	B	1390	-	4,4,4	0.47	0	6,6,6	0.22	0
4	SO4	D	1384	-	4,4,4	0.43	0	6,6,6	0.13	0
3	NAG	A	1378	1	14,14,15	0.56	0	17,19,21	1.42	2 (11%)
4	SO4	B	1387	-	4,4,4	0.45	0	6,6,6	0.31	0
3	NAG	C	1378	1	14,14,15	0.72	1 (7%)	17,19,21	1.12	2 (11%)
3	NAG	D	1381	2	14,14,15	0.56	0	17,19,21	1.38	3 (17%)
4	SO4	B	1389	-	4,4,4	0.55	0	6,6,6	0.30	0
3	NAG	D	1383	2	14,14,15	0.51	0	17,19,21	1.16	1 (5%)
3	NAG	B	1383	2	14,14,15	0.52	0	17,19,21	1.03	1 (5%)
4	SO4	B	1386	-	4,4,4	0.37	0	6,6,6	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	1384	2	-	0/6/23/26	0/1/1/1
5	GOL	B	1388	-	-	0/4/4/4	-
3	NAG	A	1379	1	-	0/6/23/26	0/1/1/1
3	NAG	C	1379	1	-	0/6/23/26	0/1/1/1
3	NAG	A	1378	1	-	0/6/23/26	0/1/1/1
3	NAG	D	1382	2	-	0/6/23/26	0/1/1/1
3	NAG	B	1385	2	-	3/6/23/26	0/1/1/1
3	NAG	C	1378	1	-	2/6/23/26	0/1/1/1
3	NAG	D	1381	2	-	0/6/23/26	0/1/1/1
3	NAG	B	1383	2	-	0/6/23/26	0/1/1/1
3	NAG	D	1383	2	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1378	NAG	C1-C2	2.16	1.55	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1385	NAG	C2-N2-C7	5.32	130.03	122.90
3	B	1385	NAG	O5-C1-C2	-4.27	104.69	111.29
3	B	1385	NAG	C8-C7-N2	4.17	123.03	116.12
3	C	1379	NAG	C1-C2-N2	-4.05	104.06	110.43
3	C	1379	NAG	C1-O5-C5	3.89	117.40	112.19
3	A	1379	NAG	C1-C2-N2	-3.65	104.69	110.43
3	B	1385	NAG	C1-C2-N2	3.30	115.64	110.43
3	A	1378	NAG	C1-O5-C5	3.03	116.25	112.19
3	B	1384	NAG	C1-O5-C5	3.02	116.23	112.19
3	D	1381	NAG	O5-C5-C4	-2.84	103.93	110.83
3	A	1379	NAG	O5-C5-C4	-2.77	104.10	110.83
3	B	1385	NAG	C1-O5-C5	2.68	115.78	112.19
3	C	1378	NAG	C1-O5-C5	2.54	115.59	112.19
3	D	1381	NAG	C2-N2-C7	2.52	126.27	122.90
3	D	1383	NAG	C4-C3-C2	2.46	114.62	111.02
3	C	1379	NAG	C4-C3-C2	2.39	114.51	111.02
3	C	1378	NAG	C2-N2-C7	2.35	126.05	122.90
3	A	1379	NAG	O5-C5-C6	2.16	111.86	107.66
3	A	1378	NAG	C1-C2-N2	2.12	113.78	110.43
3	D	1381	NAG	C1-C2-N2	-2.12	107.09	110.43
3	B	1383	NAG	C2-N2-C7	2.07	125.68	122.90
3	A	1379	NAG	C3-C4-C5	-2.03	106.55	110.23

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	1383	NAG	O5-C5-C6-O6
3	C	1378	NAG	C4-C5-C6-O6
3	C	1378	NAG	O5-C5-C6-O6
3	D	1383	NAG	C4-C5-C6-O6
3	B	1385	NAG	C8-C7-N2-C2
3	B	1385	NAG	O7-C7-N2-C2
3	B	1385	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	371/385 (96%)	0.10	10 (2%) 56 59	24, 36, 65, 89	1 (0%)
1	C	367/385 (95%)	0.10	14 (3%) 44 47	22, 35, 63, 82	1 (0%)
2	B	379/390 (97%)	0.06	8 (2%) 63 66	24, 35, 57, 78	0
2	D	377/390 (96%)	0.15	9 (2%) 59 62	26, 39, 60, 74	1 (0%)
All	All	1494/1550 (96%)	0.10	41 (2%) 56 59	22, 36, 62, 89	3 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	273	ALA	4.8
2	D	233[A]	MET	3.7
2	D	270	ALA	3.7
1	C	180	LEU	3.6
2	B	234	HIS	3.5
1	C	371	VAL	3.3
2	D	374	TYR	3.3
2	B	233	MET	3.1
2	D	234	HIS	3.0
1	C	317	TRP	3.0
1	C	302	ARG	2.9
1	C	162	ILE	2.9
1	A	371	VAL	2.7
1	C	315	VAL	2.7
1	A	317	TRP	2.7
1	C	98	ASP	2.7
2	B	51	ASP	2.7
2	D	257	GLN	2.6
1	C	225	ASP	2.6
1	C	4	ASN	2.5
1	A	302	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	98	ASP	2.4
1	A	162	ILE	2.4
1	C	313	PRO	2.4
2	D	260	VAL	2.4
1	A	135	ARG	2.3
1	A	306	ALA	2.2
1	C	307	GLY	2.2
1	C	374	MET	2.2
2	B	374	TYR	2.2
1	C	166	LYS	2.2
1	A	374	MET	2.2
1	C	181	LYS	2.1
2	B	251	MET	2.1
1	A	180	LEU	2.1
2	B	36	THR	2.1
2	B	254	GLN	2.0
1	A	372	ASP	2.0
2	D	261	ARG	2.0
2	D	312	CYS	2.0
2	B	382	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	D	1383	14/15	0.73	0.13	70,76,79,86	0
4	SO4	B	1387	5/5	0.73	0.11	82,82,88,88	0
4	SO4	B	1390	5/5	0.74	0.16	89,92,95,96	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	C	1379	14/15	0.75	0.14	44,54,60,61	0
3	NAG	A	1379	14/15	0.76	0.16	45,56,58,61	0
4	SO4	B	1389	5/5	0.77	0.12	62,70,78,78	0
3	NAG	D	1381	14/15	0.77	0.15	48,57,59,59	0
3	NAG	B	1383	14/15	0.78	0.15	47,58,62,64	0
3	NAG	B	1385	14/15	0.78	0.14	69,76,78,80	0
3	NAG	B	1384	14/15	0.83	0.12	53,61,64,68	0
4	SO4	D	1385	5/5	0.83	0.13	85,87,91,93	0
3	NAG	D	1382	14/15	0.84	0.10	48,53,55,55	0
3	NAG	C	1378	14/15	0.87	0.09	40,47,52,56	0
3	NAG	A	1378	14/15	0.88	0.09	40,46,52,54	0
4	SO4	A	1380	5/5	0.88	0.10	62,64,70,71	0
4	SO4	D	1384	5/5	0.91	0.09	70,75,76,76	0
4	SO4	B	1386	5/5	0.92	0.08	68,71,74,74	0
4	SO4	C	1380	5/5	0.93	0.11	56,56,63,64	0
5	GOL	B	1388	6/6	0.97	0.06	36,38,39,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.