



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:18 AM UTC

PDB ID : 9FW2 / pdb_00009fw2
Title : SARS CoV-2 nsp10 in complex with the ExoN domain from nsp14
Authors : Fisher, S.Z.
Deposited on : 2024-06-28
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	131	5% 79% 19% .
2	B	290	2% 84% 13% ..

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3819 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-structural protein 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	131	Total	C	N	O	S	0	3	0
			968	604	161	187	16			

- Molecule 2 is a protein called Guanine-N7 methyltransferase nsp14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	286	Total	C	N	O	S	0	6	0
			2285	1463	392	408	22			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P0DTD1

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		
3	B	2	Total	Zn	0	0
			2	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Na	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	169	Total	O	0	0
			169	169		
6	B	380	Total	O	0	0
			380	380		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Non-structural protein 11



- Molecule 2: Guanine-N7 methyltransferase nsp14



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	92.03Å 104.78Å 55.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.98 – 1.77 37.98 – 1.77	Depositor EDS
% Data completeness (in resolution range)	100.0 (37.98-1.77) 100.0 (37.98-1.77)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 1.77Å)	Xtriage
Refinement program	PHENIX (1.18.2_3874:0000)	Depositor
R, R_{free}	0.167 , 0.183 0.169 , 0.186	Depositor DCC
R_{free} test set	2669 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	18.9	Xtriage
Anisotropy	0.513	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3819	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.47	23/998 (2.3%)	1.06	10/1359 (0.7%)
2	B	1.14	26/2365 (1.1%)	0.85	4/3217 (0.1%)
All	All	1.25	49/3363 (1.5%)	0.92	14/4576 (0.3%)

All (49) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	5[A]	THR	CA-C	7.68	1.62	1.52
2	B	5[B]	THR	CA-C	7.68	1.62	1.52
2	B	16[A]	THR	CA-C	7.57	1.61	1.52
2	B	16[B]	THR	CA-C	7.57	1.61	1.52
1	A	38	ILE	C-O	-6.97	1.16	1.24
1	A	16	PHE	C-O	-6.57	1.16	1.24
1	A	21[A]	VAL	CA-C	6.46	1.62	1.52
1	A	21[B]	VAL	CA-C	6.46	1.62	1.52
2	B	34	LYS	C-O	-6.41	1.16	1.23
1	A	85[A]	ASN	CA-C	6.36	1.61	1.53
1	A	85[B]	ASN	CA-C	6.36	1.61	1.53
2	B	5[A]	THR	C-O	6.30	1.32	1.23
2	B	5[B]	THR	C-O	6.30	1.32	1.23
2	B	245	GLN	C-O	-6.17	1.16	1.24
1	A	85[A]	ASN	C-O	6.11	1.31	1.23
1	A	85[B]	ASN	C-O	6.11	1.31	1.23
2	B	42	ILE	C-O	-6.09	1.17	1.23
2	B	141	PRO	C-O	-5.86	1.18	1.24
1	A	12	THR	C-O	-5.81	1.17	1.24
2	B	231	ILE	C-O	-5.77	1.17	1.24
2	B	214	ALA	C-O	-5.69	1.17	1.24
1	A	37	PRO	C-O	-5.67	1.16	1.23
1	A	89	PHE	C-O	-5.64	1.17	1.24
1	A	93	LYS	C-O	-5.60	1.17	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	18	LEU	C-O	-5.54	1.16	1.23
2	B	7	LEU	C-O	-5.51	1.17	1.23
1	A	26	ALA	C-O	-5.50	1.17	1.24
1	A	14	LEU	C-O	-5.48	1.17	1.24
1	A	17	CYS	C-O	-5.44	1.17	1.24
1	A	31	LEU	C-O	-5.43	1.17	1.24
2	B	17	GLY	C-O	-5.42	1.17	1.23
1	A	46	CYS	C-O	-5.36	1.17	1.24
1	A	27	TYR	C-O	-5.32	1.17	1.24
2	B	12	SER	C-O	-5.32	1.17	1.23
2	B	15	ILE	C-O	-5.28	1.18	1.23
1	A	15	SER	C-O	-5.26	1.18	1.24
1	A	114	ASN	C-O	-5.25	1.16	1.23
2	B	117	LEU	C-O	-5.25	1.17	1.24
2	B	77	GLU	C-O	-5.24	1.17	1.24
1	A	30	TYR	C-O	-5.23	1.18	1.24
2	B	35	THR	C-O	-5.20	1.17	1.23
1	A	124	LYS	C-O	-5.17	1.17	1.23
2	B	72	MET	C-O	-5.16	1.18	1.24
1	A	116	VAL	C-O	-5.11	1.18	1.24
2	B	20	PRO	C-O	-5.10	1.17	1.24
2	B	69	TYR	C-O	-5.10	1.18	1.24
2	B	243	ASP	C-O	-5.08	1.17	1.23
2	B	175	LYS	C-O	-5.07	1.18	1.24
2	B	216	CYS	C-O	-5.07	1.17	1.23

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	21[A]	VAL	CA-C-O	10.79	132.12	120.47
1	A	21[B]	VAL	CA-C-O	10.79	132.12	120.47
1	A	85[A]	ASN	N-CA-C	7.57	120.99	110.01
1	A	85[B]	ASN	N-CA-C	7.57	120.99	110.01
1	A	85[A]	ASN	CA-C-O	7.09	128.70	120.69
1	A	85[B]	ASN	CA-C-O	7.09	128.70	120.69
2	B	174	LEU	N-CA-C	6.24	121.05	113.38
2	B	15	ILE	O-C-N	5.62	128.81	122.68
1	A	85[A]	ASN	O-C-N	-5.33	115.46	121.43
1	A	85[B]	ASN	O-C-N	-5.33	115.46	121.43
1	A	21[A]	VAL	N-CA-C	5.18	117.56	111.09
1	A	21[B]	VAL	N-CA-C	5.18	117.56	111.09
2	B	263	VAL	N-CA-C	5.15	117.84	112.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	234	ASP	N-CA-C	5.12	120.52	113.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	968	0	934	9	0
2	B	2285	0	2257	23	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	6	0	8	1	0
4	B	6	0	8	0	0
5	B	1	0	0	0	0
6	A	169	0	0	2	0
6	B	380	0	0	9	0
All	All	3819	0	3207	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:16[B]:THR:HG21	6:B:423:HOH:O	1.61	0.99
2:B:16[B]:THR:HG22	2:B:53:ARG:HH22	1.36	0.89
2:B:97[B]:THR:HG22	6:B:465:HOH:O	1.73	0.88
1:A:63:MET:HG3	6:A:466:HOH:O	1.86	0.74
6:A:311:HOH:O	2:B:5[A]:THR:HG21	1.93	0.68
2:B:97[B]:THR:HG21	2:B:139:LYS:HG2	1.74	0.67
2:B:26:HIS:HE1	6:B:525:HOH:O	1.77	0.66
2:B:135:ARG:HD2	6:B:431:HOH:O	1.96	0.65
2:B:26:HIS:CE1	6:B:525:HOH:O	2.50	0.64
2:B:165:LYS:HE2	6:B:496:HOH:O	1.98	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:31[B]:THR:HG23	6:B:515:HOH:O	1.98	0.62
2:B:255:SER:O	2:B:259:LEU:HD13	2.01	0.61
2:B:16[B]:THR:CG2	6:B:423:HOH:O	2.33	0.59
1:A:21[A]:VAL:HG13	2:B:200:LYS:HB3	1.85	0.57
1:A:92:LEU:HD22	1:A:97[A]:VAL:HG21	1.86	0.57
1:A:48:HIS:O	1:A:63:MET:HE2	2.06	0.55
2:B:16[B]:THR:CG2	2:B:53:ARG:HH22	2.15	0.52
2:B:97[B]:THR:HG21	2:B:139:LYS:CG	2.40	0.51
1:A:21[B]:VAL:HG23	2:B:200:LYS:HB3	1.92	0.51
2:B:95:HIS:O	2:B:97[B]:THR:HG23	2.14	0.48
2:B:96:ALA:C	2:B:97[B]:THR:HG23	2.40	0.47
2:B:257:HIS:CE1	2:B:264:HIS:HB2	2.50	0.46
1:A:85[A]:ASN:ND2	1:A:85[A]:ASN:H	2.14	0.46
2:B:32:LYS:NZ	6:B:408:HOH:O	2.45	0.45
2:B:97[B]:THR:CG2	2:B:139:LYS:HG2	2.43	0.45
1:A:92:LEU:HD22	1:A:97[A]:VAL:CG2	2.48	0.44
1:A:131:ASP:O	4:A:203:GOL:O3	2.28	0.42
2:B:141:PRO:HA	2:B:142:PRO:HD3	1.78	0.42
1:A:41:CYS:SG	1:A:107:PRO:HG2	2.60	0.41
2:B:135:ARG:HE	2:B:135:ARG:HB3	1.74	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	132/131 (101%)	128 (97%)	4 (3%)	0	100	100
2	B	290/290 (100%)	282 (97%)	7 (2%)	1 (0%)	36	22
All	All	422/421 (100%)	410 (97%)	11 (3%)	1 (0%)	43	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	98	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/105 (103%)	107 (99%)	1 (1%)	70	59
2	B	257/254 (101%)	255 (99%)	2 (1%)	73	63
All	All	365/359 (102%)	362 (99%)	3 (1%)	73	63

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	LYS
2	B	36	GLU
2	B	259	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	245	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	203	-	5,5,5	0.38	0	5,5,5	1.00	0
4	GOL	B	303	-	5,5,5	0.49	0	5,5,5	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	203	-	-	1/4/4/4	-
4	GOL	B	303	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	203	GOL	O1-C1-C2-O2
4	B	303	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	203	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	131/131 (100%)	-0.13	6 (4%) 37 43	10, 20, 32, 57	3 (2%)
2	B	286/290 (98%)	-0.18	5 (1%) 69 77	12, 19, 34, 59	6 (2%)
All	All	417/421 (99%)	-0.17	11 (2%) 57 65	10, 19, 35, 59	9 (2%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	15	ILE	5.8
2	B	16[A]	THR	3.7
2	B	14	VAL	3.5
2	B	288	LYS	3.4
1	A	86	PRO	3.3
1	A	85[A]	ASN	3.1
1	A	83	HIS	3.0
2	B	3	ASN	2.4
1	A	84	PRO	2.3
1	A	129	SER	2.3
1	A	87	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	203	6/6	0.70	0.21	35,43,50,51	0
4	GOL	B	303	6/6	0.83	0.14	28,39,49,57	0
5	NA	B	304	1/1	0.91	0.16	36,36,36,36	0
3	ZN	A	201	1/1	0.99	0.03	23,23,23,23	0
3	ZN	A	202	1/1	1.00	0.02	17,17,17,17	0
3	ZN	B	301	1/1	1.00	0.02	19,19,19,19	0
3	ZN	B	302	1/1	1.00	0.02	17,17,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN A 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.