



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:57 PM UTC

PDB ID : 5FXV / pdb_00005fxv
Title : Structure of the human RBBP4:MTA1(464-546) complex
Authors : Millard, C.J.; Varma, N.; Fairall, L.; Schwabe, J.W.R.
Deposited on : 2016-03-03
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	425	87% 10%
1	C	425	87% 11%
1	E	425	87% 11%
1	G	425	83% 6% 10%
2	B	85	74% 7% 19%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	85	4%73%8%19%
2	F	85	75%5%20%
2	H	85	2%76%•20%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 14366 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HISTONE-BINDING PROTEIN RBBP4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	381	Total	C	N	O	S	0	0	0
			3026	1912	514	590	10			
1	C	380	Total	C	N	O	S	0	0	0
			3019	1907	513	589	10			
1	E	380	Total	C	N	O	S	0	0	0
			3019	1907	513	589	10			
1	G	381	Total	C	N	O	S	0	0	0
			3026	1912	514	590	10			

- Molecule 2 is a protein called METASTASIS-ASSOCIATED PROTEIN MTA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	69	Total	C	N	O	S	0	0	0
			570	364	116	88	2			
2	D	69	Total	C	N	O	S	0	0	0
			574	366	116	90	2			
2	F	68	Total	C	N	O	S	0	0	0
			563	360	112	89	2			
2	H	68	Total	C	N	O	S	0	0	0
			569	363	115	89	2			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	462	GLY	-	expression tag	UNP Q13330
B	463	ALA	-	expression tag	UNP Q13330
D	462	GLY	-	expression tag	UNP Q13330
D	463	ALA	-	expression tag	UNP Q13330
F	462	GLY	-	expression tag	UNP Q13330
F	463	ALA	-	expression tag	UNP Q13330
H	462	GLY	-	expression tag	UNP Q13330
H	463	ALA	-	expression tag	UNP Q13330

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: HISTONE-BINDING PROTEIN RBBP4

Chain A: 




• Molecule 1: HISTONE-BINDING PROTEIN RBBP4

Chain C: 




• Molecule 1: HISTONE-BINDING PROTEIN RBBP4


Chain E: 





• Molecule 1: HISTONE-BINDING PROTEIN RBBP4

Chain G: 








• Molecule 2: METASTASIS-ASSOCIATED PROTEIN MTA1





• Molecule 2: METASTASIS-ASSOCIATED PROTEIN MTA1



• Molecule 2: METASTASIS-ASSOCIATED PROTEIN MTA1



• Molecule 2: METASTASIS-ASSOCIATED PROTEIN MTA1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.29Å 150.07Å 95.59Å 90.00° 94.54° 90.00°	Depositor
Resolution (Å)	95.29 – 3.20 95.29 – 3.20	Depositor EDS
% Data completeness (in resolution range)	97.5 (95.29-3.20) 97.5 (95.29-3.20)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.249 , 0.291 0.248 , 0.289	Depositor DCC
R_{free} test set	1753 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	80.8	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	14366	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/3107	0.67	0/4233
1	C	0.51	0/3099	0.67	0/4221
1	E	0.53	0/3099	0.69	0/4221
1	G	0.53	0/3107	0.70	0/4233
2	B	0.58	0/584	0.78	0/791
2	D	0.59	0/588	0.82	0/796
2	F	0.57	0/577	0.78	0/782
2	H	0.56	0/583	0.80	0/789
All	All	0.53	0/14744	0.70	0/20066

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	139	ILE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3026	0	2876	5	0
1	C	3019	0	2869	3	0
1	E	3019	0	2869	2	0
1	G	3026	0	2876	11	0
2	B	570	0	597	0	0
2	D	574	0	601	0	0
2	F	563	0	588	0	0
2	H	569	0	596	0	0
All	All	14366	0	13872	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:139:ILE:HG12	1:G:152:PHE:C	2.32	0.55
1:A:133:MET:HE3	1:A:139:ILE:HG21	1.89	0.54
1:A:68:LEU:HD21	1:A:399:MET:HE2	1.90	0.52
1:G:139:ILE:HG13	1:G:140:ILE:N	2.26	0.50
1:G:292:GLY:HA2	1:G:298:VAL:HG12	1.93	0.50
1:G:201:THR:CG2	1:G:224:THR:HG22	2.43	0.49
1:A:133:MET:HE2	1:A:185:TRP:CZ3	2.49	0.48
1:G:292:GLY:CA	1:G:298:VAL:HG12	2.44	0.48
1:E:298:VAL:HG23	1:E:320:ILE:HD13	1.97	0.47
1:C:189:LEU:HD21	1:C:240:GLU:HB2	1.99	0.45
1:G:139:ILE:HD11	1:G:152:PHE:HB2	1.99	0.44
1:G:201:THR:HG22	1:G:224:THR:HG22	1.99	0.44
1:G:149:VAL:HG11	1:G:193:LEU:HD21	1.99	0.43
1:G:63:ILE:HD11	1:G:83:SER:HB2	2.00	0.43
1:A:64:HIS:CE1	1:A:86:LEU:HD12	2.53	0.43
1:E:133:MET:HE2	1:E:185:TRP:CZ3	2.54	0.43
1:A:141:ALA:HB2	1:A:185:TRP:CZ2	2.54	0.43
1:C:46:THR:HG21	1:C:130:ALA:HB3	2.02	0.42
1:G:139:ILE:HG13	1:G:140:ILE:H	1.84	0.42
1:G:139:ILE:HD13	1:G:154:TYR:N	2.35	0.42
1:C:133:MET:HE2	1:C:185:TRP:CZ3	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	371/425 (87%)	362 (98%)	8 (2%)	1 (0%)	36	68
1	C	370/425 (87%)	353 (95%)	16 (4%)	1 (0%)	36	68
1	E	370/425 (87%)	352 (95%)	16 (4%)	2 (0%)	24	59
1	G	371/425 (87%)	349 (94%)	20 (5%)	2 (0%)	24	59
2	B	65/85 (76%)	59 (91%)	5 (8%)	1 (2%)	8	37
2	D	65/85 (76%)	58 (89%)	5 (8%)	2 (3%)	3	22
2	F	64/85 (75%)	59 (92%)	4 (6%)	1 (2%)	7	36
2	H	64/85 (75%)	59 (92%)	4 (6%)	1 (2%)	7	36
All	All	1740/2040 (85%)	1651 (95%)	78 (4%)	11 (1%)	21	56

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	315	SER
1	E	315	SER
1	C	315	SER
1	G	315	SER
1	E	307	LYS
2	B	532	LYS
2	D	532	LYS
2	F	532	LYS
2	H	532	LYS
2	D	545	PRO
1	G	139	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	340/375 (91%)	337 (99%)	3 (1%)	70	81
1	C	339/375 (90%)	335 (99%)	4 (1%)	63	79
1	E	339/375 (90%)	333 (98%)	6 (2%)	51	74
1	G	340/375 (91%)	325 (96%)	15 (4%)	25	59
2	B	58/71 (82%)	53 (91%)	5 (9%)	10	36
2	D	59/71 (83%)	54 (92%)	5 (8%)	10	37
2	F	58/71 (82%)	55 (95%)	3 (5%)	21	54
2	H	59/71 (83%)	57 (97%)	2 (3%)	32	64
All	All	1592/1784 (89%)	1549 (97%)	43 (3%)	39	68

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	LEU
1	A	68	LEU
1	A	126	GLU
2	B	477	LYS
2	B	479	THR
2	B	481	ILE
2	B	508	ILE
2	B	531	ARG
1	C	40	LEU
1	C	63	ILE
1	C	189	LEU
1	C	233	VAL
2	D	479	THR
2	D	481	ILE
2	D	508	ILE
2	D	531	ARG
2	D	546	ARG
1	E	75	GLU
1	E	195	SER
1	E	207	ILE
1	E	233	VAL
1	E	307	LYS
1	E	308	LEU
2	F	481	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	F	508	ILE
2	F	531	ARG
1	G	54	THR
1	G	63	ILE
1	G	115	ILE
1	G	117	ILE
1	G	167	CYS
1	G	183	LEU
1	G	194	LEU
1	G	207	ILE
1	G	210	VAL
1	G	215	LYS
1	G	233	VAL
1	G	238	LEU
1	G	240	GLU
1	G	397	ASN
1	G	410	ASN
2	H	481	ILE
2	H	531	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	88	ASN
1	A	267	HIS
1	A	277	ASN
1	C	64	HIS
1	C	192	HIS
1	C	267	HIS
1	E	128	ASN
1	E	188	ASN
1	E	267	HIS
1	E	328	HIS
2	F	474	HIS
2	F	544	HIS
1	G	78	HIS
1	G	128	ASN
1	G	136	ASN
1	G	277	ASN
1	G	397	ASN
2	H	544	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	381/425 (89%)	0.20	2 (0%) 87 76	54, 79, 108, 137	0
1	C	380/425 (89%)	0.22	6 (1%) 70 51	57, 84, 118, 135	0
1	E	380/425 (89%)	0.31	10 (2%) 57 37	60, 90, 119, 149	0
1	G	381/425 (89%)	0.47	9 (2%) 59 40	65, 109, 141, 172	0
2	B	69/85 (81%)	0.52	1 (1%) 73 54	63, 94, 128, 134	0
2	D	69/85 (81%)	0.50	3 (4%) 40 24	71, 92, 136, 147	0
2	F	68/85 (80%)	0.43	0 100 100	66, 101, 131, 146	0
2	H	68/85 (80%)	0.45	2 (2%) 53 35	66, 85, 133, 148	0
All	All	1796/2040 (88%)	0.33	33 (1%) 67 48	54, 89, 129, 172	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	175	GLY	3.5
1	C	180	GLY	3.4
1	G	107	GLY	3.2
1	G	167	CYS	3.1
1	C	210	VAL	2.9
1	E	37	THR	2.8
1	E	257	THR	2.6
1	G	103	GLY	2.6
1	G	139	ILE	2.6
1	E	308	LEU	2.5
1	G	140	ILE	2.5
1	E	113	GLY	2.5
2	D	544	HIS	2.4
1	A	111	VAL	2.4
1	C	12	VAL	2.4
1	G	150	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	399	MET	2.3
2	H	531	ARG	2.3
1	E	169	PRO	2.2
1	E	53	VAL	2.2
1	G	193	LEU	2.2
1	C	399	MET	2.2
2	H	516	LEU	2.2
2	D	545	PRO	2.2
1	E	11	ALA	2.2
1	C	175	GLY	2.1
1	E	221	THR	2.1
1	E	391	CYS	2.1
1	A	399	MET	2.1
1	C	203	CYS	2.1
1	E	216	VAL	2.0
2	D	529	ALA	2.0
2	B	516	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.