



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:13 AM UTC

PDB ID : 6FXP / pdb_00006fxp
Title : Crystal structure of S. aureus glucosaminidase B
Authors : Pintar, S.; Turk, D.
Deposited on : 2018-03-09
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	13299 (2.04-2.00)
Clashscore	190562	1022 (2.02-2.02)
Ramachandran outliers	187476	1014 (2.02-2.02)
Sidechain outliers	187428	1014 (2.02-2.02)
RSRZ outliers	180081	13314 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	254	11% 77% 19% ..
1	B	254	6% 77% 19% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PG4	B	601	-	-	X	-
2	PG4	B	603	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6774 atoms, of which 2073 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	250	Total	C	H	N	O	S	521	0	0
			2508	1284	488	345	386	5			
1	B	251	Total	C	H	N	O	S	525	0	0
			2515	1287	489	346	388	5			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	SER	-	expression tag	UNP A0A0H3JSV1
A	-2	ASN	-	expression tag	UNP A0A0H3JSV1
A	-1	ALA	-	expression tag	UNP A0A0H3JSV1
A	0	ALA	-	expression tag	UNP A0A0H3JSV1
A	1	ALA	-	expression tag	UNP A0A0H3JSV1
B	248	SER	-	expression tag	UNP A0A0H3JSV1
B	249	ASN	-	expression tag	UNP A0A0H3JSV1
B	250	ALA	-	expression tag	UNP A0A0H3JSV1
B	251	ALA	-	expression tag	UNP A0A0H3JSV1
B	252	ALA	-	expression tag	UNP A0A0H3JSV1

- Molecule 2 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	8	5		
2	A	1	Total	C	O	0	0
			13	8	5		
2	A	1	Total	C	O	0	0
			13	8	5		
2	A	1	Total	C	O	0	0
			13	8	5		
2	B	1	Total	C	O	0	0
			13	8	5		
2	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Na 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	252	Total 756	H 504	O 252	504	0
5	B	296	Total 888	H 592	O 296	592	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uncharacterized protein



• Molecule 1: Uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	151.77Å 151.77Å 122.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.95 – 2.03 19.95 – 2.03	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.95-2.03) 99.6 (19.95-2.03)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 2.03Å)	Xtriage
Refinement program	MAIN	Depositor
R, R_{free}	0.212 , 0.220 0.208 , 0.213	Depositor DCC
R_{free} test set	2101 reflections (3.90%)	wwPDB-VP
Wilson B-factor (Å ²)	32.7	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 87.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6774	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, PG4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.15	4/2065 (0.2%)	1.28	13/2778 (0.5%)
1	B	1.21	5/2071 (0.2%)	1.31	8/2785 (0.3%)
All	All	1.18	9/4136 (0.2%)	1.29	21/5563 (0.4%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	480	MET	SD-CE	-9.94	1.54	1.79
1	B	431	ILE	CA-C	7.26	1.61	1.52
1	A	150	ASP	CA-C	6.54	1.61	1.52
1	A	196	MET	CA-C	5.72	1.60	1.52
1	B	253	PHE	C-O	-5.67	1.17	1.23
1	A	146	VAL	CA-C	-5.60	1.46	1.52
1	B	361	ASN	CA-C	5.27	1.59	1.52
1	B	450	ASN	CA-C	5.26	1.59	1.52
1	A	199	ASN	CA-C	-5.22	1.48	1.53

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	178	ASP	N-CA-C	-6.94	103.73	111.71
1	A	197	ARG	N-CA-C	6.92	118.90	111.36
1	A	239	PHE	CA-CB-CG	-6.69	107.11	113.80
1	B	448	ARG	N-CA-C	6.26	118.18	111.36
1	B	320	PHE	CA-CB-CG	-5.97	107.83	113.80
1	A	31	SER	N-CA-C	5.63	119.17	111.39
1	B	260	ILE	N-CA-C	5.62	116.04	108.17
1	A	68	GLN	N-CA-C	-5.50	105.67	112.38
1	B	258	SER	N-CA-C	5.49	117.82	110.35
1	A	136	ASP	CA-CB-CG	5.43	118.03	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	69	PHE	CA-CB-CG	-5.35	108.45	113.80
1	A	22	ALA	CA-C-N	5.32	127.40	120.28
1	A	22	ALA	C-N-CA	5.32	127.40	120.28
1	A	23	LYS	N-CA-C	-5.30	105.50	111.28
1	A	211	ASP	CA-CB-CG	5.29	117.89	112.60
1	B	283	GLN	N-CA-C	-5.18	102.18	109.96
1	A	221	ILE	N-CA-C	5.16	115.37	110.42
1	B	382	ASN	CA-CB-CG	-5.12	107.48	112.60
1	B	442	ASN	OD1-CG-ND2	-5.11	117.49	122.60
1	B	459	TYR	N-CA-C	5.09	119.34	113.18
1	A	199	ASN	O-C-N	5.05	123.85	121.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2020	488	1991	38	0
1	B	2026	489	1993	41	0
2	A	65	0	90	19	0
2	B	39	0	54	17	0
3	A	1	0	0	1	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
5	A	252	504	0	7	0
5	B	296	592	0	4	0
All	All	4701	2073	4128	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:47:ALA:HB3	2:A:302:PG4:H52	1.51	0.92
1:A:206:HIS:HE1	2:A:306:PG4:H81	1.35	0.89
1:A:206:HIS:CE1	2:A:306:PG4:H81	2.07	0.89
1:B:282:SER:HA	5:B:798:HOH:O	1.75	0.84
1:B:413:LYS:HZ3	2:B:601:PG4:H82	1.43	0.81
1:B:413:LYS:HG2	2:B:601:PG4:H41	1.63	0.79
1:A:179:PHE:HD2	2:A:301:PG4:H31	1.48	0.78
1:B:413:LYS:HZ2	2:B:601:PG4:H11	1.50	0.76
1:B:325:LYS:HZ2	2:B:603:PG4:H42	1.53	0.74
1:B:486:TYR:HB3	2:B:603:PG4:H82	1.72	0.71
1:A:51:LYS:HD3	2:A:307:PG4:O3	1.92	0.70
1:A:179:PHE:HD2	2:A:301:PG4:C3	2.04	0.70
1:A:125:VAL:O	1:A:128:GLU:HG2	1.92	0.69
1:B:314:GLU:HG3	5:B:917:HOH:O	1.93	0.68
1:B:413:LYS:NZ	2:B:601:PG4:H82	2.09	0.67
1:A:51:LYS:HZ2	2:A:307:PG4:H21	1.59	0.66
1:B:435:PHE:O	1:B:447:MET:HE2	1.96	0.66
1:A:179:PHE:CD2	2:A:301:PG4:H31	2.30	0.65
1:B:282:SER:O	1:B:284:GLN:HG3	1.95	0.65
1:B:413:LYS:NZ	2:B:601:PG4:H11	2.11	0.64
1:B:413:LYS:HZ3	2:B:601:PG4:H61	1.63	0.63
1:A:47:ALA:CB	2:A:302:PG4:H52	2.27	0.62
5:A:538:HOH:O	1:B:499:LEU:HD21	2.01	0.61
2:A:306:PG4:H42	5:A:474:HOH:O	2.01	0.60
1:B:500:ASN:HA	5:B:707:HOH:O	2.02	0.60
1:A:117:HIS:O	1:A:121:GLU:HB2	2.01	0.60
1:A:134:GLU:O	1:A:135:ILE:HG13	2.01	0.60
1:A:51:LYS:HZ3	2:A:307:PG4:H42	1.68	0.58
1:B:313:ASN:HB3	1:B:316:GLN:HB2	1.87	0.56
1:B:491:VAL:HG11	1:B:499:LEU:HA	1.87	0.56
1:A:74:LYS:HZ3	2:A:305:PG4:H31	1.71	0.56
1:A:31:SER:HA	5:A:435:HOH:O	2.04	0.55
2:A:306:PG4:H41	5:A:440:HOH:O	2.06	0.54
1:A:51:LYS:NZ	2:A:307:PG4:H71	2.24	0.53
1:B:325:LYS:NZ	2:B:603:PG4:H42	2.22	0.53
1:A:207:GLN:O	2:A:306:PG4:H52	2.08	0.52
1:B:261:LYS:HE3	2:B:603:PG4:H51	1.91	0.52
1:B:436:LEU:HD23	1:B:447:MET:HE3	1.92	0.52
1:B:350:ASP:OD1	1:B:421:GLU:HG3	2.09	0.51
1:B:413:LYS:HD3	2:B:601:PG4:H11	1.92	0.51
1:B:436:LEU:HD23	1:B:447:MET:CE	2.42	0.49
1:A:30:THR:H	2:A:306:PG4:H21	1.78	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:413:LYS:NZ	2:B:601:PG4:H61	2.27	0.49
1:A:129:LEU:HB3	1:A:142:ASN:HB3	1.94	0.49
1:A:189:ASP:OD1	1:A:201:LYS:HE3	2.14	0.48
1:B:383:GLY:HA3	1:B:390:LYS:HE2	1.94	0.48
1:A:87:LEU:HD21	1:A:115:ILE:CG2	2.44	0.48
1:A:13:LYS:H	1:B:257:LYS:HD2	1.78	0.47
1:B:397:VAL:HG12	1:B:411:TYR:CE1	2.50	0.47
1:A:51:LYS:NZ	2:A:307:PG4:H21	2.27	0.47
1:B:433:LYS:O	1:B:437:SER:HB3	2.14	0.47
1:B:280:TYR:CE1	1:B:458:GLN:HG3	2.50	0.46
1:A:179:PHE:CD1	1:A:179:PHE:C	2.94	0.46
1:A:182:LYS:O	1:A:186:SER:HB3	2.16	0.45
1:B:453:ASN:HB3	1:B:456:GLU:HG3	1.99	0.45
1:A:24:LYS:HD2	5:A:630:HOH:O	2.15	0.45
1:A:189:ASP:OD1	1:A:201:LYS:CE	2.65	0.45
1:B:261:LYS:HE3	2:B:603:PG4:O4	2.15	0.45
1:A:152:ASP:OD2	1:A:155:LYS:HD3	2.17	0.44
1:B:259:ASP:O	2:B:603:PG4:H72	2.16	0.44
1:A:62:ASN:HB3	1:A:65:GLN:HB2	1.99	0.44
1:A:90:ARG:NH1	1:A:122:THR:OG1	2.51	0.44
1:A:19:ASN:OD1	1:A:51:LYS:HE3	2.18	0.44
1:A:141:TYR:CD2	1:A:141:TYR:N	2.85	0.44
1:B:464:LYS:NZ	2:B:604:PG4:O3	2.51	0.43
1:A:36:ASN:HB2	1:A:41:ALA:HB3	1.99	0.43
1:B:397:VAL:CG1	1:B:411:TYR:CE1	3.03	0.42
1:B:473:ILE:HD13	1:B:473:ILE:HA	1.79	0.42
1:A:155:LYS:HG2	1:A:159:GLU:OE1	2.19	0.42
1:B:290:ASN:O	1:B:291:ASP:HB2	2.20	0.42
1:A:51:LYS:HD3	2:A:307:PG4:C4	2.50	0.41
1:A:179:PHE:HB2	2:A:301:PG4:H21	2.03	0.41
1:A:249:ASN:O	1:A:250:LYS:C	2.63	0.41
1:B:413:LYS:CD	2:B:601:PG4:H11	2.50	0.41
1:B:499:LEU:HD11	5:B:747:HOH:O	2.21	0.41
5:A:485:HOH:O	1:B:317:LYS:HE3	2.20	0.41
1:B:259:ASP:OD1	2:B:603:PG4:H61	2.21	0.41
1:A:237:LYS:HD2	3:A:303:CL:CL	2.59	0.40
5:A:538:HOH:O	1:B:499:LEU:HD11	2.21	0.40
1:B:374:GLY:O	1:B:375:ALA:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/254 (98%)	235 (95%)	12 (5%)	1 (0%)	30	22
1	B	249/254 (98%)	239 (96%)	7 (3%)	3 (1%)	10	4
All	All	497/508 (98%)	474 (95%)	19 (4%)	4 (1%)	16	8

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	288	LYS
1	B	375	ALA
1	A	39	ASN
1	B	292	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/218 (99%)	210 (97%)	6 (3%)	38	35
1	B	216/218 (99%)	207 (96%)	9 (4%)	26	20
All	All	432/436 (99%)	417 (96%)	15 (4%)	32	27

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ASP
1	A	19	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	31	SER
1	A	36	ASN
1	A	136	ASP
1	A	150	ASP
1	B	257	LYS
1	B	258	SER
1	B	270	ASN
1	B	282	SER
1	B	291	ASP
1	B	314	GLU
1	B	425	SER
1	B	478	LYS
1	B	493	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	108	HIS
1	A	206	HIS
1	A	207	GLN
1	B	266	ASN
1	B	359	HIS
1	B	442	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 11 ligands modelled in this entry, 3 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PG4	B	603	-	12,12,12	0.40	0	11,11,11	0.27	0
2	PG4	A	306	-	12,12,12	0.57	0	11,11,11	0.33	0
2	PG4	A	307	-	12,12,12	0.38	0	11,11,11	0.67	0
2	PG4	A	305	-	12,12,12	0.96	1 (8%)	11,11,11	0.91	0
2	PG4	A	301	-	12,12,12	0.27	0	11,11,11	0.27	0
2	PG4	B	604	-	12,12,12	0.39	0	11,11,11	0.63	0
2	PG4	A	302	-	12,12,12	0.84	0	11,11,11	0.58	0
2	PG4	B	601	-	12,12,12	0.16	0	11,11,11	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PG4	B	603	-	-	3/10/10/10	-
2	PG4	A	306	-	-	1/10/10/10	-
2	PG4	A	307	-	-	3/10/10/10	-
2	PG4	A	305	-	-	4/10/10/10	-
2	PG4	A	301	-	-	3/10/10/10	-
2	PG4	B	604	-	-	3/10/10/10	-
2	PG4	A	302	-	-	4/10/10/10	-
2	PG4	B	601	-	-	3/10/10/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	305	PG4	C6-C5	2.19	1.60	1.49

There are no bond angle outliers.

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	603	PG4	C6-C5-O3-C4
2	B	604	PG4	C5-C6-O4-C7
2	B	603	PG4	O2-C3-C4-O3
2	A	305	PG4	O2-C3-C4-O3
2	B	601	PG4	O1-C1-C2-O2
2	A	305	PG4	O3-C5-C6-O4
2	A	302	PG4	C6-C5-O3-C4
2	A	307	PG4	O2-C3-C4-O3
2	A	307	PG4	O1-C1-C2-O2
2	B	604	PG4	C8-C7-O4-C6
2	A	307	PG4	C6-C5-O3-C4
2	B	603	PG4	O4-C7-C8-O5
2	A	302	PG4	O4-C7-C8-O5
2	A	301	PG4	C1-C2-O2-C3
2	A	305	PG4	O4-C7-C8-O5
2	A	302	PG4	C4-C3-O2-C2
2	A	301	PG4	C6-C5-O3-C4
2	B	604	PG4	C3-C4-O3-C5
2	A	302	PG4	O3-C5-C6-O4
2	A	305	PG4	C8-C7-O4-C6
2	A	301	PG4	C4-C3-O2-C2
2	A	306	PG4	O3-C5-C6-O4
2	B	601	PG4	O2-C3-C4-O3
2	B	601	PG4	C6-C5-O3-C4

There are no ring outliers.

8 monomers are involved in 36 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	603	PG4	7	0
2	A	306	PG4	6	0
2	A	307	PG4	6	0
2	A	305	PG4	1	0
2	A	301	PG4	4	0
2	B	604	PG4	1	0
2	A	302	PG4	2	0
2	B	601	PG4	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/254 (98%)	0.43	29 (11%) 9 8	16, 27, 67, 100	11 (4%)
1	B	250/254 (98%)	0.09	15 (6%) 27 27	17, 24, 47, 86	9 (3%)
All	All	500/508 (98%)	0.26	44 (8%) 15 15	16, 26, 57, 100	20 (4%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	499	LEU	5.8
1	A	122	THR	5.7
1	B	287	ASN	5.7
1	B	289	LYS	5.1
1	A	120	LEU	4.8
1	A	248	LEU	4.7
1	A	36	ASN	4.6
1	B	290	ASN	4.5
1	B	292	ALA	4.5
1	A	123	GLY	4.3
1	A	39	ASN	4.3
1	A	41	ALA	4.2
1	A	124	ALA	4.2
1	A	249	ASN	4.1
1	A	250	LYS	4.0
1	B	291	ASP	3.9
1	B	293	TRP	3.9
1	B	500	ASN	3.9
1	A	139	LYS	3.6
1	B	252	ALA	3.5
1	B	497	LYS	3.1
1	A	125	VAL	3.1
1	A	121	GLU	2.9
1	B	286	SER	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	501	LYS	2.9
1	B	496	SER	2.7
1	A	119	LEU	2.6
1	A	151	LYS	2.5
1	A	40	ASP	2.5
1	B	251	ALA	2.5
1	A	88	VAL	2.4
1	A	90	ARG	2.4
1	A	138	LYS	2.4
1	A	42	TRP	2.4
1	A	135	ILE	2.3
1	A	126	LYS	2.3
1	A	85	CYS	2.2
1	A	245	SER	2.1
1	A	37	LYS	2.1
1	A	149	LEU	2.1
1	A	136	ASP	2.1
1	A	35	SER	2.1
1	B	336	CYS	2.0
1	A	137	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PG4	A	307	13/13	0.68	0.20	68,80,85,86	0
2	PG4	A	305	13/13	0.75	0.27	45,75,100,101	0
2	PG4	A	306	13/13	0.77	0.30	65,82,87,92	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PG4	B	603	13/13	0.77	0.18	37,62,80,82	0
2	PG4	B	604	13/13	0.77	0.29	108,116,117,120	0
2	PG4	B	601	13/13	0.86	0.18	56,70,78,80	0
2	PG4	A	301	13/13	0.90	0.14	35,52,66,72	0
2	PG4	A	302	13/13	0.90	0.11	30,35,45,46	0
4	NA	A	304	1/1	0.93	0.21	22,22,22,22	1
3	CL	A	303	1/1	0.98	0.07	26,26,26,26	0
3	CL	B	602	1/1	0.99	0.07	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.